

## MaineCare PDL

### PDL Effective January 1, 2026

\* PLEASE NOTE: For a **search** box hit Ctrl F

\* PLEASE NOTE: All **cost effective** generics applicable to DEL are considered **PREFERRED Drugs**. "BASIC" Covered Drugs are **bolded** with the Coverage Indicator of "MC / DEL".

General Criteria for all PDL categories- For more information or help using the PDL, providers may call 1-888-445-0497; members should call 1-866-796-2463. To access PDL and PA materials via the internet: [www.maineicarepdl.org](http://www.maineicarepdl.org)

A: Preferred Drugs- Unless otherwise specified, preferred drugs are available without prior authorization. Step order may apply for preferred drugs in some drug categories as indicated on the PDL. (See item "D" below for explanation of step order.)

B: Requests for Non-preferred Drugs- Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

C: Adequate Drug Trials- 1. The minimum trial period for each preferred and step order drug is two weeks, unless otherwise stated within specific PDL drug categories; trials with less than a two week duration will be reviewed on a case-by-case basis; 2. A trial will not be considered valid if preferred or non-preferred products were readily available (by override, individual purchase, samples, etc.); 3. Certain drug trials, such as with controlled substances, may require evidence that the preferred drugs were actually tried (example: with random pill counts and with random urine drug tests, using the methods of GC/MS with no lower threshold); 4. Adequate trials require documentation of attempts to titrate dose of preferred agents toward desired clinical response. 5. Adequate trials include prevention/treatment of common adverse effects associated with preferred agents (example: antinausea, antipruritic, etc.)

D: Step Order- When numbers appear in the "step order" column, it means drugs in this category must be used in the order specified, with the lower numbers having preference over the higher numbers. Chart notes should be provided to confirm drug trials that do not appear in the member's MaineCare drug profile.

E: The Department will institute strategies to ensure cost effectiveness through the use of an enhanced Drug Benefit. Preferred brand drugs will no longer be preferred in any PDL drug category where preferred generic drugs are also available. It is expected that preferred generics will be used prior to any preferred brands. This will be operated as a form of step care. Preferred brands in these categories will require prior authorization for these high utilization / high cost members.

F: Brand Name Medication Requests- (Must be submitted on the Brand Name PA request form)- According to MaineCare Benefits Manual Chapter II (80.07-5), when medically necessary covered brand-name drugs have an A-rated generic equivalent available, the most cost effective medically necessary version will be approved and reimbursed, since the brand-name and A-rated generic drugs have been determined by the FDA to be chemically and therapeutically equivalent. The Bureau does not make determinations as to whether or not a generic drug is clinically inferior or inequivalent to its brand version. This is the proper role of the FDA. Physicians should submit their reports of generic inequivalence directly to the FDA via the MEDWATCH.

G: PA requests for non- FDA Approved Indications- Decisions will be made on a case-by-case basis until the DUR committee is able to review the evidence and make a recommendation. Interim approvals and DUR recommendations for approval of a drug for a non- FDA approved indication will require a minimum of two published, peer reviewed, non contradicted, double blind, placebo-controlled randomized clinical studies establishing both safety and efficacy.

H: Dose Consolidation Requirements- Some drugs may also be affected by dose consolidation requirements. Please see Dose Consolidation List and/or Splitting Tables provided in the PDL.

I: Trials from Multiple Drug Classes - Trial/failure/intolerance to preferred agents from multiple classes within the same category or other categories of drugs may be required prior to the approval of non-preferred agents (e.g., Cymbalta, Zofran and others).

J: Drug-specific PA Forms- Drug-specific PA forms contain medical necessity documentation requirements and/or criteria that may not be repeated in the PDL. Drug-specific PA forms may be obtained on the web at [www.maineicarepdl.org](http://www.maineicarepdl.org).

K: PA Exemptions for Prescribers- According to MaineCare Benefits Manual Chapter II (80.07-4), providers may receive a three (3) month exemption from prior authorization requirement for certain categories of drugs when they demonstrate high compliance with the Department's PDL. The Department will notify providers in writing which drug categories are included and what dates apply to the exemption. If a provider loses his/ her exemption, members who previously were not required to obtain a PA while the prescriber was exempt will be required to do so, and criteria for approval of that medication will need to be met.

L: Drug-Drug Interactions (DDI)- The DUR Committee has implemented new drug-drug interaction edits requiring prior authorization. Several drug-drug combinations and PDL drug categories are affected by new PA requirements. These will be indicated in the PDL with DDI notation. Please see the DDI document provided in the PDL.





Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
ANTI INFECTIVE COMBO'S - MISC.	MC/DEL	ERYTHROMYCIN/SULF SUSR	MC		BACTRIM DS TABS	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL	SEPTRA/DS TABS	MC		VABOMERE <sup>1</sup>		
	MC/DEL	SULFAMETHOXAZOLE/TRIMETH					
	MC/DEL	TRIMETHOPRIM/SULFAMETHOXA					
<b>ANTIPROTOZOALS</b>							
	MC/DEL	BENZNIDAZOLE <sup>2</sup>	MC	8	ALINIA <sup>1</sup>	<a href="#">Use PA Form# 20420</a>	<b>Benznidazole</b> is indicated for pediatric patients 2 to 12 years of age for the treatment of Chagas disease (American trypanosomiasis) caused by trypanosoma cruzi.
	MC/DEL	LAMPIT <sup>2</sup>					
<b>ANTI - FUNGALS</b>							
ANTIFUNGALS - ASSORTED	MC/DEL	FLUCONAZOLE <sup>1</sup>	MC/DEL	6	LAMISIL TABS <sup>4</sup>	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. The other criteria are listed on the Antifungal PA form including the required proof of a non-cosmetic fungal infection.
	MC/DEL	KETOCONAZOLE TABS <sup>7</sup>	MC/DEL	6	ITRACONAZOLE	See Quantity Limit table.	
	MC/DEL	NYSTATIN	MC	8	BREXAFEMME	Non-preferred products must be used in specified step order.	
	MC/DEL	TERBINAFINE TABS <sup>4</sup>	MC/DEL	8	CRESEMB <sup>9</sup>	Continue to use Anti-Fungal PA form for non-preferred products.	
	MC/DEL	VORICONAZOLE TABS	MC/DEL	8	GRIFULVIN V TABS		
			MC	8	GRISEOFULVIN SUSP		
			MC	8	GRISEOFULVIN ULTRAMICROSI TABS <sup>8</sup>		
			MC	8	GRIS-PEG TABS	1. QL-1/every 7-day period (150mg only).	
			MC	8	REZZAYO <sup>9</sup>	2. <b>Sporanox</b> QL 300cc/month with PA. See Quantity Limit table.	
			MC/DEL	8	SPORANOX SOLN <sup>2</sup>	3. <b>Sporanox</b> QL 30/month with PA.	
			MC/DEL	8	SPORANOX PULSEPAK CAPS <sup>3</sup>	4. Quantity limit of one tablet daily. Please see Dosage Consolidation List.	
			MC/DEL	8	SPORANOX CAPS <sup>3</sup>		
			MC/DEL	8	DIFLUCAN	5. Approved if immuno suppressed/ HIV or if the member has failed a 7-day trial of a preferred antifungal therapy.	
			MC/DEL	8	ERAXIS INJ <sup>6</sup>	6. <b>Eraxis</b> will be approved if submitting with documentation that it was initiated during a hospitalization and this request is to finish the hospital course.	
			MC	8	GRIFULVIN SUSP	7. Quantity limits allowing 30-day supply without PA. PA will be required if using > 30 days.	
			MC/DEL	8	ONMEL		
			MC/DEL	8	NOXAFL <sup>5</sup>		
			MC/DEL	8	TOLSURA		
			MC/DEL	8	VFEND TABS		
			MC	8	VIVJOA		
<b>ANTI - VIRALS</b>							
ANTIRETROVIRALS - PREP	MC	APRETUDE	MC		TRUVADA <sup>1</sup>	<a href="#">Use PA Form# 20420</a>	DDI: The concomitant use of the following drugs with Descovy is not recommended: tipranavir/ritonavir, St. John's wort, and the antimycobacterials rifabutin, rifampin, or rifapentine.
	MC	DESCOY <sup>1</sup>				1. Quantity limit of one per day	
	MC	EMTRICITABINE-TENOFOVIR					
	MC	DISOP (ORAL) TAB					
	MC	YEZTUGO					
ANTIRETROVIRALS	MC/DEL	ABACAVIR TABS	MC/DEL	8	ABACAVIR SOL	<a href="#">Use PA Form# 20420</a>	<b>Fuzeon</b> : Prescriber is either an HIV specialist provider or has consulted with one. Documentation of genotype testing is supplied and shows that there is no other potent, appropriate two or three drug oral regimen available, AND patient has a positive HIV viral load within past 6 months while on his/her current antiretroviral regimen AND the drug will be prescribed with at least two other drugs that are likely to be active based on the genotype testing.
	MC/DEL	ATAZANAVIR	MC/DEL	8	APTVUS	1. Quantity limit of one per day	DDI: Reyataz requires prior authorization if it is currently being used in combination with either prevacid, pantoprazole, prilosec, or any currently non preferred PPI .
	MC	BIKTARVY	MC	8	ATRIPLA <sup>1</sup>	2. Only preferred if <b>Norvir</b> script is in member's profile within the past 30 days of filling <b>Prezista</b>	DDI: Norvir requires prior authorization if it is currently being used in combination with either enablex 15mg or vesicare 10mg.
	MC	CABENUVA	MC/DEL	8	CIMDUO	3. <b>Isentress Chewable</b> will only be approved if between the age of 2-12 years old	DDI: Preferred Crixivan caps requires prior authorization if it is currently being used in combination with either enablex 15mg or vesicare 10mg.
	MC	COMPLERA <sup>1</sup>	MC/DEL	8	COMBIVIR TABS	4. Clinical PA required	DDI: Administration with the following drugs: the anticonvulsants carbamazepine, oxcarbazepine, phenobarbital, and phenytoin; the antimycobacterials rifampin and rifapentine; proton pump inhibitors such as dexlansoprazole, esomeprazole, lansoprazole, omeprazole, pantoprazole, rabeprazole; systemic dexamethasone (more than a single dose); and St. John's wort with Odefsey is contraindicated.
	MC/DEL	DELSTRIGO	MC/DEL	8	EDURANT	5. Only preferred for post- exposure prophylaxis	<b>Stribild</b> : PA required; must provider rationale as to why the member's medical need cannot be met with preferred agents, particularly genvoya or combinations of preferred and agents AND must be antiretroviral treatment-naïve or virologically controlled on current therapy (HIV-1RNA < copies/ml) AND be HBV negative AND not be combined with other anti-retroviral agents.
	MC	DIDANOSINE	MC/DEL	8	EPZICOM <sup>1</sup>		DDI: Tivicay will require prior authorization is used with nevirapine, oxcarbazepine, phenytoin, phenobarbital, carbamazepine, and St. John's wort.
	MC/DEL	DOVATO	MC/DEL	8	FUZEON		DDI: Aatazanavir or Darunavir and the following drugs are contraindicated (due to potential for serious and/or life-threatening events or loss of therapeutic effect):
	MC	EFAVIRENZ TAB	MC/DEL	8	INTELENCE		alfuzosin, dronedarone, rifampin, irinotecan, ergotamine, methylergonovine, cisapride, St. John's wort, lovastatin, simvastatin, pimozide, nevirapine, sildenafil (when given as revatio for treatment of PAH), indinavir, triazolam, or PO midazolam will be non-preferred and require prior authorization if it is currently being used in combination with
	MC/DEL	EFAVIRENZ CAP	MC/DEL	8	ISENTRESS <sup>3</sup>		
	MC	EFAVIRENZ-EMTRICITABINE-TENOFOVIR DF TAB	MC/DEL	8	ISENTRESS HD		
	MC	EMTRIVA <sup>1</sup>	MC	8	JULUCA		
			MC	8	KALETRA		

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
	MC	EPIVIR SOL	MC/DEL	8	LAMIVUDINE SOLN		tyost. <b>DDI:</b> Combined P-gp, UGT1A1 and strong CYP3A inhibitors may significantly increase plasma concentrations of Sunlenca. Concomitant administration of Sunlenca with these inhibitors is not recommended. <b>Sunlenca:</b> In combination with other antiretroviral(s) for the treatment of HIV-1 infection in heavily treatment-experienced adults with multidrug resistant HIV-1 infection failing their current antiretroviral regimen due to resistance, intolerance, or safety considerations.
	MC/DEL	EVOTAZ <sup>1</sup>	MC/DEL	8	LEXIVA		
	MC	GENVOYA <sup>1,4</sup>	MC/DEL	8	NEVIRAPINE		
	MC/DEL	ISENTRESS 400MG <sup>5</sup>	MC	8	NORVIR		
	MC/DEL	ISENTRESS CHEW <sup>3</sup>	MC/DEL	8	PIFELTRO		
	MC/DEL	ISENTRESS POWDER	MC	8	RETROVIR		
	MC/DEL	LAMIVUDINE TABS	MC	8	REYATAZ		
	MC/DEL	LAMIVUDINE/ZIDOVUDINE	MC/DEL	8	SELZENTRY		
	MC/DEL	LOPINAVIR-RITONAVIR SOL	MC	8	STAVUDINE		
	MC	LOPINAVIR-RITONAVIR TAB	MC	8	STRIBILD <sup>1</sup>		
	MC	ODEFSEY <sup>1</sup>	MC/DEL	8	SYMF <sup>4</sup>		
	MC/DEL	PREZCOBIX	MC/DEL	8	SYMF LO <sup>4</sup>		
	MC	PREZISTA <sup>2</sup>	MC/DEL	8	SYMTUZA		
	MC/DEL	RITONAVIR TAB 100MG	MC/DEL	8	TRIZIVIR TABS		
	MC	RUKOBIA <sup>4</sup>	MC/DEL	8	VIRACEPT TABS		
	MC	SUNLENCA <sup>4</sup>	MC	8	VITEKTA		
	MC	SUSTIVA <sup>1</sup>	MC	8	ZERIT		
	MC	TIVICAY	MC	8	VIDEX EC		
	MC	TIVICAY PD	MC	8	VIREAD TABS <sup>1</sup>		
	MC	TRIUMEQ <sup>1</sup>	MC/DEL	8	ZIAGEN TABS		
	MC	TROGARZO <sup>4</sup>	MC/DEL	8	ZIAGEN SOL		
	MC	TYBOST	MC/DEL	9	VIRAMUNE XR		
	MC	VIREAD POW					
	MC/DEL	ZIDOVUDINE					
CYTO-MEGALOVIRUS AGENTS	MC	CIDOFOVIR	MC	4	VALCYTE TABS	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC	FOSCARNET SODIUM	MC	8	FOSCAVIR		
	MC/DEL	GANCICLOVIR	MC/DEL	8	LIVTENCY <sup>1</sup>		
	MC/DEL	VALGANCICLOVIR	MC	8	PREVYMIS		
HERPES AGENTS	MC/DEL	VALACYCLOVIR HCL	MC/DEL	8	FAMCICLOVIR <sup>1</sup>	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
			MC	8	SITAVIG		
			MC	8	VALTREX TABS <sup>1</sup>		
			MC/DEL	9	FAMVIR TABS <sup>1</sup>		
INFLUENZA AGENTS	MC	AMANTADINE CAPS	MC		AMANTADINE TABS	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC	RELENZA DISKHALER AEPB	MC		FLUMADINE TABS		
	MC/DEL	OSELTAMIVIR <sup>1</sup>	MC		FLUMIST		
			MC/DEL		RIMANTADINE HCL TABS		
			MC/DEL		TAMIFLU <sup>1</sup>		
			MC/DEL		TAMIFLU SUS		
			MC/DEL		XOFLUZA		
<b>IMMUNE SERUMS</b>							
IMMUNE SERUMS	MC	HYPERRHO INJ					
<b>HEPATITIS AGENTS</b>							
HEPATITIS C AGENTS	MC	SOFSBUVIR/VELPATASVIR <sup>3</sup> (Authorized generic labeler 72626 Aseuga Therapeutics)	MC/DEL		COPEGUS TABS	<a href="#">Use PA Form #10700</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC	MAVYRET <sup>3</sup>	MC/DEL		DAKLINZA		
	MC/DEL	PEGASYS KIT <sup>1</sup>	MC		EPCLUSA <sup>2</sup>		
	MC/DEL	PEGASYS SOLN	MC		HARVONI <sup>2</sup>		
	MC/DEL	PEG-INTRON KIT <sup>1</sup>	MC/DEL		REBETOL CAPS		
	MC	RIBAVIRIN	MC		RIBAPAK		
	MC/DEL	RIBASPHERE	MC		SOVALDI <sup>2</sup>		
			MC		VIEKIRA PAK <sup>2</sup>		
			MC		VIEKIRA XR <sup>2</sup>		
			MC		VOSEVI		
			MC/DEL		ZEPATIER <sup>2</sup>		
HEPATITIS AGENTS - MISC.			MC		ACTIMMUNE	<a href="#">Use PA Form# 20420</a>	Approved for chronic granulomatous disease, osteopetrosis and idiopathic pulmonary fibrosis.



Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
					receptor (AChR) antibody positive. 6. For the treatment of patients between ages 4-17 years of age.	<b>Migraine:</b> Consideration for <b>Botox</b> approvals will only be made after failures of required trials of the following preferred medications: tricyclic or venlafaxine, beta blocker, valproic acid, topiramate.  <b>Firdapse</b> is recommended for the treatment of Lambert-Eaton myasthenic syndrome (LEMS) in adults.	
NEUROLOGICS- hATTR AGENTS			MC MC MC MC MC/DEL MC/DEL MC/DEL	8 8 8 8 8 8 8	AMVUTTRA <sup>1</sup> ATTRUBY ONPATRO <sup>1</sup> TEGSEDI <sup>1</sup> VYNDAMAX <sup>1</sup> VYNDAQEL <sup>1</sup> WAINUA <sup>1</sup>	<a href="#">Use PA Form# 20420</a> 1. PA required for appropriate diagnosis.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval.  <b>Tegsedi</b> should be non-preferred and approved for patients for whom other treatments, including Onpatro, have been ineffective. <b>Vyndamax</b> will be considered for the treatment of the cardiomyopathy of wild type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) in adults to reduce cardiovascular mortality and cardiovascular-related hospitalization.
NEUROLOGICS- SMA		GENE			GENE	<a href="#">Use PA Form# 20420</a> 1. Clinical PA is required to establish diagnosis and medical necessity. 2. For patients 2 months of age and older.	<b>Zolgensma:</b> The patient is less than 2 years of age AND The diagnosis is spinal muscular atrophy (SMA) AND the patient has bi-allelic mutations of the SMN1 gene AND the patient does not have advanced SMA (e.g. complete paralysis of limbs or permanent ventilator dependence) AND medication is prescribed per the dosing.  <b>Spinraza:</b> The diagnosis is spinal muscular atrophy (SMA) type 1, 2, or 3 (results of genetic testing must be submitted) AND The patient has at least 2 copies of the SMN2 gene AND The prescriber is a neurologist, pulmonologist, or other physician with expertise in treating SMA AND Baseline motor ability has been established using one of the following exams: Hammersmith Infant Neurological Exam (HINE) Hammersmith Functional Motor Scale Expanded (HFMSE) Upper Limb Module Test (non-ambulatory) Children's Hospital of Philadelphia Infant Test of Neuromuscular Disorders (CHOP INTEND) AND Prior to starting therapy, and prior to each dose, the following laboratory tests will be conducted: Treating provider attests the member has a platelet count > 50,000/ml or greater Treating provider agrees to do platelet count and coagulation test before each dose Treating provider agrees to do a quantitative spot urine protein test before each dose  Concomitant use of <b>Spinraza</b> and <b>Zolgensma</b> is investigational and will not be approved AND Use of Spinraza after gene replacement therapy, including <b>Zolgensma</b> is investigational and will not be approved  <b>Note:</b> Initial approval will be granted for 4 loading doses (the first 3 loading doses should be administered at 14-day intervals; the 4th loading dose should be administered 30 days after the 3rd dose). Renewal may be granted for up to 12 months with a maximum of 3 doses approved per year (12mg (5ml) every 4 months). For therapy continuation, clinical documentation must be submitted documenting improvement or maintenance of motor ability OR slower progression of disease than would otherwise be expected.
NEUROLOGICS- RETT SYNDROME			MC		DAYBUE <sup>1,2</sup>	<a href="#">Use PA Form# 20420</a> 1. Clinical PA required for appropriate diagnosis 2. For the treatment of patients 2 years of age and older.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ALS DRUGS	MC/DEL	RILUZOLE	MC MC MC MC MC MC		EXSERVAN QALSODY RILUTEK TABS RADICAVA <sup>1</sup> RELYVRI <sup>1</sup> TIGLUTIK	<a href="#">Use PA Form# 20420</a> 1. Clinical PA for indication required.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  <b>Qalsody:</b> For the treatment of amyotrophic lateral sclerosis (ALS) in adults who have a mutation in the superoxide dismutase 1 (SOD1) gene. Continued approval for this indication may be contingent upon verification of clinical benefit in confirmatory trial(s).
MOVEMENT DISORDERS	MC MC MC MC	AUSTEDO <sup>1</sup> AUSTEDO XR <sup>1</sup> INGREZZA <sup>1</sup> TETRABENAZINE <sup>1</sup>	MC/DEL		XENAZINE	<a href="#">Use PA Form# 20420</a> 1. Clinical PA required for appropriate diagnosis	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  <b>DDI:</b> Avoid concomitant use of <b>VMAT2 inhibitors</b> with MAO inhibitors (e.g. isocarboxazid, phenelzine, or selegiline). Concomitant use with strong CYP3A4 inducers (e.g. rifampin, carbamazepine, phenytoin, St. John's wort) is not recommended



Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
			MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC MC/DEL MC/DEL		METHITEST TAB METHYLTESTOSTERONE CAP OXANDROLONE STRIANT MUC ER TESTIM TESTOSTERONE GEL PACKETS TESTOSTERONE SOL TESTRED CAPS TLANDO VOGELXO XYOSTED		
ESTROGENS - PATCHES / TOPICAL	MC MC/DEL MC/DEL	EVAMIST MINIVELLE PATCH VIVELLE-DOT PTTW	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	5 8 8 8 8	ESTRADIOL PTWK DIVIGEL <sup>1</sup> CLIMARA PTWK ELESTRIN <sup>1</sup> MENOSTAR PATCH	<a href="#">Use PA Form# 20420</a> 1. Step order drugs must be used in specified step order.	Approved for failures on multiple oral estrogen agents after 90 day trials or if unable to swallow any oral medication.
ESTROGENS - TABS	MC/DEL MC/DEL	ESTRADIOL PREMARIN TABS	MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC		ENJUVIA ESTRADIOL-NORETHINDRONE ESTRACE TABS ESTRATAB TABS MENEST TABS NORETHINDRON-ETHINYL ORTHO-EST TABS	<a href="#">Use PA Form# 20420</a> Must fail preferred products before non-preferred products.	Preferred drugs must be tried for at least 90 days and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ESTROGEN COMBO'S	MC/DEL MC/DEL MC/DEL MC/DEL	ANGELIQ COMBIPATCH PTTW PREMPHASE TABS PREMPRO TABS	MC/DEL MC/DEL MC MC/DEL MC/DEL		FEMHRT 1/5 TABS <sup>1</sup> FYAVOLV LOPREEZA TAB ORTHO-PREFEST TABS <sup>1</sup> SYNTEST H.S. TABS <sup>1</sup>	<a href="#">Use PA Form# 20420</a> 1. Must fail Premphase and Prempro products before non preferred products.	Preferred drugs must be tried for at least 90 days and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
PROGESTINS	MC/DEL MC/DEL MC MC	MEDROXYPROGESTERONE ACETA <sup>1</sup> NORETHINDRONE ACETATE TABS <sup>1</sup> 17-ALPH HYDROXYPROGESTERONE PWDR PROGESTERONE CAPS	MC/DEL MC MC MC/DEL MC/DEL		AYGESTIN TABS CYCRIN TABS PROGESTERONE POWD PROMETRIUM CAPS PROVERA TABS	<a href="#">Use PA Form# 20420</a> 1. Must fail Medroxyprogesterone and Norethindrone products before non-preferred products.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
<b>ENDOMETROSIS</b>							
CENTRAL PRECOCIOUS PUBERTY AGENTS	MC	FENSOLVI <sup>1</sup>				<a href="#">Use PA Form# 20420</a> 1. For pediatric patients 2 years of age and older with central precocious puberty (CPP).	
ENDOMETROSIS- NASAL	MC/DEL	SYNAREL (NASAL) SPRAY				<a href="#">Use PA Form# 20420</a>	Synarel is also indicated for central precocious puberty.
ENDOMETROSIS/ UTERINE FIBROIDS- ORAL	MC MC MC/DEL	MYFEMBREE <sup>1,2</sup> ORIAHNN <sup>1</sup> ORILISSA <sup>1</sup>				<a href="#">Use PA Form# 20420</a> 1. Prior treatment of NSAID and hormonal contraceptives required. 2. Limited to 24 months due to the risk of continued bone loss, which may not be reversible.	
ENDOMETROSIS- INJECTABLE	MC/DEL	DEPO-SUBQ PROVERA 104				<a href="#">Use PA Form# 20420</a>	
<b>CONTRACEPTIVES</b>							
CONTRACEPTIVES - PROGESTIN ONLY	MC/DEL MC/DEL MC MC MC/DEL MC/DEL	CAMILA TABS ERRIN INCASSIA TAB HEATHER TAB NORETHINDRONE ACETATE 0.35MG TABS SLYND	MC/DEL MC/DEL MC		JOLIVETTE NORA-BE TABS ORTHO MICRONOR TABS	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  If member experienced adverse reactions, consider using Oral Contraceptives from other groups. DDI: Preferred Oral Contraceptives will now be non-preferred and require prior authorization if it is currently being used in combination with Tracleer.
CONTRACEPTIVES - INJECTABLE	MC/DEL	MEDROXYPROGESTERONE ACETATE 150mg IM	MC/DEL		DEPO-PROVERA 150 mg SUSP	<a href="#">Use PA Form# 20420</a>	The preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.





Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
DIABETIC - DPP- 4 ENZYME INHIBITOR	MC/DEL MC/DEL	JANUVIA <sup>1,2</sup> TRADJENTA <sup>2</sup>	MC MC/DEL MC/DEL MC		BRYNOVIN NESINA QTERN ZITUVIO	<a href="#">Use PA Form# 20420</a>  1. Preferred if therapeutic doses of metformin are seen in members drug profile for at least 60 days within the past 18 months or if phosphate binder is currently seen in the members drug profile. 2. Dosing limits apply. Please refer to Dose Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  <b>BRYNOVIN:</b> In addition to tried and failed Preferred Agents, Brynovin requires tried and failed Non-Preferred Agent Zituvio.
DIABETIC - DPP- 4 ENZYME INHIBITOR-COMBO	MC/DEL MC/DEL MC/DEL	JANUMET <sup>1,2</sup> JANUMET XR <sup>1,2</sup> JENTADUETO <sup>1</sup>	MC/DEL MC/DEL MC MC/DEL MC MC		JENTADUETO XR KAZANO KOMBIGLYZE XR OSENI ZITUVIMET ZITUVIMET XR	<a href="#">Use PA Form# 20420</a>  1. Preferred if therapeutic doses of metformin are seen in members drug profile for at least 60 days within the past 18 months or if phosphate binder is currently seen in the members drug profile. 2. Dosing limits apply. Please refer to Dose consolidation list.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  <b>Zituvimet/ Zituvimet XR:</b> Approvals will require trial of preferred sitagliptin/metformin products or other preferred diabetic agents.
DIABETIC - LANCET-LANCET DEVICE						<a href="#">Use PA Form# 20420</a>	Please refer to the MaineCare Preferred Diabetic Supply List available at <a href="http://www.maineicarepdi.org">www.maineicarepdi.org</a>
DIABETIC - SYRINGES-NEEDLES						<a href="#">Use PA Form# 20420</a>	Please refer to the MaineCare Preferred Diabetic Supply List available at <a href="http://www.maineicarepdi.org">www.maineicarepdi.org</a>
DIABETIC - OTHER			MC		SYMLIN	<a href="#">Use PA Form #20420</a>	
SGLT 2 INHIBITORS	MC/DEL MC/DEL	FARXIGA JARDIANCE	MC/DEL MC/DEL		INVOKANA <sup>1</sup> STEGLATRO	<a href="#">Use PA Form# 20420</a>  1. Dosing limits apply please refer to Dose Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
SGLT 2 INHIBITOR COMBINATIONS	MC/DEL MC/DEL MC/DEL	SYNJARDY SYNJARDY XR XIGDUO XR	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		GLYXAMBI INVOKAMET INVOKAMET XR SEGLUROMET STEGLUJAN TRIJARDY XR	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried for at least 3 months at full therapeutic doses and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  <b>Glyxambi /Xigduo XR- Verify prior trials and failures or intolerance of preferred treatments from other diabetic categories.</b> <b>Synjardy XR</b> is not recommended for patients with type 1 DM or for the treatment of diabetic ketoacidosis.
DIABETIC MONITOR	MC	RELION TRUEMETRIX AIR BLOOD GLUCOSE MONITORING SYSTEM  TRUEMETRIX AIR BLOOD GLUCOSE MONITORING SYSTEM  TRUEMETRIX BLOOD GLUCOSE MONITORING SYSTEM	MC MC MC MC MC MC MC MC MC		ACCUCHECK ASCENSA ASSURE CONTOUR BREEZE Z EXACTECH FREESTYLE INSULINIX FREESTYLE LITE SYSTEM KIT PRECISION XTRA METER PRODIGY	<a href="#">Use PA Form# 20420</a>	Effective October 25th 2007, approvals for all non preferred meters/ test strips will require medical necessity documenting clinically significant features that are not available on any of the preferred meters.
DIABETIC TEST STRIPS	MC/DEL MC	RELION TRUEMETRIX  TRUEMETRIX	MC MC MC MC MC MC MC MC MC MC		ACCUCHECK ASCENSA ASSURE CONTOUR BREEZE Z EXACTECH FREESTYLE FREESTYLE LITE FREESTYLE INSULINIX PRECISION XTRA PRODIGY	<a href="#">Use PA Form# 20420</a>	Effective October 25th 2007, approvals for all non preferred meters/ test strips will require medical necessity documenting clinically significant features that are not available on any of the preferred meters.  Effective October 1, 2023, a maximum of 100 blood glucose test strips every 90 days will be available without Prior Authorization for members currently utilizing continuous glucose monitors (CGM).
INCRETIN MIMETIC	MC/DEL MC MC/DEL	RYBELSUS TRULICITY VICTOZA	MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL	5 8 8 8 8 8	OZEMPIC ADLYXIN BYDUREON BCISE MOUNJARO SOLIQUA XULTOPHY	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  <b>Soliqua</b> must try both insulin and a preferred incretin mimetic and have a medical necessity for use that is not based on convenience or simply due to the fact that one injection is needed instead of two.

CATEGORY	Coverage Indicator	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
DIABETIC - ORAL SULONYLUREAS	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	CHLORPROPAMIDE TABS GLIMEPIRIDE GLIPIZIDE TABS GLIPIZIDE ER TABS GLYBURIDE MICRONIZED TABS GLYBURIDE TABS <sup>1</sup> TOLAZAMIDE TABS TOLBUTAMIDE TABS	MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL		AMARYL TABS DIABETA TABS GLUCOTROL TABS GLUCOTROL XL TBCR GLYNASE TABS MICRONASE TABS	<a href="#">Use PA Form# 20420</a> 1. PA required for members $\geq 65$ . Glyburide has a greater risk of severe prolonged hypoglycemia in older adults.	Preferred drugs must be tried for at least 3 months at full therapeutic doses and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  DDI: All sulfonylureas (except glyburide) will now be non-preferred and require prior authorization if it is currently being used with either ranitidine or cimetidine. DDI: Glimepiride will now be non-preferred and require prior authorization if it is currently being used with either fluconazole (except 150mg strength) or fluvoxamine. Amaryl is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for concurrent use with either fluconazole or fluvoxamine.
DIABETIC -ORAL BIGUANIDES	MC/DEL MC/DEL	METFORMIN HCL TABS METFORMIN ER	MC MC MC MC/DEL		GLUCOPHAGE TABS GLUCOPHAGE XR TB24 FORTAMET METFORMIN ER OSMOTIC	<a href="#">Use PA Form# 20420</a>	Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
DIABETIC - THIAZOL / BIGUANIDE COMBO			MC/DEL MC/DEL MC MC		ACTOPLUS MET <sup>1</sup> ACTOPLUS MET XR AVANDARYL <sup>1</sup> AVANDAMET TABS <sup>1</sup>	<a href="#">Use PA Form# 20420</a> 1. Requires use of Actos, Metformin, or other preferred anti-diabetics.	DDI: Actos, Avandia, or any combination product with Actos or Avandia will now be non-preferred and require prior authorization if it is currently being used with gemfibrozil.
DIABETIC - / THIAZOL	MC/DEL	PIOGLITAZONE HCL <sup>1</sup>	MC/DEL MC		ACTOS TABS <sup>3</sup> AVANDIA TABS <sup>2</sup>	<a href="#">Use PA Form# 20420</a> 1. Pioglitazone HCL is non-preferred as monotherapy. Pioglitazone HCL is preferred if therapeutic doses of metformin, sulfonylurea or insulin are seen in members drug profile for at least 60 days within the past 18 months. 2. Current users of Avandia who have tried Actos will be able to continue use of Avandia. 3. Dosing limits apply. See Dose Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  DDI: Actos, Avandia, or any combination product with Actos or Avandia will now be non-preferred and require prior authorization if it is currently being used with gemfibrozil.
DIABETIC - ALPHAGLUCOSIDASE			MC		PRECOSE TABS	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
DIABETIC - SULONYLUREA / BIGUANIDE	MC/DEL	GLYBURIDE/METFORMIN	MC MC MC/DEL		GLUCOVANCE TABS <sup>1</sup> METAGLIP TABS <sup>1</sup> DUETACT <sup>2</sup>	<a href="#">Use PA Form# 20420</a> 1. Use individual ingredients. 2. Use Actos with generic glimepiride.	Approved for patients failing to achieve good diabetic control with maximal doses of individual components.
DIABETIC - MEGLITINIDES	MC	NATEGLINIDE	MC/DEL MC/DEL		PRANDIN TABS STARLIX TABS	<a href="#">Use PA Form# 20420</a>	Preferred drugs from other diabetic sub-categories must be tried and failed due to lack of inadequate diabetic control or intolerable side effects before non-preferred drug will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  DDI: Prandin is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for current use with both sporanox and gemfibrozil, due to a significant drug-drug interaction.
<b>GLUCOSE ELEVATING AGENTS</b>							
GLUCOSE ELEVATING AGENTS	MC/DEL MC/DEL MC/DEL	BAQSIMI <sup>1</sup> GVOKE <sup>2</sup> ZEGALOGUE <sup>3</sup>	MC		GLUCAGON DIAGNOSTIC KIT	<a href="#">Use PA Form# 20420</a> 1. For the treatment of patients $\geq 4$ years of age. 2. For the treatment of patients $\geq 2$ years of age. 3. For the treatment of patients $\geq 6$ years of age.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
<b>THYROID</b>							
THYROID EYE DISEASE			MC		TEPEZZA	<a href="#">Use PA Form# 20420</a>	
THYROID HORMONES	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	ARMOUR THYROID TABS CYTOMEL TABS ERMEZA <sup>1</sup> LEVOTHROID TABS LEVOTHYROXINE SODIUM TABS LEVOXYL TABS UNITHYROID TABS	MC MC/DEL MC MC/DEL		LEVOTHYROXINE SODIUM SOLR LIOTHYRONINE SYNTHROID TABS THYQUIDITY	<a href="#">Use PA Form# 20420</a> 1. Clinical PA is required to confirm diagnosis of dysphagia.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTITHYROID THERAPIES	MC/DEL MC/DEL	METHIMAZOLE TABS PROPYLTHIOURACIL TABS	MC/DEL		TAPAZOLE TABS	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.



Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required			Criteria
SOMATOSTATIC AGENTS			MC/DEL MC MC MC/DEL MC	7 8 8 8 8	OCTREOTIDE INJ <sup>1</sup> BYNFEZIA <sup>1</sup> MYCAPSSA <sup>1</sup> SANDOSTATIN <sup>1</sup> SOMATULINE <sup>1</sup>	<a href="#">Use PA Form# 10710</a> 1. Non-preferred products must be used in specified step order.		
<b>GROWTH HORMONE ANTAGONISTS</b>								
GH ANTAGONISTS			MC		SOMAVERT	<a href="#">Use PA Form# 10710</a>	Approved for acromegaly patients failing surgery/radiation/drug therapy including bromocriptine and sandostatin.	
<b>VASOPRESSIN RECEPTOR ANTAGONIST</b>								
VASOPRESSIN RECEPTOR ANTAGONIST			MC MC/DEL		JYNARQUE <sup>1</sup> SAMSCA	<a href="#">Use PA Form# 20420</a> 1. Clinical PA required for appropriate diagnosis	<b>Samsca Drug Warning-</b> Avoid use in patients with underlying liver disease, including cirrhosis, because the ability to recover from liver injury may be impaired. Limit duration of therapy to 30 days to minimize the risk of liver injury. <b>DDI:</b> Jynarque- Concomitant use with strong CYP3A inhibitors is contraindicated. Avoid concomitant use of Jynarque with OATP1B1/B3 and OAT3 substrates (e.g. statins, bosantan, glyburide, nateglinide, repaglinide, methotrexate, furosemide).	
<b>URINARY INCONTINENCE</b>								
VASOPRESSINS	MC/DEL MC/DEL	DESMOPRESSIN TABS DDAVP SOLN	MC/DEL MC/DEL MC MC/DEL MC	5 6 8 8 8	DDAVP TABS DESMOPRESSIN SPRAY <sup>1</sup> DESMOPRESSIN ACETATE SOLN <sup>1</sup> NOCDURNA <sup>1</sup> NOCTIVA <sup>1</sup>	<a href="#">Use PA Form# 20420</a> 1. Products must be used in specified step order. Nocturnal enuresis patients will be encouraged to periodically attempt stopping DDAVP.	Approved for central diabetes insipidus and for nocturnal enuresis. For nocturnal enuresis- must be over 6 years old, must fail an adequate trial of alarm training (higher success rate, lower relapse rate) and must periodically attempt weaning (at 6 month intervals).	
ANTISPASMODICS	MC/DEL MC/DEL	OXYBUTYNIN TOLTERODINE	MC/DEL MC/DEL MC/DEL	8 8 8	DARIFENACIN ER TAB DITROPAN FLAVOXATE HCL TAB	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.	
ANTISPASMODICS - LONG ACTING	MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	FESOTERODINE GELNIQUE GEL PACKET MYRBETRIQ OXYBUTYNIN ER TABS OXYTROL SOLIFENACIN SUCCINATE TAB TROSPiUM	MC MC/DEL MC MC/DEL MC/DEL MC MC	8 8 8 8 8 8 8	DITROPAN XL TBCR ENABLEX <sup>1,2</sup> GEMTESA <sup>2</sup> TOLTERODINE TAB TOVIAZ VESICARE <sup>1</sup> VESICARE LS <sup>3</sup>	<a href="#">Use PA Form# 20420</a> 1. See Criteria Section. 2. Use a preferred long acting antispasmodic. 3. For the treatment of patients $\geq$ 2 years of age.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  <b>1. Vesicare 5mg and Enablex 7.5mg</b> maximum doses if given with drugs known to be significant CYP3A4 inhibitors (ketoconazole, sporanox, erythromycin, fluconazole, nefazodone, neflunavir, and ritonavir). <b>DDI:</b> Enablex 15mg and Vesicare 10mg will now be non-preferred and require prior authorization if they are currently being used in combination with any of the following medications: clarithromycin, erythromycin, ketek, crixivan, norvir, ketoconazole, fluconazole (except 150mg strength), sporanox, or nefazodone.	
CHOLINERGIC	MC/DEL	BETHANECHOL	MC/DEL		URECHOLINE	<a href="#">Use PA Form# 20420</a>		
HYPERAMMONIA TREATMENTS	MC	CARBAGLU TABS	MC		CARGLUMIC ACID TABS	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.	
UREA CYCLE DISORDER	MC MC	BUPHENYL TABLET PHEBURANE GRANULES	MC MC MC MC/DEL MC/DEL		BUPHENYL POWDER RAVICTI LIQUID OLPRUVA SODIUM PHENYLBUTYRATE POWDER SODIUM PHENYLBUTYRATE TAB	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  <b>Olpruva:</b> As adjunctive therapy to standard of care, which includes dietary management, for the chronic management of adult and pediatric patients weighing 20kg or greater and with a body surface area (BSA) of 1.2m <sup>2</sup> or greater, with urea cycle disorders (UCDs) involving deficiencies of carbamylphosphate synthetase (CPS), ornithine transcarbamylase (OTC), or argininosuccinic acid synthetase (AS).	
<b>METABOLIC MODIFIER</b>								
HERED. TYROSINEMIA			MC MC MC	6 6 8	ORFADIN NITYR HARLIKU <sup>1</sup>	<a href="#">Use PA Form# 20420</a> 1. Clinical PA is required to establish diagnosis and medical necessity.	Approved for Type 1 hereditary tyrosinemia patients. Must include laboratory evidence of dx at first PA. Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.	
FABRY DISEASE AGENTS			MC MC MC/DEL		ELFABRIO <sup>1</sup> FABRAZYME <sup>2</sup> GALAFOLD <sup>1</sup>	<a href="#">Use PA Form# 20420</a> 1. Clinical PA to verify appropriate diagnosis. 2. For the treatment of patients 2 years of age and older.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  <b>Elfabrio and Galfold:</b> For the treatment of adults with confirmed Fabry disease.	
<b>ANTIHYPERTENSIVES / CARDIAC</b>								
CARDIAC GLYCOSIDES	MC/DEL MC/DEL MC/DEL	DIGITEK TABS DIGOXIN LANOXIN				<a href="#">Use PA Form# 20420</a>		
CARDIAC MYOSIN INHIBITORS			MC		CAMZYOS	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  <b>Camzyos:</b> For the treatment of adults with symptomatic New York Heart Association (NYHA) class II-III obstructive hypertrophic cardiomyopathy (HCM) to improve functional capacity and symptoms. <b>DDI:</b> Concomitant use of Camzyos with a moderate to strong CYP2C19 inhibitor or a strong CYP3A4 inhibitor is contraindicated.	
CARDIAC - SINUS NODE INHIBITORS			MC		CORLANOR	<a href="#">Use PA Form# 20420</a>	In patients with stable, symptomatic chronic heart failure with left ventricular ejection fraction $\leq$ 35%, who are in sinus rhythm with resting heart rate $\geq$ 70 beats per minute (bpm) and either are on maximally tolerated doses of beta-blockers or have a contraindication to beta-blocker use.	

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
CARDIAC- ERAs			MC		TRYVIO	<a href="#">Use PA Form#20420</a>	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p> <p><b>Tryvio:</b> In combination with other antihypertensive drugs, is indicated for the treatment of resistant hypertension, to lower blood pressure (BP) in adult patients who are not adequately controlled on other drugs. Resistant HTN is defined as a patient who takes at least 3 different class antihypertensive medications with complementary mechanisms including thiazide, ACE inhibitor, ARB, long-acting calcium channel blocker, with a trial of spironolactone, unless contra-indicated.</p>
CARDIAC- SOLUBLE GUANYLATE CYCLASE STIMULATORS			MC/DEL		VERQUVO	<a href="#">Use PA Form# 20420</a>	
CARDIAC RISK REDUCTION- SGLT2/GLP-1			MC MC MC/DEL		INPEFA <sup>1</sup> LODOCO WEGOVY	<a href="#">Use PA Form #23976</a> 1. To reduce the risk of cardiovascular death, hospitalization for heart failure, and urgent heart failure visit in adults with: Heart failure or Type 2 diabetes mellitus, chronic kidney disease, and other cardiovascular risk factors.	<p>Other Preferred SGLT inhibitors must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p> <p><b>Lodoco:</b> Patient must have tried and failed generic colchicine due to lack of efficacy or intolerable side effects</p> <p><b>Wegovy:</b> Patient does not have diagnosis of diabetes, end stage renal disease/dialysis, or HFrEF (EF &lt; 45%)</p> <ul style="list-style-type: none"> <li>• Patient has BMI &gt; 27 kg/m<sup>2</sup>, and is not being used for weight loss only</li> <li>• Patient has history of at least one of the following:               <ul style="list-style-type: none"> <li>o Stroke</li> <li>o Myocardial Infarction</li> <li>o Symptomatic peripheral arterial disease</li> </ul> </li> </ul>
ANTIANGINALS--Isosorbide Dinitrate/ Mono-Nitrates	MC/DEL MC/DEL	ISOSORBIDE MONONITRATE TABS ISOSORBIDE MONONITRATE ER	MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC		DILATRATE SR CPCR ISORDIL TITRADOSE TABS ISOSORBIDE DINITRATE SUBL ISOSORBIDE DINITRATE CR TBCR ISOSORBIDE DINITRATE ER TBCR ISOSORBIDE DINITRATE TD TBCR IMDUR TB24 ISMO TABS MONOKET TABS	<a href="#">Use PA Form# 20420</a>	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p>
NITRO - OINTMENT/CAP/CR	MC/DEL MC/DEL MC MC	NITROBID OINT NITROGLYCERIN CPCR NITROL OINT NITRO-TIME CPCR				<a href="#">Use PA Form# 20420</a>	
NITRO - PATCHES	MC/DEL MC/DEL	NITROGLYCERIN PT24 NITRO-DUR PT 24 0.8MG	MC MC/DEL		NITRODISC PT24 NITRO-DUR PT24	<a href="#">Use PA Form# 20420</a>	<p>All preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p>
NITRO - SUBLINGUAL/ SPRAY	MC/DEL	NITROSTAT SUBL	MC/DEL MC MC		NITROQUICK SUBL NITROLINGUAL SOLN NITROLINGUAL TABS	<a href="#">Use PA Form# 20420</a>	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p>
BETA BLOCKERS - NON SELECTIVE	MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	CARVEDILOL LEVATOL TABS NADOLOL TABS PINDOLOL TABS PROPRANOLOL HCL SOLN <sup>1</sup> PROPRANOLOL HCL TABS <sup>1</sup> PROPRANOLOL HCL 60MG TABS PROPRANOLOL LA CAPS RANOLAZINE ER TABS SOTALOL AF SOTALOL HCL TABS TIMOLOL MALEATE TABS	MC MC/DEL MC MC MC MC/DEL MC/DEL MC/DEL MC MC MC MC MC MC MC		ASPRUZO BETAPACE TABS BETAPACE AF TABS COREG CR <sup>3</sup> COREG TABS CORGARD TABS INDERAL TABS HEMANGEOL SOL INDERAL XL CAP INDERAL LA CPCR INNOPRAN XL RANEXA	<a href="#">Use PA Form# 20420</a> 1. Recommend using BID since its effects do not last 24 hours. 2. Please use other strengths in combination to obtain this dose. 3. Dosing limits still apply. Please see dose consolidation list.	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p> <p><b>DDI:</b> Concomitant use of <b>Ranolazine</b> products with strong CYP3A inhibitors, including ketoconazole, itraconazole, clarithromycin, nefazodone, nefazodone, ritonavir, indinavir, and saquinavir, is contraindicated.</p>
BETA BLOCKERS - CARDIO SELECTIVE	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	ACEBUTOLOL HCL CAPS ATENOLOL TABS <sup>1</sup> BETAXOLOL HCL TABS BISOPROLOL FUMARATE TABS BYSTOLIC METOPROLOL TARTRATE TABS <sup>1</sup> METOPROLOL ER NEBIVOLOL HCL TAB	MC MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL		KERLONE TABS LOPRESSOR TABS SECTRAL CAPS TENORMIN TABS TOPROL XL TB24 ZEBETA TABS	<a href="#">Use PA Form# 20420</a> 1. Recommend using <b>Atenolol</b> (and <b>Metoprolol</b> ) BID since its effects do not last 24 hours.	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p>

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
BETA BLOCKERS - ALPHA / BETA	MC/DEL	LABETALOL HCL TABS	MC		TRANDATE TABS	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
BETA BLOCKERS & DURECTIC COMBOS	MC/DEL	METOPROLOL-HYDROCHLOROTHIAZIDE TAB	MC/DEL		DUTOPROL	<a href="#">Use PA Form# 20420</a>	
CALCIUM CHANNEL BLOCKERS-- Amlodipine, Felodipines, Nifedipines, Nisoldipine, and Verapamil	MC/DEL	AMLODIPINE <sup>1</sup>	MC/DEL MC MC/DEL		KATERZIA NORLIQVA NORVASC TABS <sup>1</sup>	<a href="#">Use PA Form# 20420</a> 1. Dosing limits apply, see Dose Consolidation List.	
	MC	DILTIA XT CP24	MC/DEL	5	DILACOR XR CP24 <sup>1</sup>	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed (in step-order) due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL	DILTAZEM HCL ER CP24	MC/DEL	6	TAZTIA <sup>1</sup>	<a href="#">Use PA Form# 20420</a>	1. Products must be used in specified order or PA will be required. Just write "Diltiazem 24-hour" and the pharmacy will use a preferred long acting Diltiazem that does not require PA.
	MC/DEL	DILTAZEM HCL XR CP24	MC/DEL	8	DILTAZEM HCL TABS <sup>1</sup>	<a href="#">Use PA Form# 20420</a>	DDI: All preferred Diltiazem will now be non-preferred and require prior authorization if they are currently being used in combination with either enablex 15mg or vesicare 10mg. All non-preferred Diltiazem require prior authorization, but with any prior authorization request, the member's drug profile will also be monitored for current use with enablex 15mg or vesicare 10mg.
	MC/DEL	DILTAZEM CD 300MG CP24	MC/DEL	8	DILTAZEM HCL ER CP12 <sup>1</sup>	<a href="#">Use PA Form# 20420</a>	
	MC/DEL	DILTAZEM CD 360MG CP24	MC/DEL	8	DILTAZEM HCL ER CP12 <sup>1</sup>	<a href="#">Use PA Form# 20420</a>	
	MC	CARTIA XT CP24 <sup>1</sup>					
	MC/DEL	DILTAZEM CD CP24 <sup>1</sup>					
	MC/DEL	DILTAZEM HCL ER CP24 <sup>1</sup>					
	MC/DEL	DILTAZEM XR CP24 <sup>1</sup>					
	MC/DEL	TIAZAC CP24 <sup>1</sup>					
			MC/DEL MC/DEL		PLENDIL TB24 FELODIPINE	<a href="#">Use PA Form# 20420</a>	Other Preferred calcium channel blockers must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
			MC MC		CARDENE SR CPCR NICARDIPINE HCL CAPS	<a href="#">Use PA Form# 20420</a>	Other Preferred calcium channel blockers must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL	AFEDITAB CR	MC/DEL		ADALAT CC TBCR <sup>1</sup>	<a href="#">Use PA Form# 20420</a>	Preferred drug must be tried and failed in step order due to lack of efficacy or intolerable side effects before non-preferred drugs in step order will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL	NIFEDIAC CC	MC/DEL		NIFEDIPINE CAPS	<a href="#">Use PA Form# 20420</a>	
	MC/DEL	NIFEDICAL XL TBCR	MC/DEL		PROCARDIA CAPS	<a href="#">Use PA Form# 20420</a>	
	MC/DEL	NIFEDIPINE TBCR	MC/DEL		PROCARDIA XL TBCR	<a href="#">Use PA Form# 20420</a>	
	MC/DEL	NIFEDIPINE ER TBCR					
			MC MC		SULAR TB24 SULAR CR <sup>1</sup>	<a href="#">Use PA Form# 20420</a> 1. Established users of 10MG and 20MG strengths are grandfathered.	
	MC/DEL	VERAPAMIL HCL CR TBCR	MC/DEL		CALAN TABS	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed (in step-order) due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL	VERAPAMIL HCL ER TBCR	MC/DEL		CALAN SR TBCR	<a href="#">Use PA Form# 20420</a>	
	MC/DEL	VERAPAMIL HCL SR TBCR	MC/DEL MC MC/DEL		COVERA-HS TBCR ISOPTIN-SR VERAPAMIL HCL ER CP24 VERAPAMIL HCL SR CP24 VERAPAMIL HCL TABS VERELAN CP24 VERELAN PM CP24	<a href="#">Use PA Form# 20420</a> Products must be used in specified order or PA will be required. Just write "Verapamil 24-hour" and the pharmacy will use a preferred long acting generic that does not require PA.	
ANTIARRHYTHMICS	MC/DEL	AMIODARONE HCL	MC/DEL		CORDARONE	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL	DISOPYRAMIDE	MC/DEL		DISOPYRAMIDE	<a href="#">Use PA Form# 20420</a>	
	MC/DEL	FLECAINIDE	MC/DEL		MULTAQ	<a href="#">Use PA Form# 20420</a>	
	MC/DEL	MEXILETINE HCL	MC/DEL		NORPACE	<a href="#">Use PA Form# 20420</a>	DDI: Amiodarone will now be non-preferred and require prior authorization if it is currently being used in combination with either lovastatin (doses greater than 40mg/day) or lipitor (doses greater than 20mg/day) or levofloxacin or gemifloxacin, or moxifloxacin, or ofloxacin.
	MC/DEL	PROCAINAMIDE	MC/DEL		PACERONE	<a href="#">Use PA Form# 20420</a>	
	MC/DEL	PROPAFENONE	MC		QUINIDEX	<a href="#">Use PA Form# 20420</a>	DDI: Multaq will be preferred unless the following medications are seen in the member's drug profile within the last 35 days for brand name medications or 90 days for generic medications: erythromycin, amiodarone and other antiarrhythmics, TCA's, phenothiazine, ketoconazole, itraconazole, voriconazole, cyclosporine, telithromycin, clarithromycin, nefazodone, ritonavir.
	MC	QUINAGLUTE	MC/DEL		TAMBOCOR	<a href="#">Use PA Form# 20420</a>	
	MC/DEL	QUINIDINE GLUCONATE	MC/DEL		TIKOSYN <sup>1</sup>	<a href="#">Use PA Form# 20420</a>	
	MC/DEL	QUINIDINE SULFATE	MC/DEL		RYTHMOL SR	<a href="#">Use PA Form# 20420</a>	
			MC/DEL		RYTHMOL	<a href="#">Use PA Form# 20420</a>	
ACE INHIBITORS	MC/DEL	BENAZEPRIL HCL	MC	5	MAVIK TABS	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Non-preferred products are subject to step-order requirements unless clinical circumstances warrant exception.
	MC/DEL	CAPTOPRIL TABS	MC/DEL	5	ACCUPRIL TABS	<a href="#">Use PA Form# 20420</a>	
	MC/DEL	ENALAPRIL MALEATE TABS	MC/DEL	8	ACEON TABS <sup>1</sup>	<a href="#">Use PA Form# 20420</a>	
	MC/DEL	FOSINOPRIL SODIUM	MC/DEL	8	ALTACE CAPS <sup>1</sup>	<a href="#">Use PA Form# 20420</a>	
	MC/DEL	LISINOPRIL TABS	MC	8	EPANED	<a href="#">Use PA Form# 20420</a>	
	MC/DEL	RAMIPRIL	MC/DEL	8	LOTENSIN TABS <sup>1</sup>	<a href="#">Use PA Form# 20420</a>	
	MC/DEL	QUINAPRIL HCL	MC/DEL	8	MOEXIPRIL HCL <sup>1</sup>	<a href="#">Use PA Form# 20420</a>	
			MC	8	MONOPRIL HCT TABS <sup>1</sup>	<a href="#">Use PA Form# 20420</a>	

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
			MC/DEL MC MC/DEL MC/DEL	8 8 8 8	PRINIVIL TABS <sup>1</sup> QBRELIS UNIVASC <sup>1</sup> ZESTRIL TABS <sup>1</sup>		
ANGIOTENSIN RECEPTOR BLOCKER	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	AMLODIPINE-OLMESARTAN TAB <sup>3</sup> IRBESARTAN <sup>1</sup> LOSARTAN <sup>1</sup> MICARDIS TABS <sup>3</sup> OLMESARTAN <sup>1</sup> TELMISARTAN <sup>1</sup>	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC	8 8 8 8 8 8 8	ATACAND TABS AVAPRO BENICAR TABS COZAAR DIOVAN EDARBI TEVETEN TABS	<a href="#">Use PA Form# 20420</a> 1. Dosing limits apply, please see Dose Consolidation List. 2. Use preferred active ingredients which are available without PA. 3. Preferred without a PA only if patient on a diabetic therapy or prior ACE therapy.	Per best practices patient should have trialed prior therapy of ACE inhibitor or currently on a diabetic therapy.
DIRECT RENIN INHIBITOR			MC/DEL MC/DEL MC/DEL		AMTURNIDE TEKTURN <sup>1</sup> TEKAMLO	<a href="#">Use PA Form# 20420</a> 1. Must show failure of single and combination therapy from all preferred antihypertensive categories.	
ANTIHYPERTENSIVES - CENTRAL	MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	CLONIDINE HCL TABS GUANFACINE HCL TABS HYDRALAZINE HCL TABS HYLOREL TABS METHYLDOPA TABS MINOXIDIL TABS PRAZOSIN HCL CAPS RESERPINE TABS	MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC/DEL		CLONIDINE PATCH CLONIDINE TTS GUANABENZ ACETATE TABS ISMELIN TABS MINIPRESS CAPS NEXICLON TENEX TABS	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ACE INHIBITORS AND CA CHANNEL BLOCKERS			MC/DEL MC MC MC/DEL	8 8 8 9	AMLODIPINE/BENAZEPRIL PRESTALIA <sup>1</sup> TARKA TBCR LOTREL CAPS	<a href="#">Use PA Form# 20420</a> 1. Prestalia will only be approved for patients $\geq$ 18 years of age. Use individual preferred generic medications.	
ACE AND THIAZIDE COMBO'S	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	BENAZEPRIL HCL/HYDROCHLOR CAPTOPRIL/HYDROCHLOROTHIA ENALAPRIL MALEATE/HCTZ TABS LISINOPRIL-HCTZ TABS LOTENSIN HCT TABS	MC/DEL MC MC/DEL MC/DEL MC/DEL		ACCURETIC TABS MONOPRIL HCT TABS PRINZIDE TABS UNIRETIC TABS ZESTORETIC TABS	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
BETA BLOCKERS AND DIURETIC COMBO'S	MC/DEL MC/DEL MC/DEL	ATENOLOL/CHLORTHALIDONE BISOPROLOL FUMARATE/HCTZ PROPRANOLOL/HCTZ	MC/DEL MC/DEL MC MC MC/DEL		CORZIDE TABS LOPRESSOR HCT TABS TENORETIC TIMOLIDE 10/25 TABS ZIAC TABS	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ARB'S AND CA CHANNEL BLOCKERS	MC/DEL MC/DEL MC/DEL	AMLODIPINE/VALSARTAN AMLODIPINE/VALSARTAN HCT TRIBENZOR	MC/DEL MC MC/DEL MC/DEL		AZOR BYVALSON EXFORGE EXFORGE HCT	<a href="#">Use PA Form# 20420</a>	DDI: Byvalson will be non-preferred and require a prior authorization if it is currently being used in combination with drugs known to be significant CYP2D6 inhibitors (e.g. quinidine, propafenone, fluoxetine, paroxetine). Per best practices, patient should have trialed prior therapy of ACE inhibitor or currently on a diabetic therapy.
ARB'S AND DIURETICS	MC/DEL MC/DEL MC/DEL MC/DEL	BENICAR HCT <sup>1</sup> LOSARTAN HCT <sup>1</sup> MICARDIS HCT TABS <sup>1</sup> VALSARTAN-HCT <sup>1</sup>	MC/DEL MC/DEL MC MC/DEL MC/DEL MC	7 8 8 8 8 8	IRBESARTAN HYDROCHLOROTHIAZIDE ATACAND HCT TABS AVALIDE TABS <sup>1</sup> DIOVAN HCT TABS <sup>1</sup> HYZAAR TABS TEVETEN HCT TABS	<a href="#">Use PA Form# 20420</a> 1. Dosing limits apply, see Dose Consolidation List.	Per best practices, patient should have trialed prior therapy of ACE inhibitor or currently on a diabetic therapy.
ANGIOTENSIN MODULATORS-ARB COMBINATION	MC	ENTRESTO	MC/DEL MC		EDARBYCLOR ENTRESTO SPRINKLES	<a href="#">Use PA Form# 20420</a>	
ARB'S AND DIRECT RENIN INHIBITOR COMBINATION			MC/DEL		VALTURNA	<a href="#">Use PA Form# 20420</a>	



Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
CHOLESTEROL - HMG COA + ABSORB INHIBITORS LESS POTENT DRUGS/- COMBINATIONS	MC/DEL MC/DEL MC/DEL	EZETIMIBE TABS LOVASTATIN TABS <sup>2</sup> PRAVASTATIN <sup>2</sup>	MC MC/DEL MC/DEL MC MC/DEL MC MC MC/DEL MC/DEL MC	8 8 8 8 8 8 8 8 8 8 8	ALTOPREV TB24 FLUVASTATIN TAB ER LESCOL XL TB24 LIVALO MEVACOR TABS NEXLETOL NEXLIZET PRAVACHOL TABS PRAVIGARD ZETIA TABS	<a href="#">Use PA Form# 20420</a> 2. Dosing limits apply, please see Dosage Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. <b>Zetia</b> will be approved for patients unable to tolerate all other therapies or unable to achieve cholesterol goal with maximally tolerated dose of most potent statins.  DDI: <b>Lescol</b> will now be non-preferred and require prior authorization if it is currently being used in combination with diclofenac. DDI: <b>Lovastatin</b> (doses greater than 40mg/day) will now be non-preferred and require prior authorization if it is currently being used in combination with amiodarone. DDI: <b>Lovastatin</b> (doses greater than 20mg per day) will now be non-preferred and require prior authorization if it is currently being used in combination cyclosporine. DDI: All preferred statins will now be non-preferred and require prior authorization if it is currently being used in combination with gemfibrozil.
CHOLESTEROL - HMG COA + ABSORB INHIBITORS STATIN/ NIACIN COMBO	MC	SIMCOR	MC		ADVICOR TBCR	<a href="#">Use PA Form# 20420</a>	
FAMILIAL HYPERCHOLESTEROLEMIA	MC MC	PRALUENT (LABLER 72733) PEN <sup>1,2,3,5</sup> REPATHA <sup>1,2,3</sup>	MC MC MC MC		EVKEEZA <sup>1,4</sup> JUXTAPI <sup>1</sup> KYNAMRO <sup>1</sup> LEQVIO	<a href="#">Use PA Form# 20420</a> 1. Clinical PA required for appropriate diagnosis. 2. Quantity limits apply. 3. Documented adherence to lipid lowering medications and abstinence from tobacco for previous 90 days. 4. For treatment of patients $\geq$ 12 years of age. 5. Approval of Praluent NDC's with labeler code 00024 will be considered only if labeler code 72733 NDC's are on a long-term backorder and unavailable from the manufacturer.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  <b>Juxtapid</b> is contraindicated with strong CYP3A4 inhibitors. Juxtapid dosage should not exceed 30mg daily when it is used concomitantly with weak CYP3A4 inhibitors.  <b>Kynamro</b> requires an appropriate lab testing prior to starting (ALT <ast), alkaline="" and="" bilirubin,="" every="" first="" for="" liver-related="" monthly="" months.<br="" phosphatase="" tests="" the="" then="" three="" total="" year,=""></ast),> <b>Repatha and Praluent Criteria for approval:</b> The patient's age is FDA approved for the given indication AND • Concurrent use with statin therapy AND • Documented adherence to prescribed lipid lowering medications for the previous 90 days AND • Recommended or prescribed by a lipidologist or cardiologist AND • Inability to reach goal LDL-C despite a trial of 2 or more maximum tolerated dose of statins (one of which must be atorvastatin or rosuvastatin) and ezetimibe 10mg daily.  <b>Additional criteria for the diagnosis of heterozygous familial hypercholesterolemia (HeFH): (both are required):</b> Total cholesterol > 290 mg/dL OR LDL-C > 190 mg/dL AND one of the following • Presence of tendon xanthomas OR • In 1st or 2nd degree relative-documented tendon xanthomas, MI at age $\leq$ 60 years or TC > 290 mg/dL.  <b>Additional criteria for the diagnosis of clinical atherosclerotic cardiovascular disease:</b> History of MI, angina, coronary or other arterial revascularization, stroke, TIA, or PVD of atherosclerotic origin.  <b>Additional criteria for the diagnosis of homozygous familial hypercholesterolemia (Repatha only):</b> Total cholesterol levels > 290mg/dL or LDL-C > 190mg/dL (adults) OR Total cholesterol levels > 260mg/dL or LDL-C > 155mg/dL (children $<$ 16 years) and TG within reference range OR Confirmation of diagnosis by gene testing.
FAMILIAL HYPERCHOLESTEROLEMIA AND HYPERTRIGLYCERIDEMIA					TRYNGOLZA	<a href="#">Use PA Form# 20420</a>	Tryngolza requires fasting triglycerides of $\geq$ 880 mg/dL and confirmed genetically identified familial chylomicronemia syndrome (FCS).
<b>HYPERPHAGIA - MISC</b>							
HYPERPHAGIA - MISC			MC	8	VYKAT XR		FDA approved for the treatment of hyperphagia in adults and pediatric patients 4 years of age and older with Prader-Willi syndrome (PWS).
<b>PULMONARY ANTI-HYPERTENSIVES</b>							
PULMONARY ANTI-HYPERTENSIVES	MC MC/DEL MC/DEL	EPOPROSTENOL INJ <sup>3</sup> SILDENAFIL TADALAFIL	MC/DEL MC MC/DEL MC MC MC MC MC/DEL MC MC MC MC MC MC MC MC MC/DEL MC		ADEMPAS <sup>1,3</sup> ADCIRCA <sup>4</sup> ALYQ TAB FLOLAN <sup>3</sup> LIQREV OPSUMIT <sup>1,2</sup> OPSYNVI <sup>4</sup> ORENITRAM REMODULIN <sup>3</sup> REVATIO <sup>4</sup> TADLIQ <sup>4</sup> TYVASO UPTRAVI VELVETRI <sup>3</sup> WINREVAIR <sup>4</sup> YUTREPIA	<a href="#">Use PA Form# 20420</a> 1. Requires previous trials/failure of multiple preferred medications. 2. Dosing limits apply, see the Dose Consolidation List. 3. Require WHO Group 1 diagnosis of primary PAH (Primary Pulmonary Hypertension) and NYHA functional class 3 or 4. 4. Require WHO Group 1 diagnosis of primary PAH (Primary Pulmonary Hypertension) and NYHA functional class 2 or 3.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  <b>Sildenafil</b> will be preferred with clinical PA for treatment of pulmonary arterial hypertension (WHO Group 1) in adults to improve exercise ability and delay clinical worsening. Avoid concomitant use of sildenafil with moderate or strong Cyp3A inhibitors.  <b>DDI: Uptravi</b> will require a prior authorization if it is currently being used in combination with strong inhibitors of CYP2C8 (gemfibrozil).  <b>DDI: Opsumit</b> will require a prior authorization if it is currently being used in combination with drugs known to be significant CYP3A inhibitors (ketoconazole, itraconazole, clarithromycin, indinavir, nefazodone, neflunavir, ritonavir, atazanavir, saquinavir and telithromycin).  <b>DDI: Adempas</b> will require a prior authorization if it is currently being used in combination with drugs known to be PDE inhibitors should be avoided (including dipyridamole, adcirca and tadalafil) with adempas.  <b>Liqrev:</b> treatment of pulmonary arterial hypertension (WHO Group 1) in adults to improve exercise ability and delay clinical worsening. Avoid concomitant use of liqrev with moderate or strong CYP3A inhibitors.



Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
<b>ALLERGY / ASTHMA THERAPIES</b>							
ANAPHYLACTIC DEVICES	MC MC/DEL MC/DEL MC/DEL	AUVI-Q EPINEPHRINE EPIPEN EPIPEN JR	MC MC		NEFFY TWINJECT	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ALLERGEN IMMUNOTHERAPY			MC MC MC MC MC		ODACTRA ORALAIR <sup>1</sup> PALFORZIA RAGWITEK GRASTEK	<a href="#">Use PA Form# 20420</a> 1. See criteria section	Prescriber must provide the testing to show that the patient is allergic to the components in the prescribed therapy and must provide a clinically valid rationale why single agent sublingual therapy is being chosen over subcutaneous therapy.  <b>Palforzia</b> is approved for use in patients with a confirmed diagnosis of peanut allergy. Initial dose escalation may be administered to patients aged 4 through 17 years. Up-dosing and maintenance may be continued in patients 4 years of age and older. <b>Odactra</b> is approved for use in persons 12 through 65 years of age. Note that Odactra is not indicated for the immediate relief of allergic symptoms.  Treatment must start 12 weeks before expected onset of pollen season and only after confirmed by positive skin test or in vitro testing for pollen-specific IgE antibodies for any of the 5 grass species contained in Oralair.  <b>Oralair:</b> Patient age ≥10 years and ≤65 years. Have an auto-injectable epinephrine on-hand.
ANTIASTHMATIC - ANTICHOLINERGICS - INHALER	MC MC/DEL MC/DEL	INCRUSE ELLIPTA <sup>3</sup> SPIRIVA HANDIHALER <sup>1,2</sup> SPIRIVA RESPIMAT	MC MC/DEL		LONHALA MAGNAIR TUDORZA	<a href="#">Use PA Form# 20420</a> 1. Quantity limit of 1 inhalation daily (1 capsule for inhalation daily). Spiriva will require PA if Combivent or Atrovent nebulizer solution is in member's current drug profile. 2. We ask physicians to write "asthma" on the prescription whenever Spiriva is primarily being used for that condition. 3. Quantity limit of 1 inhalation daily.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIASTHMATIC - Dipeptidyl Peptidase 1 Inhibitors	MC	BRINSUPRI				<a href="#">Use PA Form# 20420</a> 1. Clinical PA is required to establish diagnosis and medical necessity.	<b>BRINSUPRI</b> required criteria include: <ul style="list-style-type: none"><li>• Imaging confirming bronchiectasis and no overlapping asthma/COPD required.</li><li>• Documented airway clearance.</li><li>• Greater than 2 exacerbations requiring antibiotic therapy in the last 12 months.</li><li>• Must be approved by pulmonologist.</li></ul>
ANTIASTHMATIC - PHOSPHODIESTERASE 4 INHIBITORS	MC/DEL	ROFLUMILAST	MC/DEL MC		DALIRESP OHTUVAYRE <sup>1</sup>	<a href="#">Use PA Form# 20420</a> 1. For the maintenance treatment of chronic obstructive pulmonary disease (COPD) in adult patients.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIASTHMATIC - ANTICHOLINERGICS - NEBULIZER	MC/DEL	IPRATROPIUM BROMIDE SOLN	MC MC/DEL		ATROVENT SOLN YUPELRI	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIASTHMATIC - ANTIINFLAMMATORY AGENTS	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	CROMOLYN SODIUM NEBU DUPIXENT <sup>2,4</sup> FASENRA <sup>6</sup> FASENRA <sup>6</sup> AUTO INJCT XOLAIR <sup>1,4</sup>	MC MC MC MC MC	8 8 8 8	CINQAIR <sup>3</sup> NUCALA <sup>2</sup> RHAPSIDO <sup>4</sup> TEZSPIRE <sup>5</sup>	<a href="#">Use PA Form# 20420</a> 1. Need max inhaled steroids and written by pulmonary or allergy specialist. Must have elevated IgE and ≥ age 6. 2. For patients with severe asthma aged 12 years or older and eosinophilia. 3. For patients ≥ 18 years of age with eosinophilia. 4. Clinical PA required to establish diagnosis and medical necessity. 5. For adult and pediatric patients aged 12 years and older with severe asthma.  6. For patients ≥ 6 years of age for eosinophilia.	All will require suboptimal response to maximal doses of inhaled steroid as evidenced by asthmatic ER/Hospital admissions and Allergy/Pulmonary specialist management. <b>Dupixent</b> limited to patient with asthma not controlled on high dose ICS-LABA who have eosinophil greater than or equal to 150 cells or the patient is depend on an oral corticosteroid. <b>Fasenra</b> , <b>Nucala</b> and <b>Cinqair</b> are not indicated for treatment of other eosinophilic conditions and are not indicated for the relief of acute bronchospasm or status asthmaticus.  <b>Rhapsido</b> for Chronic Spontaneous Urticaria - must have had an inadequate clinical response of at least 14-days with at least two different second-generation antihistamines at 4 times standard dose. Must continue use of second-generation antihistamine. Must be prescribed by or in consultation with either allergist-immunologist, dermatologist, pulmonologist, or otolaryngologist.
ANTIASTHMATIC - NASAL STEROIDS	MC/DEL MC/DEL MC MC/DEL MC/DEL MC	BUDESONIDE SPRAY FLUTICASONE SPR <sup>3</sup> OLOPATADINE SPRAY OMNARIS SPR <sup>3</sup> TRIAMCINOLONE NS QNASL	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC	8 8 8 8 8 8	DYMISTA FLONASE SUSP <sup>2,3</sup> FLUNISOLIDE SOLN <sup>1,3</sup> NASONEX SUSP RHINOCORT AERO <sup>2,3</sup> RHINOCORT AQUA SUSP <sup>2,3</sup> RYALTRIS <sup>4</sup> TRI-NASAL SOLN <sup>2,3</sup>	<a href="#">Use PA Form# 20420</a> 1. All preferred drugs must be tried before moving to non preferred steps. 2. All step 5 medications need to be tried before moving to step 8's. 3. Dosing limits apply to whole category, see Dosage Consolidation List.	Preferred drugs and step therapy must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  <b>Xhance</b> will be considered for the treatment of nasal polyps in patients 18 years of age or older. The patient has had a documented side effect, allergy, or treatment failure of two preferred nasal glucocorticoids, one of which must be fluticasone.

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
			MC MC/DEL MC MC/DEL	8 8 8 8	VANCENASE POCKETHALER AERS <sup>2,3</sup> VERAMYST <sup>2,3</sup> XHANCE <sup>2</sup> ZETONNA <sup>3</sup>	4. Use of individual ingredients or other preferred agents.	
ANTIASTHMATIC - NASAL MISC.	MC/DEL MC/DEL MC	AZELASTINE CROMOLYN NASAL 4% IPRATROPIUM NASAL SOL <sup>1</sup>	MC/DEL MC/DEL	8 8	ASTEPRO <sup>2</sup> PATANASE	<a href="#">Use PA Form# 20420</a> 1. Ipratropium will be approved if submitted with documentation supporting use of CPAP machine. 2. Utilize Multiple preferred, as well as step therapy Azelastine.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  Approved if patient fails on nonsedating antihistamines and steroid nasal sprays.
ANTIASTHMATIC - BETA - ADRENERGICS	MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC	ALBUTEROL 0.63mg/3ml ALBUTEROL HFA ALBUTEROL NEB LEVALBUTEROL TARTRATE METAPROTERENOL PROAIR DIGITALER <sup>4</sup> PROAIR RESPICLICK PROVENTIL HFA SEREVENT STRIVERDI TERBUTALINE SULFATE TABS VENTOLIN HFA AERS	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC MC MC MC/DEL MC/DEL	8 8 8 8 4 8 8	ACCUNEB NEBU AIRSUPRA ALBUTEROL HFA (labeler 66993001968) BRETHINE VOLMAX TBCR VOSPIRE ER TB12 XOPENEX HFA <sup>3</sup> XOPENEX NEBU <sup>1,2</sup>	<a href="#">Use PA Form# 20420</a> 1. Xopenex users w/ prior asthma hospitalization due to albuterol nebulizer failure will be grandfathered.  2. Quantity Limit: 12 cc/day.. 3. Dosing limits apply, see Dosage Consolidation List. 4. For the treatment of patients $\geq$ 4 years of age.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  <b>AIRSUPRA</b> has new PA criteria that include the patient is aged $\geq$ 18, AND the patient has had a documented side effect or allergy, AND treatment failure/intolerance or contraindication to Symbicort® and Dulera® SMART therapy, AND the patient is unable to use albuterol and budesonide separately.
ANTIASTHMATIC - ADRENERGIC COMBINATIONS	MC MC MC MC MC/DEL MC/DEL	ADVAIR DISKUS <sup>1</sup> ADVAIR HFA <sup>1</sup> AIRDUO RESPICLICK <sup>2</sup> BREO ELLIPTA <sup>1</sup> DULERA FLUTICASONE-SALMETEROL SYMBICORT	MC MC/DEL MC/DEL MC	8 8 8 8	AIRDUO DIGITALER <sup>2</sup> BREYNA BREZTRI AEROSPHERE TRELEGY ELLIPTA <sup>1</sup>	<a href="#">Use PA Form# 20420</a> 1. Dosing limits apply, see Dosage Consolidation List.  2. For patients $\geq$ 12 years and older.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  <b>AirDuo® Respiclick</b> be non-preferred and require prior authorization and be available to those who are unable to tolerate or who have failed on preferred medications.  DDI: Avoid concomitant use of strong CYP3A4 inhibitors (e.g. ritonavir, atazanavir, clarithromycin, indinavir, itraconazole, nefazodone, neflifavir, saquinavir, ketoconazole, telithromycin) with <b>AirDuo® Respiclick</b> is not recommended due to increased systemic corticosteroid and increased cardiovascular adverse effects.
ANTIASTHMATIC - ADRENERGIC ANTICHOLINERGIC	MC/DEL MC MC/DEL MC/DEL	ALBUTEROL/IPRATROPIUM NEB. SOLN ANORO ELLIPTA COMBIVENT RESPIMAT STIOLTO	MC/DEL MC/DEL MC/DEL		BEVESPI AEROSPHERE <sup>2,3</sup> DUAKLIR PRESSAIR DUONEB SOLN <sup>1</sup>	<a href="#">Use PA Form# 20420</a> 1. Please use preferred individual ingredients Albuterol and Ipratropium. 2. Dosing limits apply, see Dosing Consolidation List. 3. The safety and efficacy of use in children under the age of 18 years have not been established.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  <b>Bevespi</b> should be used with extreme caution in patients being treated with MAO inhibitors, TCAs, or other drugs known to prolong the QTc interval. DDI: Avoid concomitant use of <b>Bevespi</b> with other anticholinergic-containing drugs, due to an increased risk of anticholinergic adverse events. Bevespi should be used with extreme caution in patients being treated with MAO inhibitors, TCAs, or other drugs known to prolong the QTc interval.
ANTIASTHMATIC - XANTHINES	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	AMINOPHYLLINE TABS THEOCHRON TB12 THEOLAIR-SR TB12 THEOPHYLLINE CR TB12 THEOPHYLLINE ELIX THEOPHYLLINE SOLN THEOPHYLLINE ER CP12 THEOPHYLLINE ER TB12	MC/DEL MC MC/DEL		THEO-24 CP24 THEOLAIR TABS UNIPHYL TBCR	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIASTHMATIC - STEROID INHALANTS	MC MC/DEL MC/DEL MC/DEL MC/DEL MC	ARNUITY ELLIPTA ASMANEX TWISTHALER <sup>3,4</sup> ASMANEX HFA BUDESONIDE NEB 0.25MG & 0.5MG <sup>1</sup> PULMICORT FLEXHALER <sup>3</sup> QVAR AERS <sup>3</sup>	MC MC/DEL MC MC/DEL MC/DEL	8 8 8 8 8	AEROSPA ALVESCO <sup>3</sup> ARMONAIR DIGITALER BUDESONIDE NEB 1MG PULMICORT SUSP	<a href="#">Use PA Form# 20420</a> 1. Budesonide Neb 0.25mg & 0.5mg will be preferred for members under the age of 8 years old. PA will be required for members 8 years of age and older, please consider other preferred options.  2. All preferred must be tried before moving to non preferred steps.  3. Dosing limits apply, see Dosage Consolidation List.  4. Asmanex 110mcg will be limited to member between the ages of 4-11 years old.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
ANTIASTHMATIC - 5-Lipoxygenase Inhibitors			MC		ZYFLO CR TABS	<a href="#">Use PA Form# 20420</a>	Other Preferred asthma controller drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIASTHMATIC - LEUKOTRIENE RECEPTOR ANTAGONISTS	MC/DEL MC/DEL MC/DEL	MONTELUKAST GRANULE <sup>1</sup> MONTELUKAST SODIUM TAB MONTELUKAST SODIUM CHEW TAB	MC/DEL MC/DEL MC/DEL	8 8 8	ACCOLATE TABS SINGULAIR <sup>2</sup> SINGULAIR GRANULES	<a href="#">Use PA Form# 20420</a> 1. Montelukast Granules will only be approved if between ages of 6 - 24 months. 2. Singulair Chewable 4mg from 2 years- 5 years and Singulair Chewable 5mgs from 6 years- 14 years old.	
ANTIASTHMATIC - ALPHA-PROTEINASE INHIBITOR			MC MC/DEL MC MC	8 8 8 8	ARALAST ZEMAIRA GLASSIA PROLASTIN SUSR	<a href="#">Use PA Form# 20420</a>	Prolastin and Azemaira will be approved for members with A1AT deficiency and clinically demonstrable panacinar emphysema.
ANTIASTHMATIC - HYDRO-LYTIC ENZYMES			MC/DEL		PULMOZYME SOLN	<a href="#">Use PA Form# 20420</a>	Will be approved for cystic fibrosis patients.
ANTIASTHMATIC - MUCOLYTICS	MC/DEL	ACETYLCYSTEINE <sup>1</sup>	MC		MUCOMYST	<a href="#">Use PA Form# 20420</a> 1. Acetylcysteine is covered with diagnosis of CF.	
ANTIASTHMATIC-CFTR POTENTIATOR AND COMBINATIONS			MC MC MC MC MC MC/DEL		ALYFTREK BRONCHITOL <sup>1</sup> KALYDECO ORKAMBI SYMDEKO TRIKAFTA	<a href="#">Use PA Form# 20420</a> 1. For the treatment of patients $\geq$ 18 years of age with CF.	<b>Alfytrek</b> will be considered for the treatment of patients 6 years and older with at least one responsive mutation, including 31 additional mutations not responsive to other CFTR modulator therapies.  <b>Bronchitol</b> will be considered as add-on maintenance therapy to improve pulmonary function in adult patients 18 years and older with cystic fibrosis (CF). Use Bronchitol only for adults who have passed the Bronchitol Tolerance Test (BTT). (see Recommended Dosage section for further information).  <b>Kalydeco</b> will be considered for patients with cystic fibrosis (CF) aged 1 month and older who have at least one mutation in the CFTR gene that is responsive to ivacaftor potentiation based on clinical and/or in vitro assay data. If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to detect the presence of a CFTR mutation followed by verification with bi-directional sequencing when recommended by the mutation test instructions for use.  <b>Orkambi</b> will be considered for patients with cystic fibrosis (CF) aged 1 year and older who are homozygous for the F508del mutation in the CFTR gene. If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to detect the presence of the F508del mutation on both alleles of the CFTR gene. The efficacy and safety of Orkambi have not been established in patients with CF other than those homozygous for the F508del mutation.  <b>Symdeko</b> will be considered for patients with cystic fibrosis (CF) aged 6 years and older who are homozygous for the F508del mutation or who have at least one mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to tezacaftor/ivacaftor based on in vitro data and/or clinical evidence. If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to detect the presence of a CFTR mutation followed by verification with bi-directional sequencing when recommended by the mutation test instructions for use.  <b>Trikafta</b> will be considered for the treatment of cystic fibrosis (CF) in patients aged 2 years and older who have at least one F508del mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene or mutation in the CFTR gene that is responsive based on in vitro data. If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to confirm the presence of at least one F508del mutation or a mutation that is responsive based on in vitro data.
IDIOPATHIC PULMONARY FIBROSIS	MC/DEL	OFEV <sup>1</sup> PIRFENIDONE	MC MC/DEL	8 8	ESBRIET <sup>1</sup> JASCAYD <sup>1</sup>	<a href="#">Use PA Form# 20420</a> 1. Clinical PA is required to establish diagnosis and medical necessity.	Ofev- Avoid concomitant use with P-gp and CYPA4 inducers (e.g. carbamazepine, phenytoin, and St. John's wort). Esbriet- The concomitant use with strong CYP1A2 inhibitors (e.g. fluvoxamine, enoxacin) is not recommended.
<b>COUGH/COLD</b>							
COUGH/COLD	MC/DEL MC/DEL MC/DEL MC/DEL MC MC	DEXTROMETHORPHAN CAPS <sup>1</sup> DEXTRO-GUAIF SYRP <sup>1</sup> GUAIFENESIN SYRP <sup>1</sup> PSEUDOEPHEDRINE <sup>1</sup> ROBITUSSIN DM SYRP <sup>1</sup> ROBITUSSIN SUGAR FREE SYRP <sup>1</sup>				<a href="#">Use PA Form# 20420</a> 1. All of cough cold preparations are not covered except these preferred products.	All non-preferred products are not covered as permitted by Federal Medicaid regulations and MaineCare Policy.
<b>DIGESTIVE AIDS / ASSORTED GI</b>							
GI - ANTIPERISTALTIC AGENTS	MC/DEL MC/DEL MC/DEL MC/DEL MC	DIPHENOXYLATE DIPHENOXYLATE/ATROPINE LOPERAMIDE HCL CAPS/LIQ OPIUM TINCTURE TINC PAREGORIC TINC	MC/DEL MC MC		LOFENE TABS LONOX TABS MOTOFEN TABS	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval.













Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required			Criteria
						3. Preferred stimulants will be available without PA if diagnosis of ADHD. 4. Dosing limits apply, see Dosing 5. For the treatment of Attention Deficit Hyperactivity Disorder (ADHD) in patients 7. FDA approval is currently for adults and children 6 or older. Will be available without PA for this age group if within dosing limits. Max dose of 50MG daily without a PA.  8. For the treatment of patients 6 years of age		
LONG ACTING AMPHETAMINES	MC MC/DEL MC	DEXTROAMPHET SULF CPSR <sup>1,3</sup> DEXTROAMPHETAMINE ER DYANAVEL XR SUS	MC/DEL MC MC MC MC		ADZENYS ER <sup>3</sup> ADZENYS XR- ODT ADZENYS XR <sup>3</sup> DEXEDRINE CAP SR <sup>2,3</sup> DYANAVEL XR TAB	<a href="#">Use PA Form# 20420</a>  1. Preferred stimulants will be available without PA if diagnosis of ADHD. 2. As per recent FDA alert, Adderall & Dexedrine should not be used in patients with underlying heart defects since they may be at increased risk for sudden death. 3. Dosing limits apply, see Dosing Consolidation List.	DDI: The concomitant use of <b>Adzenys XR</b> is contraindicated with monoamine oxidase inhibitors (MAOIs) or within 14 days after discontinuing MAOI treatment.	
STIMULANT - METHYLPHENIDATE	MC/DEL MC/DEL MC/DEL MC/DEL	DEXMETHYLPHENIDATE IR TABS METHYLPHENIDATE SOL METHYLPHENIDATE TAB METHYLIN TABS <sup>1,2</sup>	MC/DEL MC/DEL MC MC MC/DEL MC/DEL		FOCALIN IR TABS METADATE ER METHYLPHENIDATE HCL CHEW METHYLIN CHEWABLES METHYLIN SOL RITALIN	<a href="#">Use PA Form# 20420</a>  1. Preferred stimulants will be available without PA if diagnosis of ADHD. 2. Dosing limits apply, see Dosing Consolidation List. Maximum daily doses are as follows: 72mg daily for <b>Methylphenidate</b> and 36mg daily for <b>Dexmethylphenidate</b> .	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Please refer to General Criteria category E.	
STIMULANT - METHYLPHENIDATE - LONG ACTING	MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC MC/DEL	CONCERTA TBCR DEXMETHYLPHENIDATE CAP ER 50/50 FOCALIN XR METHYLPHENIDATE LA CAPS METHYLPHENIDATE ER CAPS 50/50 METHYLPHENIDATE ER CAPS 40/60 METHYLPHENIDATE CD CAPS 30-70 QUILLICHEW ER <sup>5,1</sup> QUILLIVANT XR SUS <sup>1,5</sup> RITALIN LA <sup>4</sup>	MC MC/DEL MC MC MC MC/DEL MC MC MC/DEL	5 8 8 8 8 8 8 8 8	METADATE CD CPCR ADHANSIA XR <sup>2,6</sup> APTENSIO XR <sup>2</sup> AZSTARYS <sup>6</sup> COTEMPLA XR <sup>2</sup> COTEMPLA XR ODT <sup>2</sup> DAYTRANA <sup>2,3</sup> JORNAY PM <sup>2,6</sup> METHYLPHENIDATE ER CAPS <sup>2,4</sup>	<a href="#">Use PA Form# 20420</a>  1. Preferred stimulants will be available without PA if diagnosis of ADHD. 2. Non-preferred products must be used in specified step order. 3. FDA approval currently only for ages 6-16. Limit of one patch daily. Max dose of 30MG daily. 4. Dosing limits apply, see Dosing Consolidation List. 5. Quillivant XR and Quillichew ER are only indicated for use in patients ≥ 6 years of age. 6. For the treatment of patients ≥ 6 years of age.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.	
STIMULANT - STIMULANT LIKE	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC	ATOMOXETINE HCL ARMODAFINIL CLONIDINE ER GUANFACINE ER MODAFINIL TABS QELBREE <sup>6,7</sup>	MC/DEL MC MC MC/DEL MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC MC	7 7 8 8 8 8 8 8 8 8 8 9 9 9 9	PROVIGIL TABS <sup>3</sup> STRATTERA <sup>1,2</sup> CAFCIT SOLN <sup>3</sup> INTUNIV KAPVAY ONYDA XR <sup>6</sup> SUNOSI WAKIX XYREM SOL XYWAV <sup>5</sup> NUVIGIL <sup>3</sup> DESOXYN TABS <sup>3</sup> DESOXYN CR <sup>3</sup>	<a href="#">Use PA Form# 20710 for Provigil, Nuvigil and Xyrem</a> <a href="#">Use PA Form# 20420 for all others</a>  1. Failure of both an amphetamine and methylphenidate is required for consideration for approval of <b>Strattera</b> , unless history of substance abuse without current use of abusive medication(s). Additionally, for patients <17 years of age, a trial of <b>Guanfacine</b> required before approval of <b>Strattera</b> . 2. Strattera currently has dosing limitations allowing one tablet per day for all strengths if obtain approval. Max daily dose of Strattera is 100mg. Please see Dosing Consolidation List. 3. Non-preferred products must be used in specified step order. 4. Please use generic Guanfacine. 5. For patients 7 years of age and older with narcolepsy. 6. For pediatric patients 6 years of age or older. 7. Preferred with a trial and fail either Atomoxetine OR any 2 preferred ADHD agents.	Provigil requests require diagnosis of Narcolepsy, ADHD, or Obstructive Sleep Apnea. Previous failures of methylphenidate and amphetamine is required for Narcolepsy and ADHD diagnosis, with additional Strattera trial needed with ADHD diagnosis. Please refer to detailed criteria on Provigil PA form.  <b>Sunsosi</b> is non-preferred and is indicated for to improve wakefulness in adult patients with excessive daytime sleepiness associated with narcolepsy or obstructive sleep apnea (OSA).  <b>Wakix</b> is non-preferred and is indicated for the treatment of excessive daytime sleepiness (EDS) in adults with narcolepsy.  <b>Xywav</b> : Diagnosis of cataplexy associated with narcolepsy OR excessive daytime sleepiness associated with narcolepsy. Diagnosis must be confirmed by submission of supporting documentation to include the specialist's interpretation of the Polysomnography (PSG) and Multiple Sleep Latency Test (MSLT) results.  FDA reminded healthcare professionals and patients that the combined use of <b>Xyrem</b> (sodium oxalate) with alcohol or central nervous system (CNS) depressant drugs can markedly impair consciousness and may lead to severe breathing problems (respiratory depression).  <b>DDI: Sunosi</b> is contraindicated with MAO inhibitors or within 14 days after discontinuing the MAO inhibitor.  <b>DDI: Concomitant use of Qelbree</b> with an MAO inhibitor or within 2 weeks after discontinuing an MAO inhibitor is contraindicated.  <b>DDI: Concomitant use of Qelbree</b> significantly increases the total exposure, but not peak exposure, of sensitive CYP1A2 substrates, which may increase the risk of adverse reactions associated with these CYP1A2 substrates. Coadministration of Qelbree with sensitive CYP1A2 substrates or CYP1A2 substrates with a narrow therapeutic range (e.g. alosetron, duloxetine, ramelteon, tasimelteon, tizanidine, theophylline), is contraindicated.	

Category	Coverage Indicator	Preferred Drugs		Coverage Indicator	Step Order	Non-Preferred Drugs PA Required				Criteria
<b>ANTI-CATALEPTIC AGENTS</b>										
PSYCHOTHERAPEUTIC AGENTS - MISC.				MC MC		NUDEXTA XENAZINE		<a href="#">Use PA Form# 20710 for Xenazine</a>		
<b>WEIGHT LOSS</b>										
WEIGHT LOSS						No longer covered: Phentermine, Xenical, Didrex, and Meridia		Weight loss drugs are not covered as permitted by Federal Medicaid regulations and Maine Medicaid (MaineCare) Policy.		
<b>ALZHEIMER DISEASE</b>										
ALZHEIMER - Cholinomimetics/Others	MC/DEL	DONEPEZIL HYDROCHLORIDE TABS <sup>1</sup>		MC	6	ARICEPT TABS <sup>2</sup>		<a href="#">Use PA Form# 20420</a>		Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL	DONEPEZIL HYDROCHLORIDE ODT <sup>1</sup>		MC	6	ARICEPT ODT <sup>2</sup>		1. PA is required to establish dementia diagnosis and baseline mental status score.		Kisunla and Leqembi: Testing to rule out reversible causes of dementia (CBC, CMP, TSH, B12, urine drug screen, RPR/VDRL, (folate if alcohol abuse is present), HIV (if risk present) and an assessment including a review of current medications as a cause of intellectual decline - Prescribed by or in consultation with a neurologist or geriatrician or geriatric psychiatrist. Diagnosis of Alzheimer's disease defined as: <ul style="list-style-type: none"> <li>Confirmed presence of amyloid pathology and mild cognitive impairment or mild dementia stage of disease, consistent with Stage 3 and Stage 4 Alzheimer's disease OR</li> <li>Confirmed presence of amyloid pathology and prodromal or mild dementia stage of disease, consistent with Stage 3 &amp; Stage 4 Alzheimer's disease</li> </ul> Testing: <ul style="list-style-type: none"> <li>Clinical Dementia Rating (CDR) global score of 0.5 or 1.0 OR</li> <li>Repeatable Battery for Assessment of Neuropsychological Status (RBANS) delayed memory index score ≤ 85 OR</li> <li>Mini-Mental State Examination (MMSE) score of 20-30 OR</li> <li>Montreal Cognitive Assessment (MoCA) score ≤ 22</li> </ul>
	MC/DEL	EXELON DIS <sup>1</sup>		MC/DEL	7	DONEPEZIL HYDROCHLORIDE TABS 23MG		2. Must fail all preferred products before moving to non-preferred.		
	MC/DEL	GALANTAMINE CAPS <sup>1</sup>		MC	8	ADALARITY <sup>3</sup>		3. Approvals will require trials and failure or clinical rationale why preferred patches can't be used.		
	MC/DEL	GALANTAMINE TAB <sup>1</sup>		MC/DEL	8	EXELON CAP				
	MC/DEL	MEMANTINE <sup>1</sup>		MC/DEL	8	GALANTAMINE HYDROBROMIDE SOL				
	MC/DEL	RIVASTIGMINE TARTRATE CAPS <sup>1</sup>		MC	8	KISUNLA <sup>1</sup>				
	MC/DEL			MC/DEL	8	LEQEMBI <sup>1</sup>				
	MC/DEL			MC/DEL	8	MEMANTINE HCL SOL				
	MC/DEL			MC/DEL	8	NAMENDA				
	MC/DEL			MC/DEL	8	NAMENDA XR CAPS				
	MC/DEL			MC	8	NAMZARIC				
	MC/DEL			MC	8	RAZADYNE <sup>2</sup>				
	MC/DEL			MC	8	ZUNVEYL				
	MC/DEL			MC	9	COGNEX CAPS <sup>2</sup>				
<b>SMOKING CESSATION</b>										
NICOTINE PATCHES / TABLETS	MC/DEL	CHANTIX TAB <sup>1</sup>		MC/DEL		NICODERM CQ PT24 <sup>1</sup>		<a href="#">Use PA Form# 20420</a>		As of July 1, 2014 per MaineCare policy, smoking cessation products will be covered without a copay(including MEDEL). No annual or lifetime limits, must follow FDA approved indications and therapy guidelines.
	MC/DEL	CHANTIX STARTER PACK						Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.		
	MC/DEL	NICOTINE DIS PT24 <sup>1</sup>						<b>Note:</b> MaineCare policy, smoking cessation product were "not covered" except for during pregnancy between 9/1/12 and 1/1/14, between 1/1/2014 and 7/1/14 smoking cessation products were covered with limitations.		
	MC/DEL	VARENICLINE TAB						Patients may qualify for the medication through The Maine Tobacco Helpline if they do not have MaineCare or MEDEL. Patients are encouraged to call The Maine Tobacco helpline at 1-800-207-1230.		
NICOTINE REPLACEMENT - OTHER	MC/DEL	NICOTINE POLACRILEX GUM <sup>1</sup>		MC/DEL	8	NICOTROL INHALER <sup>1,2</sup>		<a href="#">Use PA Form# 20420</a>		As of July 1, 2014 per MaineCare policy, smoking cessation products will be covered without a copay(including MEDEL). No annual or lifetime limits, must follow FDA approved indications and therapy guidelines.
	MC/DEL	NICOTINE LOZENGE MINI		MC/DEL	8	NICOTROL NASAL SPRAY <sup>1,2</sup>		1. See criteria section for exemptions.		Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL	NICOTINE LOZENGE		MC/DEL	8	NICORETTE GUM <sup>1,2</sup>		2. Must use non-preferred products in specified step order.		
	MC/DEL			MC	8	NICORETTE LOZENGES				
<b>ALCOHOL DETERRENTS</b>										
ALCOHOL DETERRENTS	MC/DEL	ACAMPROSATE		MC/DEL		ACAMPRO <sup>1</sup>		<a href="#">Use PA Form# 20420</a>		Preferred generic drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC	ANTABUSE TABS								
	MC	DISULFIRAM TABS								
	MC/DEL	NALTREXONE HCL TABS								
<b>MISCELLANEOUS ANALGESICS</b>										
ANALGESICS - MISC.	MC/DEL	ACETAMINOPHEN		MC		AXOCET CAPS		<a href="#">Use PA Form# 20420</a>		Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL	ASPIRIN		MC/DEL		ESGIC-PLUS		1. QL: 1. QL: No greater than 14-day supply within 90 days.		
	MC/DEL	ASPRIN/ APAP/ CAFF TAB		MC/DEL		FIORICET TABS				
	MC/DEL	BUTAL/ASA/CAFF		MC		FIORINAL CAPS				
	MC/DEL	BUTALBITAL COMPOUND		MC		FIORTAL CAPS				
	MC/DEL	BUTALBITAL/ACET TABS		MC/DEL		FORTABS TABS				

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
	<b>MC/DEL</b>	BUTALBITAL/APAP CAPS	<b>MC</b>		JOURNAVX <sup>1</sup>		
	<b>MC/DEL</b>	BUTALBITAL/APAP/CAFFEINE TABS	<b>MC</b>		PHRENILIN TABS		
	<b>MC/DEL</b>	CHOLINE MAGNESIUM TRISALI	<b>MC</b>		PHRENILIN FORTE CAPS		
	<b>MC/DEL</b>	DIFLUNISAL TABS	<b>MC</b>		TRILISATE LIQD		
	<b>MC</b>	EXCEDRIN	<b>MC</b>		TRILISATE TABS		
	<b>MC/DEL</b>	SALSALATE TABS	<b>MC</b>		ZEBUTAL CAPS		
	<b>MC</b>		<b>MC</b>		ZORPRIN TBCR		
<b>LONG ACTING NARCOTICS</b>							
NARCOTICS - LONG ACTING	<b>MC/DEL</b>	FENTANYL PATCH <sup>4</sup>	<b>MC</b>	8	ARYMO ER	<a href="#">Use PA Form# 20510</a>	Preferred drugs (Fentanyl Patch, Morphine Sulfate ER tab, and Butrans) must be tried for at least 2 weeks each & failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug & the preferred drug(s) exists. Adequate trials include prevention/treatment of common adverse effects associated w/ narcotics (antinausea, antipruritic, etc.) as well as adequate equianalgesic dosing when converting from one narcotic to another. Also, adequate documentation of attempts to titrate dose of preferred agents to achieve adequate pain relief & desired clinical response must be provided. Member's drug regimen for additions &/or discontinuations of medications that may affect absorption &/or metabolism of preferred agents must be monitored. Approvals will not be granted if patient had access to either non-preferred products or high doses of short acting narcotics during the trial period. Non-preferred drugs will not be approved for patients showing evidence of usage patterns consistent w/ controlled substance abuse such as:
	<b>MC/DEL</b>	BUTTRANS <sup>4</sup>	<b>MC</b>	8	AVINZA	<a href="#">Use PA Form #10300 for PAs over the opiate limit</a>	1. Frequent or persistent early refills of controlled drugs;
	<b>MC/DEL</b>	MORPHINE SULFATE ER TB12	<b>MC</b>	8	BELBUCA		2. Multiple instances of early refill overrides due to reports of misplacement, stolen, dropped in toilet or sink, distant travel, etc.
			<b>MC</b>	8	EXALGO		3. Breaches of narcotic contracts with any provider;
			<b>MC/DEL</b>	8	HYSINGLA ER		4. Failure to comply with patient responsibilities in attached opioid documentation (see PA form) including but not limited to failing to submit to and pass
			<b>MC</b>	8	KADIAN		5. Failing to take or pass random drug testing;
			<b>MC/DEL</b>	8	METHADONE <sup>6</sup>		6. Failing to provide old records regarding prior use of narcotics;
			<b>MC/DEL</b>	8	METHADOSE <sup>6</sup>		7. Receiving controlled substances from other prescribers that the provider submitting the PA is unaware of;
			<b>MC/DEL</b>	8	MORPHABOND ER		8. Documented history of substance abuse. Substance abuse evaluations may be required for patients with medical records displaying documented substance abuse or potential signs of narcotic misuse and abuse such as chronic early refills, short dosing intervals, frequent dose increases, multiple lost/stolen etc scripts and intolerance or "allergy" to all products but Oxycontin.
			<b>MC/DEL</b>	8	MORPHINE SULFATE ER CAP		9. Circumventing MaineCare prior authorization requirements for narcotics by paying cash for affected narcotics (prescribers failed to submit prior authorization prior to cash narcotic scripts being filled by member).
			<b>MC/DEL</b>	8	MORPHINE SULFATE SUPP		10. Requests for any Brand name controlled substance, considered by authorities to be highly abused and diverted (Oxycontin, Percocet, Tylox, Vicodin, Dilaudid, Ultracet...) with an available AB rated generic equivalent will be denied unless it will be provided in a setting that virtually eliminates the risk of diversion.
			<b>MC/DEL</b>	8	MS CONTIN TB12		11. Allergic reactions to any product within a specific narcotic class will justify and preclude use of any other product in the same class due to the risk of cross-hypersensitivity.
			<b>MC</b>	8	OPANA ER		<b>Hysingla ER</b> - Concomitant use should be avoided with mixed agonist/antagonist analgesics, partial agonist analgesics, and MAOIs. Verify prior trials and failures or intolerance of preferred treatments.
			<b>MC/DEL</b>	8	ORAMORPH SR TB12		<b>Methadone</b> - Established users must have a trial and failure of at least 2 preferred drugs for least 2 weeks. Otherwise they will be allowed 180 days to transition to a preferred product.
			<b>MC/DEL</b>	8	OXYCONTIN TB12 <sup>1</sup>		
			<b>MC</b>	8	XARTEMIS ER		
			<b>MC</b>	8	ZOHYDRO ER		
			<b>MC</b>	8	OXYCODONECONC		
			<b>MC/DEL</b>	9	OXYCODONE ER <sup>3,5</sup>		
NARCOTICS - SELECTED	<b>MC/DEL</b>	TRAMADOL HCL TABS 50 mg <sup>2</sup>	<b>MC/DEL</b>	7	RYZOLT	<a href="#">Use PA Form# 20420</a>	Preferred drugs from this and other narcotic classes must be tried for at least 2 weeks each and failed due to lack of efficacy or intolerable side effects before non-preferred drugs from this class will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Approvals will not be granted if patient had access to either non-preferred products or high doses of short acting narcotics during the trial period. Substance abuse evaluations may be required for patients with medical records displaying potential signs of narcotic misuse and abuse such as chronic early refills, short dosing intervals, frequent dose increases, multiple lost/stolen etc scripts and intolerance or "allergy" to all products but desired product. Allergic reactions to any product within a specific narcotic class will justify and preclude use of any other product in the same class due to the risk of cross-hypersensitivity.
	<b>MC/DEL</b>	TRAMADOL/APAP TABS	<b>MC</b>	8	BUPRENEX SOLN	<a href="#">Use PA form #10300 for PAs over the opiate limit</a>	Non-preferred drugs will not be approved for patients showing evidence of usage patterns consistent with controlled substance abuse such as:
			<b>MC/DEL</b>	8	BUTORPHANOL		1. frequent or persistent early refills of controlled drugs;
			<b>MC</b>	8	NALBUPHINE HCL SOLN		2. multiple instances of early refill overrides due to reports of misplacement, stolen, dropped in toilet or sink, distant travel;
			<b>MC</b>	8	QDOLO SOLN		3. breaches of narcotic contracts with any provider;
			<b>MC</b>	8	SEGLENTIS <sup>1</sup>		4. failure to comply with patient responsibilities in attached opioid documentation (see PA form) including but not limited to failing to submit to and pass pill counts;
			<b>MC</b>	8	STADOL NS SOLN		5. failing to take or pass random drug testing;
			<b>MC</b>	8	TRAMADOL ER		6. failing to provide old records regarding prior use of narcotics;
			<b>MC</b>	8	ULTRACET TABS <sup>1</sup>		7. receiving controlled substances from other prescribers that the provider submitting the PA is unaware of. Substance abuse evaluations may be required for patients with medical records displaying potential signs of narcotic misuse and abuse such as chronic early refills, short dosing intervals, frequent dose increases, multiple lost/stolen etc scripts and intolerance or "allergy" to all products but Oxycontin. Allergic reactions to any product within a specific narcotic class will justify and preclude use of any other product in the same class due to the risk of cross-hypersensitivity.
			<b>MC</b>	9	ULTRAM ER		Beginning January 2017, all current opiate users who are above the maximum combined daily dose of 100 MME must titrate their total daily dose of opioid medications below 30 MME. Also, the maximum daily supply of an opiate prescription for acute pain will be limited to 7-day supplies. The maximum day supply of an opiate prescription for chronic pain will be limited to 30-day supplies. As of July 1, 2017 all users of opioid medications must comply with the maximum combined daily dose of 100 MME.
							However, for MaineCare members, effective January 1, 2017, opioid prescription(s) for more than a 7-day supply and/or more than 30 MME/ day will require a prior authorization. Please note that MaineCare implemented a 30 MME limit January 1, 2013 that is still effective.
							Post-surgical members may receive prior authorizations for opiates up to 60 days in length if medical necessity is provided by the surgical provider.
							An MME conversion chart is available at <a href="http://www.maineicarepd.org">www.maineicarepd.org</a> . Click on "General Pharmacy Info."
							<b>Please see the Pain Management Policy tab for the complete criteria.</b>

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
<b>MISCELLANEOUS NARCOTICS</b>							
NARCOTICS - MISC.	MC/DEL	ACETAMINOPHEN/CODEINE	MC/DEL	8	ABSTRAL	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Please refer to General Criteria category E.
	MC/DEL	ASPIRIN/CODEINE TABS	MC/DEL	8	APADAZ	<a href="#">Use PA form #10300 for PAs over the opiate limit</a>	
	MC/DEL	BUTAL/ASA/CAFF/COD CAPS	MC/DEL	8	ASCOMP/CODEINE CAPS		
	MC	BUTALBITAL/ASPIRIN/CAFFE CAPS	MC/DEL	8	BUTALBITAL/APAP/CAFFEINE/ CAPS	1. Fentanyl OT loz (Barr) and Capital and Codeine Suspension products require PA for users over 18 years of age. PA is not required if under 18 years of age.	Beginning January 2017, all current opiate users who are above the maximum combined daily dose of 100 MME must titrate their total daily dose of opioid medications below 30 MME. Also, the maximum daily supply of an opiate prescription for acute pain will be limited to 7-day supplies. The maximum day supply of an opiate prescription for chronic pain will be limited to 30-day supplies. As of July 1, 2017 all users of opioid medications must comply with the maximum combined daily dose of 100 MME.
	MC	CAPITAL AND CODEINE SUSP <sup>1</sup>	MC/DEL	8	BUTALBITAL COMPOUND- CODEINE CAP		
	MC/DEL	CODEINE PHOSPHATE SOLN	MC	8	DEMEROL		
	MC/DEL	CODEINE SULFATE TABS	MC/DEL	8	DILAUDID		
	MC/DEL	ENDOCET TABS <sup>3</sup>	MC	8	DILAUDID-HP SOLN		
	MC/DEL	ENDODAN TABS	MC	8	FENTANYL CITRATE SOLN		
	MC/DEL	FENTANYL OT LOZ <sup>1</sup>	MC/DEL	8	FENTORA		
	MC/DEL	HYDROCODONE/ACETAMINOPHEN	MC/DEL	8	FIORICET/CODEINE CAPS		
	MC/DEL	HYDROMORPHONE HCL <sup>3</sup>	MC	8	FIORINAL/CODEINE #3 CAPS		
	MC	LORTAB ELX	MC	8	FIORTAL/CODEINE CAPS		
	MC/DEL	MEPERIDINE SOL	MC/DEL	8	HYDROCODONE/IBUPROFEN		
	MC/DEL	OXYCODONE TAB	MC/DEL	8	HYDROMORPHONE ER		
	MC/DEL	OXYCODONE/ACETAMINOPHEN <sup>2,3</sup>	MC/DEL	8	HYDROMORPHONE RECTAL SUPP		
	MC/DEL	ROXICET	MC	8	IBUDONE		
	MC	ROXIPRIN TABS	MC/DEL	8	LEVORPHANOL TARTRATE TAB		
			MC/DEL	8	LORCET		
			MC	8	LORTAB		
			MC	8	MAXIDONE TABS		
			MC/DEL	8	MEPERIDINE TABS		
			MC/DEL	8	NORCO TABS		
			MC/DEL	8	ONSOLIS		
			MC/DEL	8	OECTA		
			MC/DEL	8	OXYCODONE CAP		
			MC/DEL	8	OXYCODONE/APAP 10/650		
			MC/DEL	8	OXYCODONE/APAP 7.5/500		
			MC/DEL	8	PENTAZOCINE/ACET TABS		
			MC/DEL	8	PENTAZOCINE/NALOXONE TABS		
			MC	8	PERCOSET TABS		
			MC	8	PHRENILIN W/CAFFEINE/CODE CAPS		
			MC/DEL	8	ROXICET 5/500 TABS		
			MC	8	ROXICODONE TABS		
			MC/DEL	8	ROXYBOND		
			MC	8	SYNALGOS-DC CAPS		
			MC	8	TALACEN TABS		
			MC	8	TREZIX		
			MC	8	TYLENOL/CODEINE #3 TABS		
			MC	8	TYLOX CAPS		
			MC	8	XOLOX		
			MC	8	VICODIN		
			MC	8	VICOPROFEN TABS		
			MC	8	ZYDONE TABS		
			MC	9	ACTIQ LPOP		
			MC	9	CONZIP		
			MC	9	OPANA		
OPIOID DEPENDENCE TREATMENTS	MC	SUBOXONE FILM <sup>2</sup>	MC/DEL		BUPRENORPHINE <sup>1</sup>	<a href="#">Use PA form #20200 for Extended Release Buprenorphine</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL	BUPRENORPHINE/NALOXONE TABS <sup>2</sup>	MC		ZUBSOLV	<a href="#">Use PA Form #20100 for all others</a>	Members will continue to be required to follow the criteria listed below:
						1-Induction period for 30 days	
						2-Max dose of 32 mg for induction	
						3-Max dose of 24 mg for maintenance	
						4-There is not more than one opioid fill in member's drug profile between current fill of Buprenorphine and a prior Buprenorphine fill within the past 90 days	
						5- Should provide evidence of monthly monitoring including random pill counts, urine drug tests and use of Maine Prescription Monitoring Program reports.	
						6- Buprenorphine monotherapy is preferred if member is pregnant and dose not > 24 mg day and pregnancy diagnosis is noted on the prescription.	

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
EXTENDED RELEASE BUPRENORPHINE	MC MC	BRIXADI <sup>1</sup> SUBLOCADE <sup>1</sup>				<a href="#">Use PA form #20200 for Extended Release Buprenorphine</a> 1. Clinical PA required.	<b>Brixadi and Sublocade:</b> The prescriber can attest (and medical record should document) that: -member has a documented history of opioid use disorder (OUD), -XRB is being used for the treatment of OUD (rather than pain or any other non-FDA approved indication) and -member's total daily dose of sublingual buprenorphine is less than or equal to 24 mg daily. AND at least one of the following is true: -The member's previous use of sublingual buprenorphine has included misuse, overuse, or diversion. -The member is at high risk of overdose (e.g., individuals leaving incarceration or abstinence-based treatment programs; individuals who are unsheltered; or those facing potential gaps in care due to delays in care or geographically limited treatment access). -The member has experienced significant medical complications of OUD and/or injection drug use. Occurrence should be in the last 5 years, or it should be clearly documented that the risk indicated by this infection or complication is ongoing (Examples of medical complications of OUD include: threatened the function of organs or life or limb threatening and required medical and/or surgical therapy. Examples of medical complications of injection drug use include osteomyelitis, endocarditis, renal failure, joint infection or other serious medical complications directly related to OUD.) -The member has treatment-resistant OUD, including those with ongoing illicit substance use in the context of sublingual buprenorphine treatment as documented by positive urine drug screens or other clear objective evidence, and/or further functional decline with explicit documentation of the functional decline. -The member has a significant intolerance of, or documented allergy to, sublingual buprenorphine (either buprenorphine monotherapy or buprenorphine/naloxone combination therapy) that has resulted in the patient's inability to comply with continued treatment using the sublingual product. (A true allergy is usually accompanied by rash, respiratory symptoms, or anaphylaxis. Other complaints such as bad taste, mouth tingling, etc. do not constitute evidence of allergy or significant intolerance. Formulation preference or convenience are not, in and of themselves, indications for using XRB.) -The member is in ongoing treatment with XRB and would like to continue the medication.
OPIOID WITHDRAWAL AGENTS			MC		LUCEMYRA <sup>1</sup>	<a href="#">Use PA Form#20420</a> 1. Clinical PA for appropriate approved use and patient has documented contraindication to Clonidine.	
<b>NARCOTIC ANTAGONISTS</b>							
NARCOTIC - ANTAGONISTS	MC/DEL MC MC MC MC MC	NALTREXONE HCL TABS NALOXONE INJ NARCAN NS NALOXONE SPRAY OTC VIVITROL INJ ZIMHI	MC MC MC/DEL	8 8 8	OPVEE <sup>2</sup> KLOXXADO ZURNAL <sup>2</sup>	<a href="#">Use PA Form# 20420</a> 2. For the treatment of adult and pediatric patients 12 years of age and older.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
<b>COX 2 / NSAIDS</b>							
COX 2 INHIBITORS - SELECTIVE / HIGHLY SELECTIVE	MC/DEL MC/DEL MC/DEL MC/DEL	CELECOXIB <sup>4,5</sup> KETOROLAC TROMETHAMINE <sup>2,3,5</sup> NABUMETONE TABS <sup>5</sup> MELOXICAM TABS <sup>1,5</sup>	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC		CELEBREX CAPS <sup>4,5</sup> MELOXICAM CAPS <sup>5</sup> MOBIC <sup>5</sup> MOBIC SUSP <sup>5</sup> RELAFEN TABS <sup>5</sup> QMIIZ ODT VIVLODEX XIFYRM <sup>5</sup>	<a href="#">Use PA Form# 20420</a> 1. Meloxicam and Xifir have dosing limits allowing one tablet daily of all strengths without PA. 2. Ketorolac Tromethamine is indicated for the short term (up to 5 days) management of moderately severe acute pain that requires analgesic at the opioid level in adults. Not indicated for minor or chronic pain conditions. 3. Ketorolac has dosing limits allowing 24 tablets for a 5-day supply every 30 days. 4. Dosing limits will be set at a maximum of 400mg daily. 5. The FDA has issued a Public Health Advisory warning of the potential for increased cardiovascular risk & GI bleeding with NSAID use.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
NSAIDS	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	CHILDRENS IBUPROFEN DICLOFENAC POTASSIUM TABS DICLOFENAC SODIUM TABS DICLOFENAC SODIUM 1% GEL <sup>1</sup> ETODOLAC FENOPROFEN CALCIUM TABS FLURBIPROFEN TABS IBUPROFEN INDOMETHACIN KETOPROFEN	MC MC MC MC MC/DEL MC MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL		ADVIL TABS ANAPROX TABS ANAPROX DS TABS CAMBIA CATAFLAM TABS CHILDREN'S ADVIL SUSP CHILD'S IBUPROFEN SUSP CHILDREN'S MOTRIN SUSP CLINORIL TABS DAYPRO TABS	<a href="#">Use PA Form# 20420</a> The FDA has issued a Public Health Advisory warning of the potential for increased cardiovascular risk & GI bleeding with NSAID use. 1. Dosing limits apply, see Dosage Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  Approvals will be granted for other requests based on failure of at least one generic NSAID from at least 3 different NSAID classes as described in the COX-II PA form. The FDA has issued a Public Health Advisory warning of the potential for increased cardiovascular risk & GI bleeding with NSAID use. DDI: Diclofenac will now be non-preferred and require prior authorization if it is currently being used in combination with lescol.

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
	MC/DEL	MECLOFENAMATE SODIUM CAPS	MC/DEL		DICLFENAC GEL		
	MC/DEL	NAPROSYN SUSP	MC/DEL		EC-NAPROSYN TBEC		
	MC/DEL	NAPROXEN SUSP	MC/DEL		ETODOLAC ER 600MG		
	MC/DEL	NAPROXEN TABS	MC		FELDENE CAPS		
	MC/DEL	NAPROXEN SODIUM TABS	MC/DEL		FLECTOR PATCH		
	MC/DEL	NAPROXEN SODIUM CAPS	MC/DEL		IBU-200		
	MC/DEL	NAPROXEN DR TBEC	MC		INDOCIN		
	MC/DEL	OXAPROZIN TABS	MC		LICART		
	MC/DEL	SULINDAC TABS	MC/DEL		LODINE		
	MC/DEL	TOLMETIN SODIUM	MC		LOFENA		
	MC/DEL	VOLTAREN GEL	MC/DEL		MOTRIN		
			MC		NALFON CAPS		
			MC/DEL		NAPRELAN TBCR		
			MC/DEL		NAPROSYN TABS		
			MC/DEL		NAPROXEN SODIUM TBCR		
			MC		PENNSAID		
			MC/DEL		PIROXICAM CAPS		
			MC		PONSTEL CAPS		
			MC		RELAFEN DS		
			MC		SB IBUPROFEN TABS		
			MC		SPRIX		
			MC		TIVORBEX		
			MC		TOLECTIN		
			MC		V-R IBUPROFEN TABS		
			MC		ZORVOLEX		
NSAID - PPI			MC		PREVACID NAPRA-PAC	<a href="#">Use PA Form# 20420</a>	
			MC/DEL		VIMOVO <sup>1</sup>	1. Use a preferred NSAID and PPI separately.	
<b>RHEUMATOID ARTHRITIS</b>							
RHEUMATOID ARTHRITIS	MC/DEL	ACTEMRA VIALS	MC	8	AMJEVITA	<a href="#">Use PA Form# 20900</a>	See criteria as listed on Rheumatoid Arthritis PA form.
	MC/DEL	ACTEMRA SYRINGES	MC/DEL	8	ARAVA	1. Dosing limits apply, see Dosage Consolidation List.	Preferred injectable products allowed without PA if trial of a preferred oral agents (azathioprine, hydroxychloroquine, leflunomide, methotrexate, sulfasalazine tabs) are seen in the members drug profile. Dosing limits apply.
	MC/DEL	ADALIMUMAB-FKJP <sup>7</sup>	MC	8	AVTOZMA	2. Established users will be grandfathered.	<b>Jylamvo</b> will require using preferred methotrexate if unable please provide clinical rational as why inappropriate.
	MC	AVSOLA	MC/DEL	8	CIMZIA	3. Clinical PA is required to establish diagnosis and medical necessity.	<b>Xeljanz</b> is limited to adults with moderate to severe RA and UC who have had an inadequate response or intolerance to methotrexate. Should not be used concomitantly with biologic DMARDs or potent immunosuppressants.
	MC/DEL	AZATHIOPRINE	MC/DEL		CYLTEZO	4. Verification of age for appropriate indication.	<b>Zymfentra</b> : In adults for maintenance treatment of:
	MC	ENBREL <sup>2</sup>	MC/DEL		ENTYVIO	5. Treatment failure or intolerance to other forms of preferred methotrexate.	Moderately to severely active ulcerative colitis following treatment with an infliximab product administered intravenously.
	MC	ENBREL SURECLICK <sup>2</sup>	MC		HADLIMA	6. See criteria section.	Moderately to severely active Crohn's disease following treatment with an infliximab product administered intravenously.
	MC	HUMIRA <sup>1,2</sup>	MC/DEL		HULIO	7. Will require a clinical PA unless one systemic drug or a trial of a preferred oral agent (azathioprine, hydroxychloroquine, leflunomide, methotrexate, sulfasalazine tabs) are seen in the members drug profile. Dosing limits apply.	DDI: The concomitant use of <b>Xeljanz XR</b> with biologic DMARDs or potent immunosuppressants such as azathioprine and cyclosporine are not recommended. The concomitant use of <b>Xeljanz XR</b> with potent CYP3A4 inducers (e.g. rifampin) is not recommended.
	MC	KINERET SOLN	MC/DEL		HYDROXYCHLOROQUINE <sup>2</sup>	8. See additional criteria on the RA PA form.	
	MC/DEL	LEFLUNOMIDE	MC/DEL		HYRIMoz		
	MC/DEL	METHOTREXATE	MC/DEL		ILARIS <sup>1,3,4</sup>		
	MC	ORENICA	MC/DEL		INFLECTRA		
	MC/DEL	RINVOQ <sup>3</sup>	MC		INFILXIMAB VIAL		
	MC	SIMLANDI <sup>7</sup>	MC		JYLAMVO		
	MC	SIMPONI PEN	MC/DEL		KEVZARA		
	MC	SIMPONI AUTOINJECTOR	MC		OLUMIANT		
	MC/DEL	SULFASALAZINE TABS	MC		OMVOH		
	MC	TYENNE <sup>8</sup>	MC		OTREXUP		
	MC/DEL	XELJANZ <sup>3,6</sup>	MC		RASUVO <sup>6</sup>		
	MC/DEL	XELJANZ XR	MC		REMICADE		
			MC/DEL		RENFLEXIS		
			MC		TOFIDENCE		
			MC		VELSIPITY		
			MC/DEL	8	XELJANZ XR SOL		
			MC		XATMEP <sup>5</sup>		
			MC		YUFLYMA		
			MC		YUSIMRY		
			MC		ZYMFENTRA		



Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
MIGRAINE - PREVENTATIVE TREATMENT	MC MC/DEL MC/DEL MC/DEL MC/DEL	AIMOVIG <sup>1</sup> AJOVY <sup>1</sup> AJOVY AUTO INJECT <sup>1</sup> EMGALITY SYRINGE <sup>1</sup> 120mg/ml EMGALITY PEN <sup>1</sup> 120mg/ml	MC MC		QULIPTA VYEPTI <sup>2</sup>	<a href="#">Use PA Form# 10110</a> 1. See criteria section. 2. Dosing limits apply, see Dosage Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. <b>Aimovig, Ajoyv and Emgality:</b> The patient is 18 years of age or older AND patient has a diagnosis of episodic migraine (4-14 headache days per month with migraine lasting 4 hours or more) or chronic migraine ( $\geq$ 15 headache days per month, of which $\geq$ 8 are migraine days, for at least 3 months) AND patient has failed or has a contraindication to an adequate trial ( $\geq$ 60 days) of at least 2 medications for migraine prophylaxis from at least 2 different classes.
MIGRAINE - ACUTE TREATMENT	MC MC	SPASTRIN TABS UBRELVY <sup>1</sup>	MC MC MC/DEL MC/DEL MC MC MC/DEL		BELCOMP-PB SUPP ELYXYB MIGRAZONE CAPS MIGERGOT SUP NURTEC ODT REYVOW ZAVZPRET	<a href="#">Use PA Form# 10110</a> 1. Dosing limits apply, see Dosage Consolidation List.	Reyvow is non-preferred and is indicated for the acute treatment of migraine with or without aura in adults. Reyvow is not indicated for the preventive treatment of migraine. Ubrelvy is non-preferred and is indicated for the acute treatment of migraine with or without aura in adults. This is not indicated for the preventive treatment of migraine. <b>Zavzpret:</b> The patient must have a documented side effect, allergy, or treatment failure to preferred oral CGRP Inhibitor and two non-preferred oral CGRP Inhibitors.
<b>GOUT</b>							
GOUT	MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL	ALLOPURINOL TABS COLCHICINE TAB FEBUXOSTAT TAB MITIGARE PROBENECID TABS PROBENECID/COLCHICINE TABS	MC/DEL MC MC MC/DEL MC		COLCHICINE CAP COLCRYS GLOPERBA ULORIC <sup>1</sup> ZYLOPRIM TABS	<a href="#">Use PA Form# 20420</a> 1. Failure of therapeutic (300mg) dose of <b>Allopurinol</b> (failure define as not being able to get uric acid levels below 6mg/dl) or severe renal disease.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. <b>DDI:</b> The concomitant use of Gloperba and CYP3A4 inhibitors (e.g. clarithromycin, ketoconazole, grapefruit juice, erythromycin, verapamil, etc.) should be avoided due to the potential for serious and life-threatening toxicity.
<b>MISC.</b>							
ACID SPHINGOMYELINASE DEFICIENCY (ASMD)			MC		XENPOZYME <sup>1,2</sup>	<a href="#">Use PA Form# 20420</a> 1. For treatment of non-central nervous system manifestations of acid sphingomyelinase deficiency (ASMD) in adult and pediatric patients. 2. Clinical PA required for appropriate diagnosis and clinical parameters.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANESTHETICS - MISC.	MC MC MC	BUPIVACAINE HCL SOLN LIDOCAINE HCL SOLN MARCaine SOLN	MC MC/DEL MC		SENSORCAINE-MPF SOLN SYNVISC INJ XYLOCAINE SOLN	<a href="#">Use PA Form# 30130</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
COLD AGGLUTININ DISEASE (CAD)			MC		ENJAYMO <sup>1</sup>	<a href="#">Use PA Form# 20420</a> 1. Indicated to decrease the need for red blood cell transfusion due to hemolysis in adults with cold agglutinin disease (CAD).	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
CONGENITAL ADRENAL HYPERPLASIA			MC		CRENESSITY	<a href="#">Use PA Form# 30130</a>	<b>Creinessity</b> - As adjunctive treatment to glucocorticoid replacement to control androgens in adults and pediatric patients 4 years of age and older with classic congenital adrenal hyperplasia (CAH).
PRIMARY HYPEROXALURIA TYPE 1 (PH1)			MC MC/DEL		OXLUMO <sup>1</sup> RIVFLOZA	<a href="#">Use PA Form# 20420</a> 1. PA is required to establish diagnosis and medical necessity.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. <b>Rivfloza:</b> The patient has a diagnosis of Primary Hyperoxaluria Type I (PH1) confirmed via genetic testing (identification of alanine: glyoxylate aminotransferase gene (AGXT) mutation) AND urinary oxalate excretion $> 0.5\text{mmol}/1.73\text{m}^2$ or urinary oxalate: creatinine ratio is above the upper limit of normal for age AND is at least 9 years of age AND medication is being prescribed by, or in consultation, with a nephrologist or urologist.
SICKLE CELL DISEASE	MC MC MC/DEL MC	DROXIA CASGEVY <sup>2,3</sup> HYDROXYUREA LYFGENIA <sup>2,3</sup>	MC MC MC/DEL MC		ADAKVEO ENDARI <sup>1</sup> SIKLOS XROMI	<a href="#">Use PA Form# 20420</a> 1. Evidence of other preferred L-glutamine products utilization and reason for failure. 2. For the treatment of patients $\geq$ 12 years of age. 3. PA required to confirm FDA approved	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
HUTCHINSON- GILFORD PROGERIA SYNDROME (HGPS)			MC		ZOKINVY <sup>1,2</sup>	<a href="#">Use PA Form# 20420</a> 1. In patients 12 months of age and older with a body surface area (BSA) of 0.39m <sup>2</sup> and above. 2. PA required to confirm FDA approved indication.	<b>Zokinvy:</b> To reduce the risk of mortality in Hutchinson-Gilford Progeria Syndrome (HGPS). For the treatment of processing-deficient Progeroid Laminopathies with either: Heterozygous LMNA mutation with progerin-like protein accumulation OR Homozygous or compound heterozygous ZMPSTE24 mutations.
OBSTRUCTIVE SLEEP APNEA			MC		ZEPBOUND	<a href="#">Use PA Form# 20420</a>	<b>Zepbound</b> for adults with a BMI $\geq 30\text{ mg/kg}^2$ and diagnosis of moderate to severe OSA, confirmed by sleep study within the last 3 years documenting AHI $\geq 15$ , AND in which CPAP is ineffective (AHI $> 5$ during therapeutic section of sleep study) or patient is unable to tolerate CPAP for at least 90 days AND for whom lifestyle modifications have been attempted for at least 3 months with failure to achieve weight loss. <b>Note:</b> Not for patients with T1DM, T2DM.



Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
	<b>MC/DEL</b> <b>MC</b> <b>MC/DEL</b>	VALPROIC ACID SOL VALTOCO <sup>2</sup> ZONISAMIDE	<b>MC</b> <b>MC/DEL</b> <b>MC/DEL</b> <b>MC</b> <b>MC/DEL</b> <b>MC/DEL</b> <b>MC/DEL</b> <b>MC</b> <b>MC</b> <b>MC/DEL</b> <b>MC/DEL</b> <b>MC/DEL</b>	8 8 8 8 8 8 8 8 8 8 9 9 9 9	VIGAFYDE VIMPAT <sup>4</sup> VIMPAT SOL <sup>4</sup> XCOPRI ZARONTIN SYRP ZARONTIN CAP ZARONTIN SOL ZONISADE ZTALMY KEPPRA XR NEURONTIN TEGRETOL-XR TB12		
<b>ANTI-PARKINSON DRUGS</b>							
PARKINSONS - ANTICHOLINERGICS	<b>MC/DEL</b> <b>MC</b> <b>MC/DEL</b>	BENZTROPINE MESYLATE TABS COGENTIN SOLN TRIHEXYPHENIDYL				<a href="#">Use PA Form# 20420</a>	
PARKINSONS - ADENOSINE RECEPTOR ANTAGONIST			<b>MC/DEL</b>		NOURIANZ	<a href="#">Use PA Form# 20420</a>	Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  DDI: Avoid use of <b>Nourianz</b> with strong CYP3A4 inducers (e.g. carbamazepine, rifampin, phenytoin, St. John's wort).
PARKINSONS - COMT INHIBITORS			<b>MC/DEL</b> <b>MC</b>		COMTAN TABS ONGENTYS	<a href="#">Use PA Form# 20420</a>	Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
PARKINSONS - SELECTED DOPAMIN AGONISTS	<b>MC/DEL</b> <b>MC/DEL</b>	PRAMIPEXOLE ROPINIROLE NEUPRO PATCH	<b>MC/DEL</b> <b>MC</b> <b>MC/DEL</b>	5 8 8	MIRAPEX TABS <sup>1</sup> REQUIP TABS MIRAPEX ER	<a href="#">Use PA Form# 20420</a> 1. As of 12/08 users of Mirapex will be grandfathered if diagnosis is Parkinson's.	Preferred drug must be tried and failed in step-order due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
PARKINSONS- MAOIS			MC		XADAGO	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
PARKINSONS - DOPAMINERGICS/ CARBII/ LEVO	MC/DEL	AMANTADINE HCL CAPS	MC/DEL		APOKYN	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL	AMANTADINE HCL TABS	MC		AZILECT <sup>2</sup>		
	MC/DEL	BROMOCRIPTINE MESYLATE TABS	MC/DEL		CARBIDOPA/LEVODOPA RAPDIS		
	MC/DEL	BROMOCRIPTINE MESYLATE CAPS	MC		CREXONT <sup>4</sup>		
	MC/DEL	CARBIDOPA/LEVODOPA TABS <sup>3</sup>	MC		ELDEPRYL CAPS		
	MC/DEL	CARBIDOPA/LEVODOPA ER	MC		GOCOVRI		
	MC/DEL	CARBIDOPA/LEVO/ENTACAPONE TAB	MC/DEL		INBRIJA		
	MC	LARODOPA TABS	MC		KYNMOBI		
	MC/DEL	SELEGILINE CAPS HCL	MC/DEL		ONAPGO		
			MC/DEL		OSMOLEX ER		
			MC/DEL		PARLODEL CAPS		
			MC/DEL		PARLODEL TABS		
			MC		RYTARY		
			MC		SINEMET TABS		
			MC		SINEMET TBCR		
			MC		VYALEV		
PARKINSONS - COMBO.			MC/DEL		STALEVO <sup>1</sup>	<a href="#">Use PA Form# 20420</a>	
			MC		CARBIDOPA/LEVODOPA/ENTACA <sup>1</sup>	1. Clinical PA is required to establish diagnosis and medical necessity.	
<b>MUSCLE RELAXANTS</b>							
MUSCLE RELAXANTS	MC/DEL	BACLOFEN TABS	MC/DEL	7	ORPHENADRINE CITRATE	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL	CHLORZOXAZONE TABS	MC/DEL	8	CARISOPRODOL 350MG TABS		
	MC/DEL	CYCLOBENZAPRINE HCL 5mg & 10mg TABS	MC/DEL	8	AMRIX		
	MC	LIORESAL INTRATHECAL KIT	MC/DEL	8	DANTRIUM CAPS		
	MC/DEL	METHOCARBAMOL TABS	MC	8	FLEQSVUY		
	MC/DEL	TIZANIDINE HCL TABS	MC	8	LIORESAL TABS		
			MC	8	LORZONE		
			MC/DEL	8	LYVISPAN		
			MC	8	METAXALONE		
			MC	8	NORFLEX TBCR		
			MC	8	OZOBAX		
			MC	8	ROBAXIN-750 TABS		
			MC	8	VECUROMIUM INJ		
			MC/DEL	8	ZANAFLEX TABS		
			MC/DEL	9	CARISOPRODOL 250MG TABS		
			MC/DEL	9	CHLORZOXAZONE 250mg TABS		
			MC/DEL	9	SKELAXIN TAB		
			MC/DEL	9	SOMA TABS		
			MC	9	TANLOR		
MUSCLE RELAXANT - COMBO.			MC/DEL		CARISOPRODOL/ASPIRIN TABS	<a href="#">Use PA Form# 20420</a>	Individual components are available with PA described in the section above. 1. frequent or persistent early refills of non-controlled drugs; 2. multiple instances of early refill overrides due to reports of misplacement stolen, dropped in toilet or sink, distant travel, etc.
			MC/DEL		CARISOPRODOL/ASPIRIN/CODE		
			MC		NORGESIC TABS		
			MC/DEL		ORPHENADRINE COMPOUND		
			MC/DEL		ORPHENADRINE/ASA/CAFF		
			MC		ORPHENGESIC		
<b>PARATHYROID HORMONE</b>							
PARATHYROID HORMONE			MC		NATPARA <sup>1</sup>	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
			MC		YORVIPATH <sup>1</sup>		

Category	Coverage Indicator	Preferred Drugs		Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
<b>VITAMINS</b>								
VITAMINS	MC	CYANOCOBALAMIN SOLN	MC			AQUASOLE SOLN	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval.
	MC	FERIVA CAP	MC			AQUAVIT-E SOLN	<a href="#">Please refer to OTC list for covered products.</a>	
	MC	FERIVAF A CAP	MC			DHT SOLN		
	MC/DEL	FOLIC ACID TABS	MC			FUSION PLUS CAP		
	MC/DEL	MEPHYTON TABS	MC			HEMOCYTE PLU CAP		DDI: B-12 will now be non-preferred and require prior authorization if it is currently being used in combination with either prevacid, pantoprazole, prilosec, or any currently non preferred PPI.
	MC/DEL	NIACIN	MC			INTEGRA CAP		
	MC	NIACOR TABS	MC			INTEGRA F CAP		
	MC/DEL	NICOTINIC ACID SR CPCR	MC			INTEGRA PLUS CAP	<a href="#">Click here for the OTC List</a>	<b>Please refer to OTC list for covered products.</b>
	MC	PYRIDOXINE HCL TABS	MC			NASCOBAL GEL		Preferred products that used to require diag codes still require diag codes unless indicated otherwise.
	MC	TANDEM CAP	MC			TANDEM PLUS CAP		
	MC/DEL	THIAMINE HCL SOLN						
	MC/DEL	VITAMIN B-1 TABS						
	MC/DEL	VITAMIN B-12						
	MC	VITAMIN B-6 TABS						
	MC/DEL	VITAMIN C						
	MC/DEL	VITAMIN E CAPS						
	MC/DEL	VITAMIN E/D-ALPHA CAPS						
	MC	VITAMIN K1 SOLN						
	MC	V-R VITAMIN E CAPS						
VITAMIN D's	MC/DEL	CALCITRIOL CAPS <sup>1</sup>	MC			CALCIJEX	<a href="#">Use PA Form# 20420</a>	Preferred products require dialysis/renal failure diagnosis.
	MC/DEL	ROCALTROL	MC/DEL			DOXERCALCIF CAP		Rayaldee requires clinical PA to verify stage 3 or 4 CKD.
	MC/DEL	VITAMIN D2 <sup>2</sup>	MC/DEL			DOXERCALCIF INJ		
	MC/DEL	VITAMIN D3 <sup>2</sup>	MC/DEL			PARICALCITROL CAP		
	MC/DEL	VITAMIN DROPS	MC/DEL			PARICALCITROL INJ		
	MC	PARICALCITOL CAPS	MC/DEL			HECTOROL (ORAL)		
			MC			HECTOROL (PARENTERAL)		
			MC			RAYALDEE		
			MC			ZEMPLAR INJ		
			MC			ZEMPLAR CAPS		
<b>ENZYMES</b>								
POMPE DISEASE AGENTS			MC			NEXVIAZYME <sup>1</sup>	<a href="#">Use PA Form# 20420</a>	All preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
			MC			LUMIZYME		Pombili and Opfolda are for the treatment of adult patients with late-onset Pompe disease (lysosomal acid alpha-glucosidase [GAA] deficiency) weighing ≥40kg and
			MC			OPFOLDA		who are not improving on their current enzyme replacement therapy (ERT).
			MC			POMBILITI		
<b>MISC MULTI-VITAMINS</b>								
VITAMINS - MISC.	MC	CENTRUM TABS	MC			ADEKS	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval.
	MC	CENTRUM JR/IRON CHEW	MC/DEL			ADVANCED NATALCARE TABS		
	MC	CENTRUM-LUTEIN TABS	MC			AQUADEKS		
	MC	CEROVITE ADVANCED FO TABS	MC			CENTRUM JR/EXTRA C CHEW		
	MC/DEL	CHEWABLE MULTIVIT/FL CHEW	MC			CENTRUM PERFORMANCE TABS		
	MC	COD LIVER OIL CAPS	MC			CENTRUM SILVER TABS		
	MC/DEL	COMPLETE NATAL DHA (ORAL)	MC			DALYVITE LIQD		
		COMBO PKG	MC			EMBREX 600 MISC		
	MC	COMPLETE SENIOR TABS	MC			FERRALET 90		
	MC	DAILY MULTI VIT/IRON	MC			IBERET		
	MC/DEL	DIALYVITE 1MG	MC			MATERNA TABS		
	MC/DEL	DIALYVITE 800MG	MC			MAXARON		
	MC/DEL	FULL SPECTRUM B	MC			MULTIRET FOLIC -500 TBCR		
	MC	M.V.I-12 INJ	MC/DEL			NATAFORT TABS		
	MC	MULTI-VIT/FLUORIDE	MC/DEL			NATALCARE CFE 60 TABS <sup>1</sup>		
	MC/DEL	NATALCARE RX TABS	MC/DEL			NATALCARE GLOSS TABS <sup>1</sup>		
	MC/DEL	NEPHRONEX	MC			NATALCARE PIC TABS <sup>1</sup>		
	MC/DEL	NIVA-PLUS (ORAL) TABLET	MC			NATALCARE PIC FORTE TABS <sup>1</sup>		
	MC/DEL	ONE DAILY TABS	MC/DEL			NATALCARE PLUS TABS <sup>1</sup>		
	MC/DEL	ONE-DAILY MULTIVITAMINS	MC			NATALCARE THREE TABS <sup>1</sup>		
	MC/DEL	ONE-TABLET-DAILY	MC/DEL			NATACHEW CHEW		

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required			Criteria
	<b>MC/DEL</b>	POLY-VIT/IRON/FLUORID SOLN	<b>MC</b>		NATALFIRST TABS			
<b>MINERALS</b>								
	<b>MC/DEL</b>	POLY-VITAMIN/FLUORIDE SOLN	<b>MC</b>		NATATAB RX TABS			
	<b>MC/DEL</b>	POLY-VITAMINS/IRON SOLN	<b>MC/DEL</b>		NEPHPLEX RX TABS			
	<b>MC</b>	PRENATA (ORAL) TAB CHEW	<b>MC/DEL</b>		NEPHROCAPS CAPS			
	<b>MC/DEL</b>	PRENATAL TABS <sup>1</sup>	<b>MC/DEL</b>		NEPHRO-VITE TABS			
	<b>MC/DEL</b>	PRENATAL FORMULA 3 TABS <sup>1</sup>	<b>MC</b>		NESTABS RX TABS			
	<b>MC/DEL</b>	PRENATAL PLUS TABS <sup>1</sup>	<b>MC/DEL</b>		NIFEREX			
	<b>MC/DEL</b>	PRENATAL PLUS NF TABS <sup>1</sup>	<b>MC/DEL</b>		OCUVITE TABS			
	<b>MC</b>	PRENATAL PLUS/27MG IRON <sup>1</sup>	<b>MC</b>		POLY-VI-FLOR SOLN			
	<b>MC</b>	PRENATAL PLUS/IRON TABS <sup>1</sup>	<b>MC</b>		POLY-VI-SOL SOLN			
	<b>MC</b>	PRENATAL VITAMIN PLUS LOW IRON (ORAL) TABLET	<b>MC</b>		POLY-VI-SOL/IRON SOLN			
	<b>MC/DEL</b>	PRENATAL RX/BETA-CAROTENE <sup>1</sup>	<b>MC</b>		POLY-VITAMIN DROPS SOLN			
	<b>MC/DEL</b>	PREPLUS (ORAL) TABLET	<b>MC</b>		PRECARE			
	<b>MC/DEL</b>	RENAL CAPS	<b>MC</b>		PREFERA OB			
	<b>MC/DEL</b>	RENAPHRO CAPS	<b>MC</b>		PREMESIS RX TABS			
	<b>MC</b>	STRESS TAB NF TABS	<b>MC</b>		PRENATABS CBF TABS <sup>1</sup>			
	<b>MC</b>	THERAPEUTIC-M TABS	<b>MC</b>		PRENATAL CARE TABS <sup>1</sup>			
	<b>MC</b>	THERAVITE LIQD	<b>MC/DEL</b>		PRENATAL MR 90 TBCR <sup>1</sup>			
	<b>MC/DEL</b>	TRINATAL RX 1 (ORAL) TABLET	<b>MC</b>		PRENATAL MTR/SELENIUM TABS <sup>1</sup>			
	<b>MC/DEL</b>	TRIVEEN-DUO DHA (ORAL) COMBO. PKG	<b>MC</b>		PRENATAL OPTIMA ADVANCE TABS <sup>1</sup>			
	<b>MC/DEL</b>	TRI-VITAMIN/FLUORIDE SOLN	<b>MC/DEL</b>		PRENATAL PC 40 TABS <sup>1</sup>			
	<b>MC</b>	VITA CON FORTE CAPS	<b>MC</b>		PRENATAL RX TABS <sup>1</sup>			
	<b>MC</b>	VITAPLEX PLUS TABS	<b>MC</b>		PRENATE <sup>1</sup>			
			<b>MC</b>		PRENATE ELITE <sup>1</sup>			
			<b>MC</b>		PRIMACARE MISC			
			<b>MC</b>		PROTEGRA CAPS			
			<b>MC</b>		STUARTNATAL PLUS 3 TABS <sup>1</sup>			
			<b>MC</b>		TRI-VI-SOL SOLN			
			<b>MC</b>		TRI-VI-SOL/IRON SOLN			
			<b>MC/DEL</b>		ULTRA NATALCARE TABS			
			<b>MC</b>		ULTRA-NATAL TABS <sup>1</sup>			
			<b>MC</b>		VICON FORTE CAPS			
			<b>MC</b>		VINATAL FORTE TABS <sup>1</sup>			
			<b>MC</b>		VINATE <sup>1</sup>			
			<b>MC/DEL</b>		VINATE ADVANCED TABS <sup>1</sup>			
<b>MISCELLANEOUS MINERALS</b>								
	<b>MC</b>	CALCARB	<b>MC</b>		ANEMAGEN	<a href="#">Use PA Form# 20420</a>		
	<b>MC</b>	CALCI-MIX CAPSULE CAPS	<b>MC</b>		CALCET TABS	<a href="#">Please refer to OTC list.</a>		
	<b>MC</b>	CALCIQUID SYRP	<b>MC/DEL</b>		CALCIUM 600-D TABS			
	<b>MC</b>	CALCITRATE/VITAMIN D TABS	<b>MC</b>		CALCIUM/VITAMIN D TABS			
	<b>MC/DEL</b>	CALCIUM	<b>MC</b>		CALTRATE 600 PLUS/VIT D TABS			
	<b>MC/DEL</b>	CALCIUM CARBONATE	<b>MC</b>		CALTRATE PLUS TABS			
	<b>MC/DEL</b>	CALCIUM CITRATE TABS	<b>MC</b>		CHROMAGEN			
	<b>MC/DEL</b>	CALCIUM GLUCONATE TABS	<b>MC</b>		CITRACAL PLUS TABS			
	<b>MC/DEL</b>	CALCIUM LACTATE TABS	<b>MC</b>		CONTRIN CAPS			
	<b>MC</b>	CALCIUM/MAGNESIUM TABS	<b>MC</b>		FEOGEN FORTE CAPS			
	<b>MC/DEL</b>	CALCIUM/VITAMIN D TABS	<b>MC</b>		FEROCON CAPS			
	<b>MC</b>	CALTRATE 600 TABS	<b>MC/DEL</b>		FERREX 150 CAPS			
	<b>MC/DEL</b>	CHEWABLE CALCIUM CHEW	<b>MC</b>		FERRO-SEQUELS TBCR			
	<b>MC</b>	CITRACAL TABS	<b>MC</b>		FE-TINIC CAPS			
	<b>MC</b>	CITRACAL + D TABS	<b>MC</b>		FE-TINIC 150 FORTE CAPS			
	<b>MC</b>	CITRUS CALCIUM TABS	<b>MC/DEL</b>		FLUOR-A-DAY SOLN			
	<b>MC</b>	CITRUS CALCIUM 1500 + D TABS	<b>MC</b>		HEMOCYTE TABS			
	<b>MC</b>	EFFERVESCENT POTASSIUM TBDF	<b>MC/DEL</b>		K-DUR TBCR			
	<b>MC/DEL</b>	FEOSTAT CHEW	<b>MC</b>		KLOR-CON PACK			
	<b>MC</b>	FERATAB TABS	<b>MC</b>		K-LYTE			
	<b>MC/DEL</b>	FER-GEN-SOL SOLN	<b>MC/DEL</b>		K-PHOS TABS NEUTRAL			
	<b>MC</b>	FER-IRON SOLN	<b>MC</b>		K-TABS TBCR			
	<b>MC</b>	FERRONATE TABS	<b>MC</b>		K-VESCENT PACK			



Category	Coverage Indicator	Preferred Drugs		Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
<b>MISC. ELECTROLYTES/NUTRITIONALS</b>								
ELECTROLYTES/ NUTRITIONALS	<b>MC</b>	INTRALIPID EMUL <sup>1</sup>	<b>MC</b>		BOOST <sup>1</sup>	<a href="#">Use PA Form# 20420 &amp; SGA Form</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval.	
	<b>MC</b>	P.T.E. -5 SOLN <sup>1</sup>	<b>MC</b>		CASEC POWD <sup>1</sup>		Medical foods are not to be authorized solely for the purpose of enhancing nutrient intake or managing body weight if the participant is able to eat conventional foods adequately. Medical foods may be approved if the member has a medical condition which precludes or restricts the use of conventional foods and necessitates the use of a formula. Concurrent Stimulant therapy is not an acceptable medical reason/condition for use of medical foods for enhancing nutrient intake or managing body weight.	
	<b>MC</b>	SEA-OMEGA CAPS <sup>1</sup>	<b>MC</b>		CHOICE DM LIQD <sup>1</sup>			
	<b>MC</b>		<b>MC</b>		DELIVER 2.0 LIQD <sup>1</sup>			
	<b>MC</b>		<b>MC</b>		DOJOLVI			
	<b>MC</b>		<b>MC</b>		ENFAMIL <sup>1</sup>			
	<b>MC</b>		<b>MC</b>		ENSURE <sup>1</sup>			
	<b>MC</b>		<b>MC</b>		GLUCERNA <sup>1</sup>			
	<b>MC</b>		<b>MC</b>		ISOCAL LIQD <sup>1</sup>			
	<b>MC</b>		<b>MC</b>		KINDERCAL TF LIQD <sup>1</sup>			
	<b>MC</b>		<b>MC</b>		KINDERCAL TF/FIBER LIQD <sup>1</sup>			
	<b>MC</b>		<b>MC</b>		L-CARNITINE CAPS <sup>1</sup>			
	<b>MC</b>		<b>MC</b>		LIPISORB LIQD <sup>1</sup>			
	<b>MC</b>		<b>MC</b>		LOVAZA <sup>1,2</sup>			
	<b>MC</b>		<b>MC</b>		MODULEN IBD POWD <sup>1</sup>			
	<b>MC</b>		<b>MC</b>		NUTRAMIGEN POWD <sup>1</sup>			
	<b>MC</b>		<b>MC</b>		NUTREN <sup>1</sup>			
	<b>MC</b>		<b>MC</b>		NUTRITIONAL SUPPLEMENT LIQD <sup>1</sup>			
	<b>MC</b>		<b>MC</b>		NUTRIVENT 1.5 LIQD <sup>1</sup>			
	<b>MC</b>		<b>MC</b>		PEPTAMEN <sup>1</sup>			
	<b>MC</b>		<b>MC</b>		PHENYLADE <sup>1</sup>			
	<b>MC</b>		<b>MC</b>		PHENYL-FREE <sup>1</sup>			
	<b>MC</b>		<b>MC</b>		PKU 3 POWD <sup>1</sup>			
	<b>MC</b>		<b>MC</b>		PREGESTIMIL POWD <sup>1</sup>			
	<b>MC</b>		<b>MC</b>		PROBALANCE LIQD <sup>1</sup>			
	<b>MC</b>		<b>MC</b>		PROSOBEE <sup>1</sup>			
	<b>MC</b>		<b>MC</b>		SCANDISHAKE PACK <sup>1</sup>			
	<b>MC</b>		<b>MC</b>		VASCEPA			
ERYTHROPOEITINS	<b>MC</b>	EPOGEN SOLN	<b>MC</b>	8	ARANESP SOLN <sup>1</sup>	<a href="#">Use PA Form# 10520</a>	Non-Preferred drugs must be tried and failed in step-order, due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Please see the EPO PA form for other approval and renewal criteria.	
	<b>MC</b>	MIRCERA SYRINGE	<b>MC</b>	8	PROCRIT SOLN <sup>1</sup>			
	<b>MC</b>	RETACRIT						
<b>GRANULOCYTE CSF</b>								
GRANULOCYTE CSF	<b>MC</b>	FULPHILA	<b>MC</b>	8	FYLNETRA	<a href="#">Use PA Form# 20520</a>	See approval criteria detailed on Granulocyte Colony Stimulating Factor PA form.	
	<b>MC</b>	NEUPOGEN SYRINGE	<b>MC</b>	8	GRANIX SYRINGE			
	<b>MC</b>	NEUPOGEN VIAL	<b>MC</b>	8	GRANIX VIAL			
	<b>MC/DEL</b>	NYVEPRIA SYRINGE	<b>MC</b>	8	LEUKINE			
			<b>MC/DEL</b>	8	NIVESTYM			
			<b>MC</b>	8	ROLVEDON			
			<b>MC</b>	8	RYZNEUTA			
			<b>MC</b>	8	STIMUFEND			
			<b>MC/DEL</b>	8	ZARXIO			
			<b>MC/DEL</b>	8	ZIEXTENZO			
			<b>MC</b>	9	NEULASTA <sup>1</sup>			
<b>GAUCHER DISEASE</b>								
GAUCHER DISEASE			<b>MC</b>		CERDELGA <sup>1</sup>	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Exceeding days supply limits for LMWH class requires PA.	
			<b>MC</b>		YARGESA <sup>1</sup>			
							<b>Yargesa:</b> As monotherapy for the treatment of adult patients with mild to moderate type 1 Gaucher disease for whom enzyme replacement therapy is not a therapeutic option (e.g., due to allergy, hypersensitivity, or poor venous access).	
<b>NIEMANN-PICK DISEASE AGENTS</b>								
NIEMANN-PICK DISEASE AGENTS			<b>MC</b>		AQNEURSA <sup>1</sup>	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.	
			<b>MC</b>		MIPLYFFA <sup>1</sup>			

Category	Coverage Indicator	Preferred Drugs		Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
<b>ANTICOAGULANTS / PLATELET AGENTS</b>								
ANTICOAGULANTS		<b>MC</b> COUMADIN TABS <b>MC/DEL</b> ENOXAPARIN <sup>1</sup> <b>MC</b> ELIQUIS <b>MC</b> ELIQUIS STARTER PACK <b>MC</b> HEPARIN SODIUM/NACL 0.9% SOLN <b>MC</b> HEP-LOCK SOLN <b>MC</b> INNOHEP <b>MC</b> HEPARIN LOCK SOLN <b>MC/DEL</b> HEPARIN LOCK FLUSH SOLN <b>MC/DEL</b> HEPARIN SODIUM SOLN <b>MC/DEL</b> HEPARIN SODIUM LOCK FLUSH SOLN <b>MC/DEL</b> PRADAXA <b>MC/DEL</b> JANTOVEN <b>MC/DEL</b> WARFARIN SODIUM TABS <b>MC/DEL</b> XARELTO <b>MC/DEL</b> XARELTO STARTER PACK		<b>MC</b> ARIXTRA SOLN <b>MC/DEL</b> FONDAPARINUX <b>MC/DEL</b> FRAGMIN INJ <b>MC/DEL</b> FRAGMIN VIAL <b>MC</b> LOVENOX SOLN <b>MC/DEL</b> LOVENOX 300 <sup>2</sup> <b>MC/DEL</b> LOVENOX SUBQ SYRINGE <b>MC/DEL</b> PRADAXA ORAL PELLETS <sup>4</sup> <b>MC</b> IPRIVASK <b>MC/DEL</b> SAVAYSAS <sup>3</sup>		<a href="#">Use PA form# 20420</a> 1. Enoxaparin therapy durations greater than 7 days every 30 days require PA. 2. Use other strengths available to obtain desired dose. 3. Diagnosis required. 4. For the treatment of patients aged 3 months to less than 12 years of age.		Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Exceeding days supply limits for LMWH class requires PA.  <b>DDI:</b> Warfarin will require prior authorization if being used in combination with fluconazole, miconazole, or voriconazole. <b>DDI:</b> Warfarin will require prior authorization if being used in conjunction with gemfibrozil or fenofibrate. <b>DDI:</b> Rifampin will require prior authorization if being used in combination with savaysa.
ANTIHEMOPHILIC AGENTS		<b>MC</b> ALPHANATE <b>MC</b> ALPHANINE SD <b>MC/DEL</b> ALPROLIX VIAL <b>MC/DEL</b> BEBULIN VIAL <b>MC/DEL</b> BENEFIX SOLR <b>MC/DEL</b> HELIXATE FS KIT <b>MC</b> HEMOFIL - M <b>MC</b> HUMATE-P SOLR <b>MC/DEL</b> IXINITY VIAL <b>MC/DEL</b> JIV <sup>3</sup> <b>MC</b> KOATE-DVI <b>MC</b> KONYNE - 80 <b>MC/DEL</b> KOVALTRY <b>MC/DEL</b> REBINYN <b>MC</b> MONARC - M <b>MC</b> MONOCLOATE - P <b>MC</b> MONONINE <b>MC/DEL</b> NOVOEIGHT <b>MC</b> NOVOSEVEN SOLR <b>MC</b> NUWIQ <b>MC/DEL</b> PROFILNINE <b>MC</b> RECOMBIMATE SOLR <b>MC</b> REFACTO <b>MC/DEL</b> RIXUBIS VIAL <b>MC</b> WILATE INJ <b>MC/DEL</b> XYNTHA		<b>MC/DEL</b> ADYNOVATE VIAL <b>MC</b> ADVATE <sup>1,2,5</sup> <b>MC</b> ALTUVIIO <sup>4</sup> <b>MC/DEL</b> AFSTYLA <b>MC/DEL</b> BEQVEZ <b>MC/DEL</b> ESPEROCT <b>MC/DEL</b> ELOCTATE <b>MC/DEL</b> HEMGENIX <b>MC/DEL</b> IDELVION <b>MC/DEL</b> KOGENATE FS <sup>5</sup> <b>MC</b> RECOMBIMATE VIAL <sup>5</sup> <b>MC</b> ROCTAVIAN <sup>4</sup> <b>MC</b> SEVENFACT		<a href="#">Use PA Form# 20420</a> 1. Only if other products unavailable. 2. Advate may be available with PA in cases of large volume dosing in patients with poor venous access. 3. Not indicated for use in children <12 years of age due to greater risk for hypersensitivity reactions and is not indicated for use in previously untreated patients. 4. Clinical PA required for appropriate diagnosis. 5. Established users will be grandfathered.		Non-preferred will only be approved if other preferred products are unavailable.  <b>Beqvez</b> - FDA Approved Indication: An adeno-associated virus vector-based gene therapy indicated for the treatment of adults with moderate to severe hemophilia B (congenital factor IX deficiency) who: <ul style="list-style-type: none"> <li>• Currently use factor IX prophylaxis therapy, or</li> <li>• Have current or historical life-threatening hemorrhage, or</li> <li>• Have repeated, serious spontaneous bleeding episodes, and,</li> <li>• Do not have neutralizing antibodies to adeno-associated virus serotype Rh74var (AAVRh74var) capsid as detected by an FDA-approve test.</li> </ul> <b>Hemgenix</b> is an adeno-associated viral vector-based gene therapy for IV infusion after dilution. For treatment of adults with Hemophilia B (congenital Factor IX deficiency) who: Currently use Factor IX prophylaxis therapy, or have current or historical life-threatening hemorrhage, or have repeated, serious spontaneous bleeding episodes.  <b>Altuviio</b> is a von Willebrand Factor (VWF) independent recombinant DNA-derived, Factor VIII concentrate indicated for use in adults and children with hemophilia A (congenital factor VIII deficiency) for: Routine prophylaxis to reduce the frequency of bleeding episodes, On-demand treatment and control of bleeding episodes, Perioperative management of bleeding.  <b>Roctavian</b> : For the treatment of adults with severe hemophilia A (congenital factor VIII deficiency with factor VIII activity <1 IU/dL) without antibodies to adeno-associated virus serotype 5 (AAV5) detected by an FDA-approved test.  <b>Inclusion:</b> <ul style="list-style-type: none"> <li>• Severe factor VIII deficiency (less than 1% native factor VIII).</li> </ul> <b>Exclusion Criteria:</b> <ul style="list-style-type: none"> <li>• Antibodies to the virus AAV5</li> <li>• Factor VIII inhibitors (or history of)</li> <li>• Known significant fibrosis of cirrhosis of the liver, or unexplained elevated LFTs</li> <li>• History of inadequate compliance with prophylaxis, or regular bleeds despite adequate prophylaxis</li> <li>• Conditions in which high-dose steroids are contraindicated.</li> <li>• Inability to abstain from alcohol for one year</li> <li>• Plan to impregnate a partner within 6 months of infusion</li> <li>• Hypersensitivity to mannitol</li> <li>• Active infections, either acute or uncontrolled chronic</li> <li>• HIV infection (limited information on use in this population)</li> </ul>
NON-FACTOR REPLACEMENT THERAPY		<b>MC</b> HEMLIBRA		<b>MC/DEL</b> <b>MC/DEL</b> <b>MC</b> <b>MC</b>		ALHEMO HYMPAVI QFITLIA QFITLIA PEN		<a href="#">Use PA Form# 20420</a> Subsequent changes made to Antihemophilic Agents: Factor Therapy to move Hemlibra to Non-Factor Therapy
PLATELET AGGREGATION INHIBITORS		<b>MC/DEL</b> ASPIRIN <b>MC</b> ASPIRIN-DIPYRIDAMOLE ER CPMP 12HR <b>MC/DEL</b> BRILINTA 90mg <b>MC/DEL</b> DIPYRIDAMOLE TABS <b>MC/DEL</b> CLOPIDOGREL 75MG <b>MC/DEL</b> PRASUGREL HCL TAB		<b>MC/DEL</b> 7 TICLOPIDINE HCL TABS <b>MC/DEL</b> 8 BRILINTA 60mg <b>MC</b> 8 DURLAZA <b>MC</b> 8 EFFIENT <b>MC/DEL</b> 8 PERSANTINE TABS <b>MC/DEL</b> 8 PLAVIX TABS <b>MC/DEL</b> 8 ZONTIVITY		<a href="#">Use PA Form# 20715 for Plavix, Effient &amp; Brilinta</a> <a href="#">Use PA form# 20420 for other requests</a> 1. Dosing limits apply, see Dose Consolidation List.		Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  A special PA may be obtained at the pharmacy for members scheduled for "stent" placement or have had placement if in the last 12months. Please indicate on prescription date of stent placement.  <b>Brilinta</b> - Concomitant use with strong CYP3A4 inhibitors should be avoided (including ketoconazole, itraconazole, atazanavir, and telithromycin). Doses of simvastatin and lovastatin >40mg should be avoided. <b>DDI:</b> exists for using maintenance ASA dose >100mg, as it reduces the effectiveness of Brilinta. <b>DDI:</b> Plavix will require prior authorization if being used in combination with omeprazole, esomeprazole, cimetidine, fluconazole, ketoconazole, intelence, fluoxetine, ticlopidine, and fluvoxamine.

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
PLATELET AGGR. INHIBITORS / COMBO'S - MISC.	MC/DEL MC/DEL	CILOSTAZOL PENTOXIFYLLINE ER TBCR	MC/DEL MC/DEL MC/DEL MC MC		AGRYLIN CAPS ANAGRELIDE CAPS PLETAL TABS TRENTAL TBCR YOSPRALA	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
<b>HEMATOLOGICALS</b>							
MONOCLONAL ANTIBODY			MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC		BKEMV EMPAVELI ENSPRYNG EPYSQLI FABHALTA GAMIFANT PIASKY SOLIRIS ULTOMIRIS UPLIZNA VOYDEYA	<a href="#">Use PA Form# 20420</a>	A diagnosis of Paroxysmal nocturnal hemoglobinuria (PNH) using the HAM test or flow cytometry is required. In addition, the patient must show evidence of having received a meningitis vaccine at least 2 weeks prior to the start of therapy.  <b>Gamifant</b> is recommended for the treatment of adult and pediatric (newborn and older) patients with primary hemophagocytic lymphohistiocytosis (HLH) with refractory, recurrent, or progressive disease or intolerance with conventional HLH therapy.  <b>Fabhalta</b> and <b>Ultomiris</b> are recommended for the treatment of adults with paroxysmal nocturnal hemoglobinuria (PNH).  <b>Bkemv</b> and <b>Epysqli</b> have updated criteria for a diagnosis of generalized myasthenia gravis (gMG): must have confirmation that patients are anti-acetylcholine receptor (AChR) antibody positive.
IMMUNE GLOBULIN	MC MC/DEL MC MC/DEL MC/DEL MC/DEL MC	BIVIGAM <sup>1</sup> CUTAQUIG <sup>1</sup> GAMMAGARD S-D <sup>1</sup> HIZENTRA <sup>1</sup> PANZYGA <sup>1</sup> PRIVIGEN <sup>1</sup>	MC MC MC/DEL MC MC/DEL MC MC/DEL		ALYGLO ASCENIV <sup>2</sup> CUVITRU GAMMAPLEX INJ HYQVIA OCTAGAM INJ <sup>1</sup> XEMBIFY	<a href="#">Use PA Form# 20420</a> 1. Clinical PA required. 2. For the treatment of patients between 12 to 17 years of age.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  <b>Alyglo</b> is indicated for treatment of primary humoral immunodeficiency in adults ages 17 or older. <b>Cutaquig</b> is indicated as replacement therapy for primary humoral immunodeficiency (PI) in adults. <b>Xembify</b> is indicated for treatment of primary humoral immunodeficiency (PI) in patients 2 years of age and older. <b>Asceniv</b> indicated for the treatment of primary humoral immunodeficiency (PI) in adults and adolescents (12 to 17 years of age). PI includes but is not limited to the humoral immune defect in congenital agammaglobulinemia, common variable immunodeficiency (CVID), X-linked agammaglobulinemia, Wiskott-Aldrich syndrome, and severe combined immunodeficiencies (SCID).
HEREDITARY ANGIOEDEMA	<b>PROPHYLAXIS</b>		<b>PROPHYLAXIS</b>		<a href="#">Use PA Form# 20420</a>		Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  <b>Haegarda</b> is indicated for routine prophylaxis to prevent Hereditary Angioedema (HAE) attacks in adolescent and adult patients
	MC MC MC MC/DEL	CINRYZE <sup>1</sup> HAEGARDA <sup>1</sup> ORLADEYO <sup>1,2</sup> TAKHYRO <sup>1</sup>	MC MC	8 8	ANDEMBRY DAWNZERA <sup>2</sup>	1. Clinical PA is required to establish diagnosis and medical necessity. 2. For the treatment of patients $\geq$ 12 years of age	
HEMATOLOGICAL AGENTS-THROMBOPOIETIN RECEPTOR AGONISTS	<b>TREATMENT</b>		<b>TREATMENT</b>		<a href="#">Use PA Form# 20420</a>		<b>Doptelet</b> and <b>Mulpelta</b> : For the treatment of thrombocytopenia in adults with chronic liver disease who are scheduled to undergo a procedure.
	MC/DEL MC MC/DEL	BERINERT KIT <sup>1</sup> FIRAZYR <sup>1</sup> RUCONEST VIAL <sup>1</sup>	MC/DEL MC	8 8	KALBITOR VIAL EKTERLY <sup>2</sup>	1. Clinical PA is required to establish diagnosis and medical necessity. 2. For the treatment of patients $\geq$ 12 years of age	
HEMATOLOGICAL AGENTS-IgAN	MC MC	PROMACTA <sup>1</sup> NPLATE <sup>1</sup>	MC MC/DEL MC/DEL		ALVAIZ DOPTELET MULPLETA	<a href="#">Use PA Form# 20420</a> 1. Clinical PA required. Must see prior trial with insufficient response to corticosteroids and immunoglobulins.	<b>Doptelet</b> and <b>Mulpelta</b> : For the treatment of thrombocytopenia in adults with chronic liver disease who are scheduled to undergo a procedure.  All preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists  PA required to confirm FDA-approved indication. <b>Vanrafia</b> is for adults with biopsy proven primary IgAN AND eGFR $\geq$ 30 cc/min/1.73m <sup>2</sup> AND urine protein $\geq$ 1 g/day AND on stable dose of maximally tolerated renin-angiotensin system inhibitor.
ANEMIA- BETA THALASSEMIA			MC MC		REBLOZYL ZYNTEGLO	<a href="#">Use PA Form# 20420</a>	<b>Reblozyl</b> is indicated for three (3) treatments of anemia in adults: 1. in adult patients with beta thalassemia who require regular red blood cell (RBC) transfusions; 2. without previous erythropoiesis stimulating agent use (ESA-naïve) in adult patients with very low- to intermediate-risk myelodysplastic syndromes (MDS) who may require regular RBC transfusions; and 3. failing an ESA and requiring 2 or more RBC units over 8 weeks in adult patients with very low- to intermediate-risk MDS with ring sideroblasts (MDS-RS) or with myelodysplastic/myeloproliferative neoplasm with ring sideroblasts and thrombocytosis (MDS/MPN-RS-T). <b>It is not indicated for use as a substitute for RBC transfusions in patients who require immediate correction of anemia.</b>  <b>Zynteglo</b> is indicated for the treatment of adult and pediatric patients with $\beta$ -thalassemia who require regular red blood cell (RBC) transfusions.
HEMATOLOGIC DISORDER TREATMENT AGENTS			MC/DEL MC MC		CABLIVI WAYRILZ <sup>1</sup> TAVALISSE	<a href="#">Use PA Form# 20420</a> 1. Clinical PA is required to establish diagnosis and medical necessity.	<b>Tavalisse</b> is recommended for patients at risk of bleeding when one line of therapy (steroids, IVIG, splenectomy) has failed.  <b>Cablivi</b> is recommended for the treatment of adult patients with acquired thrombotic thrombocytopenic purpura (aTTP), in combination with plasma exchange and immunosuppressive therapy.  <b>Wayrilz</b> : Baseline platelet count is less than 30,000/mcL and prescribed in consultation or by a hematologist/oncologist.
COMPLEMENT RECEPTOR ANTAGONIST			MC		TAVNEOS	<a href="#">Use PA Form# 20420</a>	
WHIM SYNDROME AGENTS			MC		XOLREMDI	<a href="#">Use PA Form# 20420</a>	<b>Xolremdi</b> : In patients 12 years of age and older with WHIM syndrome (warts, hypogammaglobulinemia, infections, and myelokathexis) to increase the number of circulating mature neutrophils and lymphocytes.







Category	Coverage Indicator	Preferred Drugs		Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
<b>DERMATOLOGICAL</b>								
ISOTRETINION, ACNE		MC AMNESTEEM <sup>1</sup>			MC	ABSORICA ABSORICA LD	<a href="#">Use PA Form# 20420</a> 1. Users 24 or under, PA will not be required.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - ACNE PREPARATIONS		MC ERYDERM SOLN	MC/DEL	AKLIEF <sup>6</sup>		<a href="#">Use PA Form# 10220 for Brand Name requests</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.	
		MC/DEL ERYTHROMYCIN GEL	MC	ALTINAC CREA		<a href="#">Use PA Form# 20420 for all other requests</a>		
		MC/DEL ERYTHROMYCIN SOLN	MC	AMZEEQ <sup>6</sup>				
		MC/DEL EVOCLIN	MC	AVITA CREA				
		MC ISOTRETINOIN	MC	BENZAC				
		MC METRONIDAZOLE GEL <sup>2</sup>	MC/DEL	BENZACLIN GEL <sup>3</sup>				
		MC METRONIDAZOLE LOTN <sup>2</sup>	MC/DEL	BENZAGEL-10 GEL				
		MC/DEL TRETINOIN .025%, .01% GEL <sup>1</sup>	MC	BENZEFOAM				
		MC TRETINOIN CREA <sup>1,2</sup>	MC/DEL	BREVOXYL				
			MC	CLEOCIN-T <sup>2</sup>				
			MC	CLINAC BPO GEL				
			MC	CLINDETS SWAB				
			MC	DESQUAM-E GEL				
			MC	DESQUAM-X				
			MC	DIFFERIN 0.3% GEL				
			MC	DIFFERIN				
			MC	EMGEL GEL				
			MC	EPIDUO				
			MC	EPSOLAY				
			MC	ERYCETTE PADS				
			MC	FINEVIN CREA				
			MC	METROCREAM CREA <sup>2</sup>				
			MC	METROGEL GEL <sup>2</sup>				
			MC	METROLOTION LOTN <sup>2</sup>				
			MC	NEOBENZ MICRO				
			MC/DEL	PLIXDA				
			MC	RHOFADE				
			MC/DEL	SODIUM SULFACET/SULF LOTN				
			MC	SOOLANTRA <sup>4</sup>				
			MC/DEL	TRIAZ				
			MC	TWYNEO				
			MC	VELTIN				
			MC	WINLEVI <sup>5</sup>				
			MC	ZENCIA WASH				
			MC	ZETACET				
			MC	ZILXI				
TOPICAL- ATOPIC DERMATITIS		MC/DEL PIMECROLIMUS	MC	ANZUPGO	<a href="#">Use PA Form# 20420</a>	Preferred drugs also indicated for this condition, including topical steroids, cyclosporin AND calcineurin inhibitors must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Note: If unable to use TCIs then a trial of Eucrisa could be recommended before Dupixent.		
		MC/DEL PROTOPIC OINT	MC/DEL	CIBINQO	1. Avoid live vaccines if treated with Dupixent.			
		MC/DEL TACROLIMUS OINT	MC	NEMLUVIO	2. Clinical PA required.			
		MC ADBRY <sup>2,4</sup>			3. For the treatment of patients $\geq$ 12 years of age.			
		MC EBGLYSS <sup>2,3,4</sup>			4. Preferred only after a trial and failure of TCI.			
		MC/DEL DUPIXENT <sup>1,2,4</sup>			5. Clinical PA is required to establish diagnosis and medical necessity.			
		MC EUCRISA <sup>2,4</sup>				<b>ANZUPGO:</b> use of Anzupgo in combination with other JAK inhibitors or potent immunosuppressants is not recommended.		
		MC OPZELURA <sup>2,3,4</sup>						
		MC RINVOQ <sup>5</sup>						



Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
TOPICAL - CORTICOSTEROIDS		LOW POTENCY			LOW POTENCY	<a href="#">Use PA Form# 20420</a>	At least 1 drug from each potency of preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC	DERMA-SMOOTH-E FS BODY	MC/DEL		ACLOVATE		1. Dosing limits apply, see Dosing Consolidation List.
	MC/DEL	HYDROCORTISONE CREA	MC		DESONATE GEL		2. Treatment beyond 4 weeks is not recommended.
	MC	HYDROCORTISONE LOTN	MC/DEL		FLUOCINOLONE ACETONIDE		3. For the treatment of patients $\geq$ 12 years of age.
	MC	HYDROCORTISONE LOTN	MC/DEL		FLUOCINOLONE		4. For the treatment of patients $\geq$ 18 years of age.
	MC	TEXACORT SOLN	MC		HALOG		
	MC		MC		HYDROCORTISONE POWD		
	MC		MC		LIDA MANTLE HC CREA		
	MC		MC		PROCTOCORT CREA		
	MC		MC/DEL		VERDESO		
		MEDIUM POTENCY			MEDIUM POTENCY		
	MC/DEL	DESOXIMETASONE 0.05% CREA/GEL	MC/DEL		BESER LOTION <sup>3</sup>		
	MC	FLUTICASONE PROPIONATE CREA/OINT	MC		CLODERM CREA		
	MC	HYDROCORTISONE BUTYRATE	MC/DEL		CORDRAN		
	MC	HYDROCORTISONE OINT	MC/DEL		CUTIVATE CREA / OINT		
	MC	HYDROCORTISONE VALERATE	MC/DEL		CUTIVATE LOTN		
	MC	MOMETASONE FUROATE OINT	MC/DEL		DERMATOP		
	MC	TRIAMCINOLONE ACETONIDE .025-.1%	MC		ELOCON OINT		
	MC		MC/DEL		KENALOG AERS		
	MC		MC/DEL		LOCOID		
	MC		MC		LUXIQ FOAM		
	MC		MC/DEL		PANDEL CREA		
	MC		MC		TOPICORT		
	MC		MC/DEL		TOPICORT LP CREA		
	MC		MC		TOVET FOAM <sup>3</sup>		
		HIGH POTENCY			WESTCORT		
	MC/DEL	DESONIDE <sup>1</sup>	MC				
	MC	TRIAMCINOLONE ACETONIDE .5%	MC		AMCINONIDE CREA		
	MC		MC/DEL		BETAMETHASONE DIPROPIONATE		
	MC		MC		DESOXIMETASONE 0.25% CREA/OINT		
		VERY HIGH POTENCY			VERY HIGH POTENCY		
	MC/DEL	AUGMENTED BETA DIP	MC/DEL		CLOBETASOL PROPIONATE LOTN		
	MC/DEL	BETAMETHASONE VALERATE	MC/DEL		CLOBETASOL PROPIONATE SHAMPOO 0.05%		
	MC	DIFLORASONE DIACETATE	MC/DEL		CORMAX		
	MC/DEL		MC/DEL		DIPROLENE		
	MC/DEL		MC/DEL		IMPEKLO <sup>4</sup>		
	MC/DEL		MC/DEL		LEXETTE		
	MC/DEL		MC/DEL		OLUX FOAM		
	MC/DEL		MC/DEL		PSORCON		
	MC/DEL		MC		PSORCON E		
	MC		MC/DEL		SERNIVO SPRAY <sup>2</sup>		
	MC		MC		TEMOVATE		
		MISCELLANEOUS			ULTRAVATE		
	MC	PROCTO-KIT CREA 1%					
TOPICAL - STEROID LOCAL ANESTHETICS			MC		EPIFOAM FOAM	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - STEROID COMBINATIONS	MC	DERMA-SMOOTH-E FS SCALP	MC		CARMOL-HC CREA	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

CATEGORY	Coverage Indicator	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
TOPICAL - EMOLLIENTS	MC/DEL MC MC	AMMONIUM LACTATE CREA <sup>1</sup> AMMONIUM LACTATE LOTN 12% <sup>1</sup> VITAMIN A & D MEDICATED OINT	MC MC MC MC MC		LAC-HYDRIN CREA <sup>1</sup> LAC-HYDRIN LOTN 12% MEDERMA GEL MIMYX RENOVA CREA	<a href="#">Use PA Form# 20420</a> 1. Dosing limits still apply, see Dose Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - ENZYMES / KERATOLYTICS / UREA			MC MC MC		CARMOL 40 CREA SALEX CREA SALEX LOTN	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  Ziox, Panafil and Papain products have been removed from the PDL due to FDA safety concerns regarding drugs containing Papain.
TOPICAL - GENITAL WARTS	MC/DEL	IMIQUIMOD 5% <sup>2</sup>	MC/DEL MC/DEL MC/DEL MC MC	5 8 8 8 8	PODOFILOX SOLN CONDYLOX <sup>1</sup> ALDARA <sup>1</sup> PICATO VEREGEN <sup>1</sup>	<a href="#">Use PA Form# 20420</a> 1. Non-preferred products must be used in specified order.  2. Dosing limits still apply, see Dose Consolidation List.	
TOPICAL - LOCAL ANESTHETICS	MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	AF CAPSICUM OLEORESIN CREA CAPSAICIN CREA CAPSAICIN PATCH DIBUCAIN OINT ELA-MAX <sup>1</sup> LIDOCAINE/PRilocaine CREA <sup>1</sup> LIDOCAINE CREAM LIDOCAINE GEL LIDOCAINE PTCH 5%	MC/DEL MC/DEL MC MC MC MC/DEL		EMLA PADS EMLA CREA LIDA MANTLE CREA PONTOCAINE SOLN SYNERA ZOSTRIX ZTLIDO <sup>2</sup>	<a href="#">Use PA Form# 20420</a> 1. Lidocaine/Prilocaine cream and Ela-Max products require PA for users over 18 years of age.  2. Dosing limits still apply, see Dose Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - DEPIGMENTING AGENTS			MC MC MC MC/DEL MC/DEL MC MC MC	8 8 8 8 8 8 8 9	ALUSTRA CREA EPIQUIN MICRO GLYQUIN CREA HYDROQUINONE CREA HYDROQUINONE/SUNSCREENS SOLAQUIN FORTE CREA TRI-LUMA CREA ELDOQUIN	<a href="#">Use PA Form# 20420</a>	As per Medicaid Policy, cosmetic drugs are not covered. Non-cosmetic clinical applications will be considered by prior authorization on a case by case basis.
TOPICAL - SCABICIDES AND PEDICULICIDES	MC/DEL MC MC/DEL MC/DEL MC	ACTICIN CREA LICE KILLING SHAM LICE TREATMENT CREME RINS LIQD PERMETHRIN LOTN NATROBA <sup>1</sup>	MC MC MC/DEL MC MC MC/DEL		ELIMITE CREA EURAX LINDANE MALATHION OVIDE LOTN SPINOSAD SUSP	<a href="#">Use PA Form# 20420</a> 1. Dosing limits apply, see Dosage Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - WOUND / DECUBITUS CARE			MC MC MC		FILSUEZ REGRANEX GEL VYJUVEK	<a href="#">Use PA Form# 20420</a>	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  Regranex will be approved for diabetic patients in good control (hgbA1c <8), who are not smoking, with a stage III or IV WOON AND NPUAP lower extremity diabetic ulcer and with an adequate blood supply (TcP 02 >30, ABI>0.7 or ASP> 70), and where the underlying cause has been corrected. The wound must be free of infection and have been previously treated with preferred standard therapies for at least 2 months. Maximum approval for 20 weeks.  Accuzyme and Ethezyme products have been removed from the PDL due to FDA concerns regarding drugs containing papain.  Filsuvez: The patient has a diagnosis of dystrophic or junctional epidermolysis bullosa. The patient is at least 6 months old and does not have current evidence or history of squamous cell carcinoma or active infection in the area requiring filsuvez application. The patient has used standard wound care treatments, including silicone or foam dressings without wound resolution  Vijuvek: For the treatment of wounds in patients 6 months of age and older with dystrophic epidermolysis bullosa (DEB) with mutation(s) in the collagen type VII alpha 1 chain (COL7A1) gene.
TOPICAL - ASTRINGENTS / PROTECTANTS			MC MC MC		MOISTURIN DRY SKIN CREA PROSHIELD PLUS SKIN PROTE CREA SURGILUBE GEL	<a href="#">Use PA Form# 20420</a> 1. Dosing limits apply, see Dosage Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
HYPERHIDROSIS THERAPY - AXILLARY	MC	XERAC AC SOLN	MC	8	SOFDRA <sup>1,2</sup>	1. Clinical PA is required to establish diagnosis and medical necessity.  2. For adults and pediatric patients 9 years of age and older.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.  SOFDRA: prescribed by a dermatologist.





Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
ANTINEOPLASTIC AGENTS- LHRH ANALOGS	MC/DEL MC/DEL MC/DEL MC/DEL	LUPRON DEPOTSYRINGEKIT <sup>1</sup> LUPRON DEPOT- PED KIT <sup>1</sup> (1-month) LUPRON DEPOT-PED SYRINGEKIT (3-month) TRIPTODUR VIAL	MC/DEL MC/DEL MC/DEL MC		FIRMAGON <sup>2</sup> SUPPRELIN LA (IMPLANT) KIT TRELSTAR VANTAS <sup>2</sup>	<a href="#">Use PA Form# 20420</a> 1. Dosing limits apply, please refer to Dosage Consolidation List. 2. PA required to confirm FDA approved indication.	
ANTINEOPLASTIC AGENTS - TYROSINE KINASE INHIBITORS			MC MC/DEL MC		SPRYCEL <sup>1</sup> TYKERB <sup>2</sup> GLEEVEC <sup>1</sup>	<a href="#">Use PA Form# 20420</a> 1. Verification of diagnosis is required. 2. PA required to confirm FDA approved indication and to monitor for potential drug-drug interactions.	
ANTINEOPLASTICS- MISCELLANEOUS	MC MC/DEL MC/DEL	AMIFOSTINE MERCAPTOPURINE OXALIPLATIN	MC MC/DEL MC/DEL MC MC/DEL MC/DEL		DOCEFREZ ELOXATIN ETHYOL LEUPROLIDE PURINETHOL ZOLINZA	<a href="#">Use PA Form# 20420</a>	
ANTINEOPLASTICS- MONOCLONAL ANTIBODIES	MC/DEL	TRAZIMERA	MC/DEL MC/DEL MC MC/DEL MC MC MC/DEL		ENHERTU HERCEPTIN HERCESSI HERZUMA KANJINTI OGIVRI ONTRUZANT	<a href="#">Use PA Form# 20420</a>	
<b>CANCER</b>							
CANCER	MC	ALIMTA	MC		ABECMA	<a href="#">Use PA Form# 20420</a>	All non-preferred: A clinical PA is required to confirm appropriate clinical indication for the individual drug request. Specific to each drug all age, clinical testing requirements, previous step therapies, adjunctive drug therapy requirements, and response without disease progression will be also be evaluated for clinical appropriateness. The standard for the appropriate indication will include the FDA label as well as current NCCN guidelines.  <b>Scemblix</b> is for the treatment of adult patients with: Philadelphia chromosome-positive chronic myeloid leukemia (Ph+ CML) in chronic phase (CP), previously treated with two or more tyrosine kinase inhibitors (TKIs).
	MC/DEL	ANASTROZOLE TABS	MC		AKEEGA	1. PA required to confirm appropriate diagnosis and testing.	
	MC	ERBITUX	MC		ALECENSA	2. Avoid CYP3A drug interaction.	
	MC	IMATINIB MESYLATE	MC/DEL		ALIQOPA <sup>3</sup>	3. Clinical PA required for appropriate diagnosis.	
	MC/DEL	LETROZOLE	MC		ALUNBRIG <sup>1</sup>		
	MC	RUXIENCE	MC		ALYMSYS		
	MC/DEL	VIDAZA	MC/DEL		ARIMIDEX		
	MC	ZIRABEV	MC		AUCATZYL		
			MC		AUGTYRO		
			MC		AVMAPKI-FAKZYNJA		
			MC		AYVAKIT		
			MC/DEL		AVASTIN		
			MC/DEL		BALVERSA		
			MC		BAVENCIO <sup>1,8</sup>		
			MC		BEIZRAY		
			MC/DEL		BENDEKA <sup>3</sup>		
			MC/DEL		BESPONSA <sup>3</sup>		
			MC		BESREMI <sup>1</sup>		
			MC		BIZENGRI		
			MC		BLENREP		
			MC/DEL		BOSULIF		
			MC/DEL		BRAFTOVI <sup>1</sup>		
			MC		BREYANZI		
			MC		BRUKINSA		
			MC		CABOMETYX <sup>3</sup>		
			MC		CAMCEVI		
			MC/DEL		CALQUENCE <sup>3</sup>		
			MC		COMETRIQ <sup>3,4,5</sup>		
			MC		COTELLIC		
			MC/DEL		COPIKTRA		
			MC		DANZITEN		
			MC/DEL		DARZALEX <sup>3</sup>		
			MC/DEL		DATROWAY		
			MC/DEL		DAURISMO		
			MC/DEL		ELREXFIO		

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required			Criteria
			MC/DEL		EMPLICITI(IV) <sup>8</sup>			
			MC		EMRELIS			
			MC		EPKINLY			
			MC/DEL		ERLEADA			
			MC/DEL		ERIVEDGE			
			MC		EXKIVITY			
			MC		FARYDAK			
			MC/DEL		FEMARA			
			MC		FOLOTYN			
			MC		FOTIVDA			
			MC		FRUZAQLA			
			MC		GAVRETO			
			MC/DEL		GILOTrif <sup>4,5</sup>			
			MC		GOMEKLI			
			MC		GRAFAPEX			
			MC/DEL		HERNEXEOS			
			MC/DEL		IBRANCE			
			MC		IBTROZI			
			MC		ICLUSIG <sup>3</sup>			
			MC/DEL		IDHIFA <sup>3</sup>			
			MC		IMBRUVICA			
			MC		IMDELLTRA			
			MC/DEL		IMFINZI			
			MC/DEL		IMJUDO			
			MC		IMKELDI			
			MC		IMLYGIC			
			MC		INLURIYO			
			MC/DEL		INLYTA			
			MC/DEL		INREBIC			
			MC		INQOVI			
			MC		ITOVEBI			
			MC		IWILFIN			
			MC		JAKAFI			
			MC		JAYPIRCA <sup>1,2</sup>			
			MC		JEMPERLI			
			MC		JOBEVNE			
			MC/DEL		KEYTRUDA <sup>1</sup>			
			MC		KEYTRUDA QLEX			
			MC		KIMMTRAK			
			MC		KISQALI <sup>1</sup>			
			MC/DEL		KOSELUGO			
			MC		KRAZATI <sup>3</sup>			
			MC		KYMRIAH <sup>3,9</sup>			
			MC		KYPROLIS <sup>1</sup>			
			MC		LARTRUVO <sup>1</sup>			
			MC		LAZCLUZE			
			MC		LENVIMA			
			MC/DEL		LIBTAYO <sup>1</sup>			
			MC		LONSURF			
			MC/DEL		LORBRENA			
			MC		LOQTORZI			
			MC		LUMAKRAS			
			MC/DEL		LUMOXITI <sup>1</sup>			
			MC		LUNSUMIO <sup>1</sup>			
			MC		LYNOZYFIC			
			MC		LYNPARZA <sup>1</sup>			
			MC		LYTGOBI			
			MC		NEXAVAR <sup>1</sup>			
			MC		NERLYNX <sup>3</sup>			

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required			Criteria
			MC		NINLARO(PO)			
			MC/DEL		NUBEQA			
			MC		MARGENZA			
			MC/DEL		MEKINIST <sup>3,4</sup>			
			MC/DEL		MEKTOVI <sup>1</sup>			
			MC		MODEYSO			
			MC		MONJUVI			
			MC/DEL		MYLOTARG <sup>3</sup>			
			MC/DEL		MVASI			
			MC		ODOMZO <sup>1,2,5</sup>			
			MC		OGSIVEO			
			MC		OJEMDA			
			MC		OJJAARA			
			MC		OMISIRGE			
			MC		ONUREG			
			MC/DEL		OPDIVO <sup>3</sup>			
			MC		OPDIVO QVANTIG			
			MC		OPDUALAG			
			MC		ORGOVYX			
			MC		ORSERDU <sup>2,3</sup>			
			MC		PADCEV			
			MC		PEMAZYRE			
			MC		PEPAXTO			
			MC		PHESGO			
			MC		PHYRAGO			
			MC/DEL		PIQRAY			
			MC		POLIVY			
			MC		POMALYST			
			MC		PORTRAZZA <sup>3</sup>			
			MC		QINLOCK			
			MC		RETEVMO			
			MC		REVUFORJ			
			MC/DEL		ROMVIMZA			
			MC		REZLIDHIA			
			MC/DEL		ROZLYTREK			
			MC		RUBRACA			
			MC		RITUXAN			
			MC		RYBREVANT			
			MC		RYDAPT			
			MC		RYLAZE			
			MC		RYTELO			
			MC/DEL		SARCLISA			
			MC		SCEMBLIX <sup>1</sup>			
			MC/DEL		STIVARGA			
			MC/DEL		SUTENT <sup>1,2</sup>			
			MC/DEL		SYLATRON			
			MC		TABRECTA			
			MC		TALVEY			
			MC/DEL		TAFINLAR <sup>3,4,5,6</sup>			
			MC		TAZVERIK			
			MC/DEL		TALZENNA <sup>1</sup>			
			MC/DEL		TAGRISSO			
			MC		TECARTUS			
			MC		TECELRA			
			MC		TECENTRIQ <sup>1</sup>			
			MC		TECENTRIQ HYBREZA			
			MC		TEPMETKO			
			MC		TEVIMBRA			
			MC/DEL		TIBSOVO <sup>1</sup>			

New drugs are initially non-preferred until reviewed by the DUR Committee and the State. According to State policy, any drug requiring specific diagnosis still requires the specific diagnosis unless otherwise noted within this document.

Last update 11/4/2025

**PDL DOSAGE CONSOLIDATION LIST**

Tabs/Caps/Patches: Quantities in units

Sprays/Inhalers/Nebulizers: Quantities in GM, ML, OR MCG

Injectibles: Quantities in ML

Shaded areas are non-preferred agents - Quantities of these

non-preferred agents are available up the limit only with

prior authorization

Drug Name	Strength	Limit/Day	Limit/Days
ABILIFY SOLUTION	1MG/ML	30ML	1020/34
ACCUPRIL	5MG	1	35/35
ACCUPRIL	10MG	1	35/35
ACCUPRIL	20MG	1	35/35
ACEON	2MG	1	35/35
ACEON	4MG	1	35/35
ACTONEL	5MG	1	35/35
ACTONEL	35MG	1/WK	5/35
ACTOS	All Strengths	1	35/35
ADDERALL XR	5MG	3	90/30
ADDERALL XR	10MG	3	90/30
ADDERALL XR	15MG	3	90/30
ADDERALL XR	20MG	2	60/30
ADDERALL XR	30MG	1	35/35
ADEMPAS	All Strengths	1	35/35
ADVAIR DISKUS	All Strengths	2	60/30
ADVAIR HFA	All Strengths	4	120/30
ADZENYS XR	All Strengths	1	30/30
AEROBID	250MCG	8 INHALATIONS	21/35
AEROBID-M	250MCG	8 INHALATIONS	21/35
ALAVERT-NON DROW	TAB	1	96/96
ALENDRONATE	All Strengths	1/WK	35/35
ALTABAX	5GM		1 TUBE/30
ALTABAX	15GM		1 TUBE/30
ALTABAX	30GM		1 TUBE/30
ALTACE	1.25MG	1	35/35
ALTACE	2.5MG	1	35/35
ALTACE	5MG	1	35/35
AMARYL	1MG	1	35/35
AMARYL	2MG	1	35/35
AMBIEN	5MG		12/34
AMBIEN	10MG		12/34
AMBIEN CR	6.25MG		12/34
AMBIEN CR	12.5MG		12/34
AMERGE (Step 8)	1MG		12/30
AMERGE (Step 8)	2.5MG	2.5MG	12/30
AMLODIPINE	2.5MG	1.5	53/35 DAYS
AMLODIPINE	5MG	1.5	53/35 DAYS
AMMONIUM LACTATE CREA	12%		1 TUBE/10
AMMONIUM LACTATE LOTN	12%		1TUBE/8
AMPHETAMINE/DEXTROAMPHET ER	5MG	3	90/30
AMPHETAMINE/DEXTROAMPHET ER	10MG	3	90/30
AMPHETAMINE/DEXTROAMPHET ER	15MG	3	90/30
AMPHETAMINE/DEXTROAMPHET ER	20MG	2	60/30
AMPHETAMINE/DEXTROAMPHET ER	30MG	1	90/90
AMPHETAMINE SALT	5,10,15MG	3	105/35
AMPHETAMINE SALT	20MG	2	70/35
AMPHETAMINE SALT	30MG	1	35/35
ANDRODERM	2.5MG	2	60/30
ANDRODERM	5MG	1	30/30
ARAVA	10MG	1	35/35
ARCAPTA	75MCG	1 INHALATION	35/35
ARICEPT	5MG	1	35/35
ARICEPT	10MG	1	35/35
ARIPIPRAZOLE	2MG	2	180/90
ARIPIPRAZOLE	5MG	2	180/90
ARIPIPRAZOLE	10MG	2	180/90
ARIPIPRAZOLE	15MG	2	180/90
ARIPIPRAZOLE	20MG	1.5	135/90
ARIPIPRAZOLE	30MG	1	90/90
ARIIXTRA INJECTION	2.5MG/0.5ML		7/30
ARIIXTRA INJECTION	5MG/0.4ML		7/30
ARIIXTRA INJECTION	7.5MG/0.6ML		7/30
ARIIXTRA INJECTION	10MG/0.8ML		7/30
ARMONAIR	All Strengths	1 INHALATION	60U/30
ASMANEX 30 UNITS	220MCG	1 INHALATION	30U/30
ASMANEX 60 UNITS	220MCG	2 INHALATIONS	60U/30
ASMANEX 120 UNITS	220MCG	4 INHALATIONS	120U/30
ATACAND	4MG	1.5	53/35

Drug Name	Strength	Limit/Day	Limit/Days
ATROVENT HFA	17MCG	12 INHALATIONS	25.8/34
ATROVENT 30ML	0.03%	12 SPRAYS	30/30
ATROVENT 15ML	0.06%	16 SPRAYS	45/30
AVANDIA	2MG	1.5	53/35
AVANDIA	4MG	1	35/35
AVAPRO	75MG	1.5	53/35
AVAPRO	150MG	1	35/35
AXERT (Step 8)	6.25MG		12/30
AXERT (Step 8)	12.5MG		12/30
AZEXEL	20%		1 TUBE/18
AZILECT	All Strengths	1	35/35
BACTROBAN CREAM			1 TUBE/30
BECONASE AQ	42MCG	8 INHALATIONS	50/30
BENICAR-HCT	All Strengths	1	30/30
BENAZEPRIL	5MG	1	35/35
BENAZEPRIL	10MG	1.5	53/35
BENAZEPRIL	20MG	1	35/35
BENAZEP/HCTZ	5-6.25	1	35/35
BENAZEP/HCTZ	10/12.5	1	35/35
BEVESPI AERO		4 INHALATIONS	120/30
BONIVA	2.5MG	1	35/35
BOTOX (ADULTS)	100U/ML	1 session/90 days	600U/90
BOTOX (CHILDREN>12)	100U/ML	1 session/90 days	400U/90
BREO ELLIPTA	100/25MCG	1 INHALATIONS	60/60
BRILINTA	All Strengths	2	70/35
BRINTELLIX	All Strengths	1	35/35
BUTRANS		1 patch/WK	4/28
BYETTA	5mcg inj	0.04ML	1.2ML/30
BYETTA	10mcg inj	0.08ML	2.4ML/30
CALAN SR	120MG	1	35/35
CALAN SR	180MG	2	70/35
CALAN SR	240MG	2	70/35
CARDIZEM CD	120MG/24	1	35/35
CARDIZEM CD	180MG/24	1	35/35
CARDIZEM CD	240MG/24	1	35/35
CARDIZEM CD	300MG/24	1	35/35
CARDIZEM CD	360MG/24	1	35/35
CARDIZEM LA	120MG/24	1	35/35
CARDIZEM LA	180MG/24	1	35/35
CARDIZEM LA	240MG/24	1	35/35
CARDIZEM LA	300MG/24	1	35/35
CARDIZEM LA	360MG/24	1	35/35
CARDURA	1MG	1	35/35
CARDURA	2MG	1.5	53/35
CARDURA	4MG	1.5	53/35
CARTIA XT	120MG	1	90/90
CARTIA XT	180MG	1	90/90
CARTIA XT	240MG	1	90/90
CARTIA XT	300MG	1	90/90
CATAPRES-TTS1	0.1 MG/24HR		5/35
CATAPRES- TTS2	0.2 MG/24HR		5/35
CATAPRES- TTS3	0.3 MG/24HR		5/35
CEFIXIME	400MG	2	2/7
CELEBREX	100MG	1	35/35
CELEBREX	200MG	2	70/35
CELEBREX	400MG	1	35/35
CELEXA	20mg	0.5	17/34
CELEXA	40mg	1	51/34
CITALOPRAM	10MG	2	180/90
CITALOPRAM	20MG	2	180/90
CITALOPRAM	40MG	1	90/90
CLARINEX	REDI TAB	1	35/35
CLEOCIN-T		1 PACKAGE	1/30
CLINDAMYCIN PHOSPHATE		1 PACKAGE	1/30
COMBIVENT	103-18MCG	12 INHALATIONS	30/35
Drug Name	Strength	Limit/Day	Limit/Days
EFFEXOR XR	37.5MG	1	35/35
EFFEXOR XR	75MG	1	35/35
EMSAM	All Strengths	1	34/34

ATACAND	8MG	1.5	53/35
ATACAND	16MG	1	35/35
ATRIPLA	600MG	1	35/35
ATOMOXETINE	All Strengths	1	90/90
COMETRIQ	80MG	1	35/35
COMETRIQ	20MG	3	105/35
CONCERTA	18MG	1	30/30
CONCERTA	27MG	1	30/30
CONCERTA	36MG	2	60/30
COPAXONE INJ	20MG		1/32
COPAXONE KIT	20MG/ML		1/30
COREG CR	All Strengths	1	34/34
COSENTYX	150MG	1	1/30
CRESTOR	5MG	1	35/35
CRESTOR	10MG	1	35/35
CRESTOR	20MG	1	35/35
CRESTOR	40MG	1	35/35
CYMBALTA	All Strengths	1	35/35
DALMANE	15MG		10/30
DALMANE	30MG		10/30
DAYPRO	600MG	2	70/35
DAYTRANA	10mg/9hr (27.5mg)	1	34/34
DAYTRANA	15mg/9hr (41.3mg)	1	34/34
DAYTRANA	20mg/9hr (55.0mg)	1	34/34
DAYTRANA	30mg/9hr (82.5mg)	1	34/34
DDAVP	5ML		15/34
DENAVIR CREAM			2gm/30
DEPO-PROVERA	150MG/ML		1/90
DEPO-PROVERA	400MG/ML		2.5/90
DEPO-TESTOSTERONE	200MG/ML		20/90
DESMOPRESSIN	0.1MG	12	420/35
DESMOPRESSIN	0.2MG	6	210/35
DESONIDE	0.05%		2 TUBES/30
DESOWEN	0.05%		2 TUBES/30
DETROL LA	2MG	1	35/35
DEXEDRINE	All Strengths	3	90/30
DEXILANT	All Strengths	1	35/35
DEXTRAMPHETAMINE	All Strengths	3	90/30
DICLOFENAC 1% GEL	1% GEL		2 TUBES/30
DIFLUCAN	150MG		1/7
DILACOR XR	240MG/24	1	35/35
DILACOR XR	120MG/24	1	35/35
DILACOR XR	180MG/24	1	35/35
DILTIA - XT	120MG/24	1	90/90
DILTIA - XT	180MG	1	90/90
DILTIA - XT	240MG/24	1	90/90
DILTIAZEM CAP ER	120MG	1	90/90
DILTIAZEM CAP XR	120MG	1	90/90
DILTIAZEM CAP	120MG/24	1	90/90
DILTIAZEM CAP	180MG/24	1	90/90
DILTIAZEM CAP ER	240MG	1	90/90
DILTIAZEM CAP XR	240MG	1	90/90
DILTIAZEM XR CAP	240MG/24	1	90/90
DILTIAZEM CAP	240MG/24	1	90/90
DILTIAZEM CAP	300MG/24	1	90/90
DILTIAZEM CAP	360MG/24	1	90/90
DIOVAN	80MG	1	35/35
DIOVAN - HCT	80 - 12.5	1	35/35
DITROPAN XL	5MG	1	35/35
DITROPAN XL	10MG	2	70/35
DORAL	7.5MG		10/30
DOXAZOSIN	1MG	1	90/90
DOXAZOSIN	2MG	1.5	135/90
DOXAZOSIN	4MG	1.5	135/90
DRYSOL SOL	20%		1 BOTTLE/30DAYS
DURAGESIC PATCHES	12.5MCG/HR		11/33
DURAGESIC PATCHES	25MCG/HR		11/33
DURAGESIC PATCHES	50MCG/HR		11/33
DURAGESIC PATCHES	75MCG/HR		11/33
DURAGESIC PATCHES	100MCG/HR		22/33
DULOXETINE	20MG	3	270/90
DULOXETINE	30MG	3	270/90
DULOXETINE	60MG	2	180/90
EDEX	All Strengths		1/30

ENALAPRIL	2.5	1	90/90
ENALAPRIL	5MG	1.5	135/90
ENALAPRIL	10MG	1.5	135/90
ENALAPR/HCTZ	5-12.5	1	90/90
ENBREL	25MG/ML		8/28
ENBREL SURECLICK			8/28
ESTAZOLAM	1MG		10/30
ESTAZOLAM	2MG		10/30
ESTRING MIS	2MG		1/90
EVENITY		12 DOSES/LIFETIME	12 DOSES/LIFETIME
EVOTAZ	All Strengths	1	30/30
FELODIPINE	2.5MG	1	90/90
FELODIPINE	5MG	1.5	135/90
FENTANYL	25MCG/HR		11/33
FENTANYL	50MCG/HR		11/33
FENTANYL	75MCG/HR		11/33
FENTANYL	100MCG/HR		22/33
FETZIMA	All Strengths	1	35/35
FINASTERIDE	5MG	1	90/90
FLONASE	50MCG	4 SPRAYS	32/34
FLOVENT HFA 44MCG	44MCG	4 INHALATIONS	10.6/30
FLOVENT HFA 110MCG	110MCG	4 INHALATIONS	12/30
FLOVENT HFA 220MCG	220MCG	8 INHALATIONS	24/30
FLOVENT DISKUS	50MCG, 100MCG	4 INHALATIONS	60/30
FLOVENT DISKUS	250MCG	3 INHALATIONS	120/30
FLUCONAZOLE	150MG		1/7
FLUNISOLIDE SOLN	0.025%	16 SPRAYS	75/30
FLUOXETINE CAP	40MG	2	180/90
FLUOXETINE CAP	20MG	4	360/90
FLUOXETINE CAP	10MG	3	270/90
FLURAZEPAM	15MG		10/30
FLURAZEPAM	30MG		10/30
FLUTICASONE SPR		4 SPRAYS	48/90
FLUVOXAMINE	25MG	3	270/90
FLUVOXAMINE	50MG	3	270/90
FOCALIN	All Strengths	3	105/35
FOCALIN XR	All Strengths	1	35/35
FORFIVO XL	All Strengths	1	35/35
FOSAMAX	5MG	1	35/35
FOSAMAX	10MG	1	35/35
FOSAMAX	70MG	1/WK	5/35
FOSAMAX	40MG	2/WK	10/35
FOSINOPRIL	10MG	1.5	135/90
FOSINOPRIL	20MG	2	180/90
FRAGMIN INJ	10000U/ML	2ML	14/7
FRAGMIN INJ	2500U/.2ML	0.4ML	2.80/7
FRAGMIN INJ	25000U/ML	0.8ML	5.6/7
FRAGMIN INJ	5000U/.2ML	0.4ML	2.80/7
FRAGMIN INJ	7500U/.3ML	0.6ML	4.2/7
FROVA TAB (Step 8)	2.5MG		12/30
FULYZAQ	125MG	2	70/35
FUZEON	KIT	1	1/30
FYCOMPRA	All Strengths	1	35/35
GABAPENTIN	300MG	9	810/90
GABAPENTIN	400MG	9	810/90
GABAPENTIN	600MG	6	540/90
GABAPENTIN	800MG	4	360/90
GEODON	20MG	2	70/35
GEODON	40MG	2	70/35
GEODON	60MG	2	70/35
GEODON	80MG	2	70/35
GEODON	INJ	2	70/35
GILOTRIF	All Strengths	1	35/35
GLIMEPIRIDE	1MG	1	90/90
GLIMEPIRIDE	2MG	1	90/90
GLUCOSE TES STRP		12	420/35
GLUCAGEN INJ. HYPKIT			2/30
GLYCOLAX*	255GM		255GM/90

\* Available for once daily dosing to members under the age of 18 years

Drug Name	Strength	Limit/Day	Limit/Days
LUNESTA	2MG		12/34
LUNESTA	3MG		12/34
LUPRON DEPOT INJ	11.25MG	KIT	1/90

Drug Name	Strength	Limit/Day	Limit/Days
ILARIS			2/28
HALCION	0.125MG		10/35
HALCION	0.25		10/35
HUMIRA	40mg/0.8ml		4/28
HYDROXYZINE TAB	All Strengths	3	270/90
HYTRIN	1MG	1	35/35
HYTRIN	5MG	1	35/35
HYZAAR	50-12.5	1	35/35
IMDUR	30MG	1.5	53/35
IMDUR	60MG	1.5	53/35
IMITREX (step 8)	25MG		12/30
IMITREX (step 8)	50MG		12/30
IMITREX (step 8)	100MG		12/30
IMITREX VIAL	All Strengths		6 boxes/30
IMITREX CARTRIDGE	All Strengths		12/30
IMITREX NASAL SPRAY	All Strengths		12/30
IMITREX PEN INJCTR	All Strengths		12/30
IMIQUIMOD	5%		12/28
INTAL	800MCG	8 INHALATIONS	28.4/34
INVOKANA	All Strengths	1	35/35
IPRATROPIUM 30ML	0.03%	12 SPRAYS	90/90
IPRATROPIUM 15ML	0.06%	16 SPRAYS	135/90
ISOPTIN SR	180MG	2	70/35
IRBESARTAN	All Strengths	1	90/90
ISOPTIN SR	240MG	2	70/35
ISOSORBIDE MONO	30MG	2	180/90
ISOSORBIDE MONO	60 MG	1.5	135/90
JANUMET	All Strengths	2	70/35
JANUVIA	All Strengths	1	35/35
JUVISYNC	All Strengths	1	35/35
KETOPROFEN	100MG	2	180/90
KETOPROFEN	200MG	1	90/90
KETOROLAC	10MG	4.8	24/30
KHEDEZLA	All Strengths	1	35/35
LAC-HYDRIN CREAM	12%		1TUBE/30
LAMICTAL	25MG	6	210/35
LAMICTAL	25MG CHW	6	210/35
LAMICTAL	100MG	2	70/35
LAMISIL	250MG	1	35/35
LAMOTRIGINE	25MG	6	540/90
LAMOTRIGINE	100MG	2	180/90
LANSOPRAZOLE CAPS	All Strengths	2	180/90
LATUDA	All Strengths	1	17/34
LEFLUNOMIDE	10MG	1	90/90
LEVAQUIN	250MG	1	35/35
LEXAPRO	5MG	0.5	15/30
LIPITOR	10MG	1	35/35
LIPITOR	20MG	1	35/35
LIPITOR	40MG	1.5	53/35
LISINOP/HCTZ	10/12.5MG	1	90/90
LINEZOLID	600mg		28/60
LINZESS	All Strengths	1	35/35
LOSARTAN	All Strengths	1	90/90
LOSARTAN- HCT	All Strengths	1	90/90
LOTENSIN	5MG	1	35/35
LOTENSIN	10MG	1.5	35/35
LOTENSIN	20MG	1	53/35
LOTENSIN - HCT	5 - 6.25	1	35/35
LOTENSIN - HCT	10 - 12.5	1	35/35
LOVASTATIN	10MG	1.5	135/90
LOVASTATIN	20MG	1.5	135/90
LOVENOX INJ	30MG/.3ML	0.6	14 injections/7
LOVENOX INJ	40MG/.4ML	0.8	14 injections/7
LOVENOX INJ	60MG/.6ML	1.2	14 injections/7
LOVENOX INJ	80MG/.8ML	1.6	14 injections/7
LOVENOX INJ	100MG/ML	2	14 injections/7
LOVENOX INJ	120MG/.8ML	1.6	14 injections/7
LOVENOX INJ	150MG/ML	2	14 injections/7
LUNESTA	1MG		12/34
NIFEDIPINE ER	90MG	1	90/90
NIFEDIPINE ER,CR	30MG	1	90/90
NORVASC	2.5MG	1.5	53/35 DAYS
NORVASC	5MG	1.5	53/35 DAYS

LUPRON DEPOT INJ	22.5	KIT	1/90
LUPRON DEPOT INJ	30MG		1/90
LUPRON DEPOT INJ	30MG	KIT	1/90
LYRICA	25,50,75MG	3	102/35
LYRICA	100,150,200MG	3	102/35
LYRICA	225,300MG	2	70/35
MAVIK	1MG	1	35/35
MAVIK	2MG	1	35/35
MAXAIR AUTO	200MCG	12 INHALATIONS	14/30
MAXALT (step 8)	5MG		12/30
MAXALT (step 8)	10MG		12/30
MAXALT MLT (step 1)	5MG		12/30
MAXALT MLT (step 1)	10MG		12/30
MEDROXYPR AC	150MG/ML		1/90
MELOXICAM TABS	All Strengths	1	90/90
METADATE ER	10,20MG	3	90/30
METFORMIN ER	500MG	4	360/90
METHYLIN	All Strengths	3	90/30
METHYLPHENIDATE ER	36mg	2	180/90
METHYLPHENIDATE	All Strengths	3	90/30
METROCREAM		1 PACKAGE	1/30
METROGEL		1 PACKAGE	1/30
METROLOTION		1 PACKAGE	1/30
METRONIDAZOLE GEL		1 PACKAGE	1/30
METRONIDAZOLE LOTION		1 PACKAGE	1/30
MEVACOR	10MG	1.5	53/35
MEVACOR	20MG	1.5	53/35
MIACALCIN		3.75ml	1 bottle/34
MICARDIS	All Strengths	1	30/30
MICARDIS-HCT	All Strengths	1	30/30
MIGRANAL NASAL SPRAY	All Strengths		12/30
MIRALAX	255G	8.5G	1 bottle/30
MIRALAX	17G/PACKET	0.5 packet	15 packets/30
MIRTAZAPINE	15mg	3	270/90
MOBIC	7.5 MG	1	35/35
MOBIC	15MG	1	35/35
MOEXIPRIL	7.5	1.5	135/90
MONOPRIL	10MG	1.5	53/35
MONOPRIL	20MG	2	70/35
MUPIROCIN			1 TUBE/30
NABUMETONE	500MG	2	180/90
NABUMETONE	750MG	2	180/90
NARATRIPTAN			12/30
NASACORT AERS	55 MCG	4 SPRAYS	9.3/25
NASACORT AQ	55MCG	4 SPRAYS	17/30
NATROBA		120ML	1 bottle/30
NAYZILAM	All Strengths		5/30
NEUPOGEN INJ	300MCG/ML		10/30
NEUPOGEN INJ	480MCG/1.6		16/30
NEUPOGEN INJ	300MCG/.5ML		5/30
NEUPOGEN INJ	480MCG/.8ML		8/30
NEURONTIN	300MG	9	315/35
NEURONTIN	600MG	9	315/35
NEXIUM	20MG	1	35/35
NEXIUM	40MG	2	70/35
NEXIUM SUS	All Strengths	1	30/30
NIFEDIPINE CR	90MG	1	90/90
NIFEDIPINE ER	60MG	1	90/90
NIFEDIPINE ER	30MG	1	90/90
NIFEDIPINE ER	60MG	1	90/90
RELPAX	All Strengths		12/30
REMODULIN	All Strengths		1 MDV/30
RESTORIL	7.5MG		10/30
RESTORIL	15MG		10/30
RESTORIL	30MG		10/30
REVLIMID	All Strengths	1	35/35
REYVOW	All Strengths		4/30
RHINOCORT AQ	32MCG	8 SPRAYS	18/30
REFRESH PLUS		15 ML	1 bottle/30
REFRESH PLUS		30 ML	2 bottles/30
REFRESH TEARS		15 ML	1 bottle/30
REFRESH TEARS		30 ML	2 bottles/30
RESCULA			2 bottles/35
REYATAZ	All Strengths	1	35/35

NUVARING		1/MO	1/28
ODOMZO	200mg	1	30/30
OLMESARTAN	All Strengths	1	90/90
OLANZAPINE	2.5MG	3	270/90
OLANZAPINE	5MG	3	270/90
OLANZAPINE	7.5MG	3	270/90
OLANZAPINE	10MG	3	270/90
OLANZAPINE	15MH	2	180/90
OLANZAPINE	20MG	1.5	135/90
OLANZAPINE ODT	All Strengths	1	90/90
OMEPRAZOLE	10MG	2	180/90
OMEPRAZOLE	20MG	2	180/90
OMEPRAZOLE	40MG	2	180/90
OMNARIS	50MCG	4 sprays	12.5/30
OPSUMIT	All Strengths	1	35/35
ORUVAIL	100MG	2	70/35
ORUVAIL	200MG	1	35/35
OXAPROZIN	600MG	2	180/90
OXYCODONE ER	10,20,40MG	2	70/35
OXYCODONE ER	80MG	4	140/35
OXYCONTIN**	10,20,30,40MG	2	70/35
OXYCONTIN**	80MG	4	140/35
PANTOPRAZOLE	All Strengths	2	180/90
PAROXETINE	10MG	2	180/90
PAROXETINE	20MG	2	180/90
PAXIL	10MG	1.5	53/35
PAXIL	20MG	1	35/35
PEGASYS KIT		KIT	1/28
PLAN B			2/15 or 4/30
PLENDIL	2.5MG	1	35/35
PLENDIL	5MG	1.5	53/35
PRAVACHOL	10MG	1	35/35
PRAVACHOL	20MG	1	35/35
PRAVACHOL	40MG	1	35/35
PRAVACHOL	80MG	1	35/35
PRAVASTATIN	10MG	1	35/35
PRAVASTATIN	20MG	1	35/35
PRAVASTATIN	40MG	2	180/90
PRAVASTATIN	80MG	1	35/35
PREVPAC MIS	500MG-30MG		14/30
PRILOSEC OTC	20MG	2	168/84
PRINVIL	2.5MG	1	35/35
PRINVIL	5MG	1	35/35
PRINVIL	10MG	1.5	53/35
PRINVIL	20MG	1.5	53/35
PRINZIDE	10-12.5	1	35/35
PROAIR HFA	90mcg	12 INHALATIONS	17/34
PROTONIX	20MG	2	70/35
PROTONIX	40MG	2	70/35
PROZAC	10MG	1.5	53/35
PULMICORT	200MCG	8 INHALATIONS	1/25
PULMICORT FLEX	All Strengths	8 Inhalations	2/30
QUETIAPINE	25MG	3	270/90
QUETIAPINE	50MG	3	270/90
QUETIAPINE	100MG	3	270/90
QUETIAPINE	200MG	3	270/90
QUINAPRIL	5MG	1	90/90
QUINAPRIL	10MG	1	90/90
QUINAPRIL	20MG	1	90/90
QVAR AERS	All Strengths	8 Inhalations	14.6/25
RANITIDINE SYRUP***	15MG/ML	20ML	700ML/35
RELAFEN	500MG	2	70/35
RELAFEN	750MG	2	70/35
REMERON	15MG	1.5	53/35
Drug Name	Strength	Limit/Day	Limit/Days
SULAR	10MG	1.5	53/35
SULAR	20MG	1	35/35
SUMATRIPTAN PEN INJ	All Strengths		12/30
SUMATRIPTAN NASAL SPRAY	All Strengths		12/30
SUMATRIPTAN SYRINGE	All Strengths		12/30
SUMATRIPTAN TAB	All Strengths		12/30
SYNViSC INJ	8MG/ML		2/30
SYRINGES		10	1000/100
TAFINLAR	50MG	6	210/35

RISPERDAL	0.5MG	1.5	53/35
RISPERDAL	0.25MG	1.5	53/35
RISPERDAL	1MG	1.5	53/35
RISPERDAL	2MG	1.5	53/35
RISPERDAL	3MG	2	70/35
RISPERDAL	4MG	2	70/35
RISPERDAL INJ	25MG		2/28
RISPERDAL INJ	37.5		2/28
RISPERDAL INJ	50MG		2/28
RISPERDAL M-TAB	0.5MG	1.5	53/35
RISPERDAL M-TAB	1MG	1.5	53/35
RISPERDAL M-TAB	2MG	4	140/35
RISPERDAL SOL.	1MG/ML	8ML	280/35
RISPERIDONE	0.5MG	3	270/90
RISPERIDONE	0.25MG	3	270/90
RISPERIDONE	1MG	3	270/90
RISPERIDONE	2MG	3	270/90
RISPERIDONE	3MG	2	180/90
RISPERIDONE	4MG	2	180/90
RISPERIDONE SOL.	1MG/ML	8ML	280/35
RITALIN LA	All Strengths	1	35/35
RITALIN LA	30mg	2	70/35
SAVELLA	All Strengths	2	70/35
SEREVENT DISKUS	50MCG	2 INHALATIONS	60/30
SEROQUEL	100MG		45/30
SEROQUEL XR	150MG	1	35/35
SEROQUEL XR	200MG	1	35/35
SEROQUEL XR	300MG	2	70/35
SEROQUEL XR	400MG	2	70/35
SERTRALINE	25MG	3	270/90
SERTRALINE	50MG	3	270/90
SERTRALINE	100MG	3	270/90
SIMVASTATIN	5MG	1	35/35
SIMVASTATIN	10MG	1.5	53/35
SIMVASTATIN	20MG	1.5	53/35
SIMVASTATIN	40MG	1.5	53/35
SIMVASTATIN	80MG	1	35/35
SINGULAIR	4MG	1	35/35
SINGULAIR	5MG	1	35/35
SINGULAIR	10MG	1	35/35
SONATA	5MG		12/34
SONATA	10MG		12/34
SPIRIVA	HANDIHLR	1 INHALTION	30/30
SPORANOX SOL	10MG/ML	10ML/ML	300cc/30
SPORANOX PULSEPAK	F		30/30
SPORANOX	100MG		30/30
STADOL INJ	1MG/ML		9/35
STADOL INJ	2MG/ML		9/35
STRATTERA	All Strengths	1	35/35
SUPRAX	400MG	1	1/7
XIFYRM	All Strengths	1	90/90
XOPENEX HFA		12 INHALATIONS	2 INHALERS/34
XOPENEX NEB		12CC	408/34
ZALEPLON	All Strengths		30/30
ZECURITY	6.5		4/28
ZEMBRACE	All Strengths		3boxes/30
ZESTORETIC	10-12.5	1	35/35
ZESTRIL	2.5MG	1	35/35
ZESTRIL	5MG	1	35/35
ZESTRIL	10MG	1.5	53/35
ZESTRIL	20MG	1.5	53/35
ZETONNA	37MCG	2	60/30
ZIPRASIDONE	20MG	3	270/90
ZIPRASIDONE	40MG	3	270/90
ZOCOR	5MG	1	35/35
ZOCOR	10MG	1.5	53/35
ZOCOR	20MG	1.5	53/35
ZOCOR	40MG	1.5	53/35
ZOFRAN*	4MG	3	90/30
ZOFRAN*	8MG	1.5	45/30
ZOFRAN*	24MG	0.5	15/30
ZOFRAN*	4MG/5ML	15ML	450/30
ZOLMITRIPTAN TAB	All Strengths		12/30

TAFINLAR	75MG	4	140/35
TAMIFLU CAPS	75MG		10/30
TAZTIA XT CAP	120MG/24	1	90/90
TAZTIA XT CAP	180MG/24	1	90/90
TAZTIA XT CAP	240MG/24	1	90/90
TAZTIA XT CAP	300MG/24	1	90/90
TAZTIA XT CAP	360MG/24	1	90/90
TELMISARTAN	All Strengths	1	90/90
TEMAZEPAM	7.5MG		10/30
TEMAZEPAM	15MG		10/30
TEMAZEPAM	30MG		10/30
TEQUIN	200MG	1	35/35
TERAZOSIN	1MG	1	90/90
TERAZOSIN	5MG	1	90/90
TERBINAFINE	250MG	1	35/35
TEST STRIPS	Blood Glucose	12	420/35
TIAZAC	120MG/24	1	35/35
TIAZAC	180MG/24	1	35/35
TIAZAC	240MG/24	1	35/35
TIAZAC	300MG/24	1	35/35
TIAZAC	360MG/24	1	35/35
TIAZAC	420MG/24	1	35/35
TILADE	1.75MG	8 INHALATIONS	48.6/35
TOPAMAX SPRINKLES	All Strengths	1	35/35
TOPROL XL	25MG	1.5	53/35
TOPROL XL	50MG	1.5	53/35
TRADJENTA	All Strengths	1	35/35
TRAMADOL	50MG	8	720/90
TRAMADOL/ APAP	37.5/325MG	8	720/90
TRELEGY ELLIPTA	All Strengths	1 INHALATION	30U/30
TREXIMET	85/500	2.5	12/30
TRIAZOLAM	0.125MG		10/30
TRIAZOLAM	0.25MG		10/30
TROKENDI XR	25MG	1	35/35
TROKENDI XR	50MG	1	35/35
TROKENDI XR	100MG	1	35/35
TROKENDI XR	200MG	2	70/35
UBRELVY	All Strengths		10/30
ULTRAM	50MG	8	280/35
UNIVASC	7.5MG	1.5	53/35 DAYS
UTIBRON	7.5mcg/15.6mcg	2 INHALATIONS	60/30
VALTOCO	All Strengths		10/30
VALSARTAN-HCT	All Strengths	1	90/90
VASERETIC	5-12.5MG	1	35/35
VASOTEC	2.5MG	1	35/35
VASOTEC	5MG	1.5	53/35
VASOTEC	10MG	1.5	53/35
VENLAFAXINE TABS	25	3	270/90
VENLAFAXINE TABS	37.5	3	270/90
VENLAFAXINE TABS	100	3	270/90
VENLAFAXINE ER CAPS	37.5	3	270/90
VENLAFAXINE ER CAPS	75	3	270/90
VENLAFAXINE ER	150	2	180/90
VENTOLIN HFA	90MCG	12 INHALATIONS	36/34
VERAPAMIL ER, SR	120MG	1	90/90
VERAPAMIL ER, CR, SR	180MG	2	90/90
VERAPAMIL ER, CR, SR	240MG	2	90/90
VERELAN	180MG	1	35/35
VERELAN SR	120MG	1	35/35
VERELAN SR	180MG	1	35/35
VERELAN SR	240MG	2	70/35
VERAMYST	27.5MCG	4 sprays	10/30
VYEPTI	All Strengths		4/30
VYVANSE	All Strengths	1	35/35
VYVANSE CHEW	All Strengths	1	35/35

ZOLOFT	25MG	0.5	18/35
ZOLOFT	50MG	0.5	18/35
ZOLOFT	100MG	3	105/35
ZOLPIDEM (step 1)	5MG		30/30
ZOLPIDEM (step 1)	10MG		30/30
ZOMIG (Step 8)	5MG		12/30
ZTLIDO	All Strengths	3	90/30
ZYPREXA	2.5MG	1.5	53/35
ZYPREXA	5MG	1	35/35
ZYPREXA	7.5MG	1	35/35
ZYPREXA	10MG	1	35/35
ZYPREXA	15MG	1	35/35
ZYPREXA	20MG	1	35/35
ZYPREXA ZYDIS	5MG	1	35/35
ZYPREXA ZYDIS	10MG	1	35/35
ZYPREXA ZYDIS	15MG	1	35/35
ZYPREXA ZYDIS	20MG	1	35/35

\*Cancer diagnosis with non-daily chemotherapy required

\*\*Available without pa with CA and HO diag.

\*\*\* Ranitidine syrup available without PA to users less than 6 years old.

MDV=Multidose Vial