

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
PDL Effective October 1, 2025								
* PLEASE NOTE: For a search box hit Ctrl F								
* PLEASE NOTE: All <i>cost effective</i> generics applicable to DEL are considered PREFERRED Drugs. "BASIC" Covered Drugs are bolded with the Coverage Indicator of "MC / DEL".								
General Criteria for all PDL categories- For more information or help using the PDL, providers may call 1-888-445-0497; members should call 1-866-796-2463. To access PDL and PA materials via the internet: www.mainearepdl.org								
A: Preferred Drugs- Unless otherwise specified, preferred drugs are available without prior authorization. Step order may apply for preferred drugs in some drug categories as indicated on the PDL. (See item "D" below for explanation of step order.)								
B: Requests for Non-preferred Drugs - Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.								
C: Adequate Drug Trials - 1. The minimum trial period for each preferred and step order drug is two weeks, unless otherwise stated within specific PDL drug categories; trials with less than a two week duration will be reviewed on a case-by-case basis; 2. A trial will not be considered valid if preferred or non-preferred products were readily available (by override, individual purchase, samples, etc.); 3. Certain drug trials, such as with controlled substances, may require evidence that the preferred drugs were actually tried (example: with random pill counts and with random urine drug tests, using the methods of GC/MS with no lower threshold); 4. Adequate trials require documentation of attempts to titrate dose of preferred agents toward desired clinical response. 5. Adequate trials include prevention/treatment of common adverse effects associated with preferred agents (example: antinausea, antipruritic, etc.)								
D: Step Order - When numbers appear in the "step order" column, it means drugs in this category must be used in the order specified, with the lower numbers having preference over the higher numbers. Chart notes should be provided to confirm drug trials that do not appear in the member's MaineCare drug profile.								
E: The Department will institute strategies to ensure cost effectiveness through the use of an enhanced Drug Benefit Preferred brand drugs will no longer be preferred in any PDL drug category where preferred generic drugs are also available. It is expected that preferred generics will be used prior to any preferred brands. This will be operated as a form of step care. Preferred brands in these categories will require prior authorization for these high utilization / high cost members.								
F: Brand Name Medication Requests - (Must be submitted on the Brand Name PA request form)- According to MaineCare Benefits Manual Chapter II (80.07-5), when medically necessary covered brand-name drugs have an A-rated generic equivalent available, the most cost effective medically necessary version will be approved and reimbursed, since the brand-name and A-rated generic drugs have been determined by the FDA to be chemically and therapeutically equivalent. The Bureau does not make determinations as to whether or not a generic drug is clinically inferior or inequivalent to its brand version. This is the proper role of the FDA. Physicians should submit their reports of generic inequivalence directly to the FDA via the MEDWATCH.								
G: PA requests for non- FDA Approved Indications - Decisions will be made on a case-by-case basis until the DUR committee is able to review the evidence and make a recommendation. Interim approvals and DUR recommendations for approval of a drug for a non- FDA approved indication will require a minimum of two published, peer reviewed, non contradicted, double- blind, placebo-controlled randomized clinical studies establishing both safety and efficacy.								
H: Dose Consolidation Requirements - Some drugs may also be affected by dose consolidation requirements. Please see Dose Consolidation List and/or Splitting Tables provided in the PDL.								
I: Trials from Multiple Drug Classes - Trial/failure/intolerance to preferred agents from multiple classes within the same category or other categories of drugs may be required prior to the approval of non-preferred agents (e.g., Cymbalta, Zofran, Elidel and others).								
J: Drug-specific PA Forms - Drug-specific PA forms contain medical necessity documentation requirements and/or criteria that may not be repeated in the PDL. Drug-specific PA forms may be obtained on the web at www.mainearepdl.org .								
K: PA Exemptions for Prescribers - According to MaineCare Benefits Manual Chapter II (80.07-4), providers may receive a three (3) month exemption from prior authorization requirement for certain categories of drugs when they demonstrate high compliance with the Department's PDL. The Department will notify providers in writing which drug categories are included and what dates apply to the exemption. If a provider loses his/ her exemption, members who previously were not required to obtain a PA while the prescriber was exempt will be required to do so, and criteria for approval of that medication will need to be met.								
L: Drug-Drug Interactions (DDI) - The DUR Committee has implemented new drug-drug interaction edits requiring prior authorization. Several drug-drug combinations and PDL drug categories are affected by new PA requirements. These will be indicated in the PDL with DDI notation. Please see the DDI document provided in the PDL.								
CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
AROMATIC L-AMINO ACID DECARBOXYLASE DEFICIENCY (AADC)								
AADC DEFICIENCY AGENTS				MC	8	KEBILIDI (INJECTION) VIAL 280000000000 VG/0.5ML ELDOCAGENE EXUPARVOVEC-TNEQ	Use PA Form# 20420	
ASSORTED ANTIBIOTICS								
BETA-LACTAMS / CLAVULANATE COMBO'S	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC		AMOXICILLIN AMOXICILLIN/POTASSIUM CLA CHEW AMOXICILLIN/POTASSIUM CLA SUSR AMOXICILLIN/POTASSIUM CLA TABS AMPICILLIN BICILLIN L-A SUSP DICLOXACILLIN SODIUM CAPS OXACILLIN SODIUM SOLR PENICILLIN V POTASSIUM UNASYN SOLR	MC/DEL MC/DEL		AUGMENTIN ³ AUGMENTIN XR TB12 ⁴	Use PA Form# 20420 3. Chewable 125mg & 250mg and Solution 125mg/5ml and 250mg/5ml available without PA. 4. Use preferred generic amoxicillin/-clavulanate potassium alternatives.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Ampicillin will now be non-preferred and require prior authorization if it is currently being used in combination with either Prevacid, pantoprazole, Prilosec, or any currently non preferred PPI.
CEPHALOSPORINS	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		CEFADROXIL HEMIHYDRATE CEFAZOLIN SODIUM SOLR CEFDINIR CEFEPIME CEFPODOXIME CEFPODOXIME PROXETIL SUS CEFPODOXIME PROXETIL TAB CEFIXIME 400MG ² CAP CEFPROZIL CEPHALEXIN 250MG & 500MG CAPS CEFTAZIDIME 6MG CEFTIN SUSP CEFTRIAXONE CEFUROXIME AXETIL TABS CEPHALEXIN MONOHYDRATE	MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL		CEDAX CEFACTOR ¹ CEFADROXIL MONOHYDRATE TABS CEFIXIME SUS CEPHALEXIN TABS CEPHALEXIN 750MG CAPS CEFTIN DAXBIA FETROJA ³ FORTAZ FORTAZ SOLN KEFLEX CAPS OMNICEF ROCEPHIN SUPRAX ²	Use PA Form# 20420 1. Both brand and generic are clinically non-preferred. 2. Dosing limits apply, see Dosage Consolidation List. 3. Approvals will only be considered for patients 18 yrs of age or older who have limited or no alternative treatment options for the treatment of complicated urinary tract infections (cUTIs).	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Vantin will now be non-preferred and require prior authorization if it is currently being used in combination with either Prevacid, pantoprazole, Prilosec, or any currently non preferred PPI. As outlined in the US CDC Guidance on the Use of Expedited Partner Therapy (EPT) in the Treatment of Gonorrhea , MaineCare will cover a single 800 mg dose of cefixime for the treatment of gonorrhea as part of EPT.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
	MC MC/DEL MC		FORTAZ SOLR SUPRAX CHEWABLE TAZICEF 6GM	MC MC/DEL		TAZICEF SOLR TEFLARO		
MACROLIDES / ERYTHROMYCIN'S	MC/DEL MC/DEL MC MC MC MC MC/DEL		AZITHROMYCIN TABS AZITHROMYCIN SUSP E.E.S. ERYPED 200 SUSR ERYPED 400 SUSR ERY-TAB TBEC ERYTHROCIN STEARATE TABS ERYTHROMYCIN	MC/DEL MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		AZITHROMYCIN POW CLARITHROMYCIN SUSP CLARITHROMYCIN TABS DIFICID PCE TBEC ZITHROMAX TABS ZITHROMAX 1GM PAK ZITHROMAX TRI-PAK ZITHROMAX SUSP ZMAX ZINPLAVA	Use PA Form# 20420 1. 7-Day supply per month without PA.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Preferred erythromycin will now be non-preferred and require prior authorization if it is currently being used in combination with either carbamazepine, enablex 15mg or vesicare 10mg. Any non preferred formulation of erythromycin will require prior authorization and the member's drug profile will also be monitored for concurrent use with either carbamazepine, enablex 15mg or vesicare 10mg. DDI: Preferred Clarithromycin formulations (clarithromycin tablets) will now be non-preferred and require prior authorization if they are currently being used in combination with either carbamazepine, onglyza 5mg, enablex 15mg or vesicare 10mg. Any non preferred formulation of clarithromycin will require prior authorization and the member's drug profile will also be monitored for concurrent use with either carbamazepine, onglyza 5mg, enablex 15mg or vesicare 10mg. Zinplava will be non-preferred and require clinical prior authorization to verify it is prescribed or consulted by GI or ID specialist, diagnosis, and concurrent use of an antibacterial agent as well as limiting its use to those who have recurrent C. diff disease that has recurred despite use of guideline recommended vancomycin taper or for whom this would be contraindicated.
TETRACYCLINES	MC/DEL MC/DEL MC/DEL		DOXYCYCLINE MONOHYDRATE 100mg & 50mg CAPS MINOCYCLINE HCL CAPS TETRACYCLINE HCL CAPS	MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC		DECLOMYCIN TABS DORYX CPEP DOXYCYCLINE HYCLATE DOXYCYCLINE MONOHYDRATE 150mg & 75mg CAPS DYNACIN CAPS MINOLIRA ER NUZYRA ¹ ORACEA PERIOSTAT SEYSARA ² SOLODYN ER XIMINO	Use PA Form# 20420 1. For the treatment of patients ≥ 8 years of age. 2. For the treatment of patients ≥ 9 years of age	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
FLUOROQUINOLONES	MC/DEL MC/DEL MC/DEL		CIPROFLOXACIN LEVOFLOXACIN OFLOXACIN	MC MC MC MC MC MC MC MC MC		AVELOX SOLN AVELOX ABC PACK TABS BAXDELA CIPRO FACTIVE LEVAQUIN TABS SOLN/INJ LEVAQUIN TABS ¹ NOROXIN TABS PROQUIN XR	Use PA Form# 20420 1. Dosing limits apply, see Dosage Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Preferred Ofloxacin will now be non-preferred and require prior authorization if they are currently being used in combination with amiodarone. DDI: Preferred Levofloxacin will now be non-preferred and require prior authorization if they are currently being used in combination with amiodarone. DDI: Preferred Avelox will now be non-preferred and require prior authorization if they are currently being used in combination with amiodarone. DDI: All preferred Fluoroquinolones will require clinical PA for patients over 60 that are currently on immunosuppressants or steroid therapy. DDI: Factive is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for concurrent use with amiodarone.
AMINO GLYCOSIDES	MC MC MC/DEL MC/DEL		GENTAMICIN KITABIS PAK NEOMYCIN SULFATE TABS TOBRAMYCIN AMPUL-NEB	MC/DEL MC MC/DEL MC MC/DEL MC/DEL		ARIKAYCE ^{1,2} BETHKIS ¹ TOBI PODHALER ^{1,2} TOBI NEBU TOBRAMYCIN SULFATE SOLN ZEMDRI ²	Use PA Form# 20420 1. Clinical PA to verify appropriate diagnosis 2. See Criteria section	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. TOBI Podhaler is limited to patients with significant impairment from using nebulized version of medication Arikayce will require clinical PA to confirm MAC lung disease and for use in adults who have limited or no alternative treatment options. Zemdri will be reserved for patients with limited or no alternative treatment of care.
ANTI-MYCOBACTERIALS / ANTI-TUBERCULOSIS	MC/DEL MC/DEL MC/DEL MC/DEL		ETHAMBUTOL HCL TABS MYAMBUTOL TABS RIFABUTIN CAPS RIFAMPIN	MC/DEL MC/DEL MC MC		MYCOBUTIN CAPS PRETOMANID RIFADIN CAPS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Pretomanid is indicated as part of a combination regimen with bedaquiline and linezolid for the treatment of adults with pulmonary extensively drug resistant (XDR) or treatment-intolerant or non-responsive multidrug-resistant (MDR) tuberculosis (TB). Approval of this indication is based in limited clinical safety and efficacy data. This drug is indicated for use in a limited and specific population of patients. DDI: Preferred Rifampin will be non-preferred and require prior authorization if it is currently being used in combination with either pradaxa or latuda.
ANTIMALARIAL AGENTS	MC/DEL MC MC/DEL MC/DEL		DARAPRIM TABS KRINTAFEL ² MEFLOQUINE HCL TABS QUININE SULFATE	MC MC/DEL MC/DEL MC MC MC/DEL		ARALEN TABS CHLOROQUINE PHOSPHATE TABS³ HYDROXYCHLOROQUINE TABS³ ISONARIF ¹ MALARONE TABS PLAQUENIL TABS	Use PA Form# 20420 1. Ingredients available as preferred without PA. 2. Krintafel is preferred for ≥ 16 years of age. 3. Established users will be grandfathered.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Avoid coadministration of Krintafel with Organic Cation Transporter 2 (OCT2) and Multidrug and Toxin Extrusion (MATE) substrates (e.g. dofetilide, metformin).

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
ANTHELMINTICS	MC/DEL MC/DEL MC/DEL		ALBENDAZOLE PRAZIQUANTEL TAB STROMECTOL TABS	MC MC MC/DEL		ALBENZA TABS EMVERM BILTRICIDE TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIBIOTICS - MISC.	MC MC MC MC/DEL MC MC/DEL MC MC/DEL MC/DEL MC/DEL		AZACTAM SOLR COLY-MYCIN-M SOLR COLISTIMETHATE SODIUM SOLR FIRVANQ ⁴ FUROXONE TABS METRONIDAZOLE ¹ PENTAMIDINE ISETHIONATE SOLR SOLOSEC TRIMETHOPRIM TABS VANCOMYCIN 5GM INJ. VANCOMYCIN CAPS	MC MC MC MC/DEL MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC		AEMCOLO COLISTIMETHATE SODIUM SOLR CAYSTON ³ FLAGYL CAPS FLAGYL TABS FLAGYL ER TBCR KETEK LIKMEZ METRONIDAZOLE 375MG CAPS ¹ METRONIDAZOLE 750MG TABS ¹ NEBUPENT SOLR REBYOTA ⁵ TINDAMAX VANCOMYCIN 10GM INJ. ² XENLETA VOWST ⁵	Use PA Form# 20420 1. 375mg caps and 750mg tabs are non-preferred. Please use available preferred strengths (250mg & 500mg tabs) to obtain required dose without PA. 2. Please use multiple 5gm which are preferred to obtain dose without PA. 3. Clinical PA is required to establish CF diagnosis and medical necessity. Prior trial and failure of preferred Tobi before approval will be granted. 4. Quantity limit of one per 150ml bottle. 5. For the treatment of patients 18 years of age and older.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. 1. For macrolide resistant infections when quinolones inappropriate DDI: Ketek is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for concurrent use with either enablex 15mg or vesicare 10mg or carbamazepine. Cayston is only indicated to improve respiratory symptoms in CF patients with Pseudomonas aeruginosa. Dosing limits, as should be given TID X28 days (followed by 28 days OFF Cayston therapy). A bronchodilator should be used before administration of Cayston. Xenleta will be considered for the treatment of adults with community-acquired bacterial pneumonia (CABP) caused by the following susceptible microorganisms: Streptococcus pneumoniae, Staphylococcus aureus (methicillin-susceptible isolates), Hemophilus influenzae, Legionella pneumophila, Mycoplasma pneumoniae, and Chlamydomphila pneumoniae. Vowst: To prevent the recurrence of C.difficile infection (CDI) in individuals 18 years of age and older following antibacterial treatment for recurrent CDI (rCDI). Likmez: patient has a medical necessity for a non-solid oral dosage form. Rebyota: For the prevention of recurrence of C. difficile infection (CDI) in individuals 18 years of age and older following antibiotic treatment for recurrent CDI. The limitation of use is that Rebyota® is not indicated for treatment of CDI.
CARBAPENEMS				MC MC MC/DEL MC/DEL		INVANZ SOLR MERREM SOLR PRIMAXIN RECARBRIO	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
LINCOSAMIDES / OXAZOLIDINONES / LEPROSTATICS	MC/DEL MC/DEL MC/DEL MC MC/DEL		CLEOCIN SOLN CLEOCIN SUSR CLINDAMYCIN HCL 150CAPS DAPSONE TABS LINEZOLID 600mg TABS ²	MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL	8 8 8 8 9 9	CLEOCIN CAPS CLINDAMYCIN HCL 300CAPS ¹ SIVEXTRO VIBATIV ZYVOX SUSR ZYVOX TABS	Use PA Form# 30820 for Zyvox & Vibativ Use PA Form# 20420 for all others 1. Use multiple 150's for Clindamycin instead of 300's. 2. Quantity limit of 14 days supply within a 60-day period.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. For Zyvox or Vibativ , please see the criteria listed in the Antibacterial Antibiotics PA form.
ANTI INFECTIVE COMBO'S - MISC.	MC/DEL MC/DEL MC/DEL MC/DEL		ERYTHROMYCIN/SULF SUSR SEPTRA/DS TABS SULFAMETHOXAZOLE/TRIMETH TRIMETHOPRIM/SULFAMETHOXA	MC MC		BACTRIM DS TABS VABOMERE ¹	Use PA Form# 20420 1. For the treatment of patients ≥ 18 years of age.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIPROTOZOALS	MC/DEL MC/DEL		BENZNIDAZOLE ² LAMPIT ²	MC		ALINIA ¹	Use PA Form# 20420 1. Alina is preferred for children less than 12 years of age. 2. Clinical PA required for appropriate diagnosis.	Benznidazole is indicated for pediatric patients 2 to 12 years of age for the treatment of Chagas disease (American trypanosomiasis) caused by trypanosoma cruzi.
ANTI - FUNGALS								
ANTIFUNGALS - ASSORTED	MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		ANCOBON CAPS FLUCONAZOLE ¹ KETOCONAZOLE TABS ⁷ NYSTATIN TERBINAFINE TABS ⁴ VORICONAZOLE TABS	MC/DEL MC/DEL MC MC/DEL MC MC MC MC MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC	6 6 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	LAMISIL TABS ⁴ ITRACONAZOLE BREXAFEMME CRESEMBA ⁹ GRIFULVIN V TABS GRISEOFULVIN SUSP GRISEOFULVIN ULTRAMICROSI TABS ⁸ GRIS-PEG TABS REZZAYO ⁹ SPORANOX SOLN ² SPORANOX PULSEPAK CAPS ³ SPORANOX CAPS ³ DIFLUCAN ERAXIS INJ ⁶ GRIFULVIN SUSP ONMEL NOXAFIL ⁵ TOLSURA VFEND TABS VIVJOA	Use PA Form# 20420 See Quantity Limit table. Non-preferred products must be used in specified step order. Continue to use Anti-Fungal PA form for non-preferred products. 1. QL-1/every 7-day period (150mg only). 2. Sporanox QL 300cc/month with PA. See Quantity Limit table. 3. Sporanox QL 30/month with PA. 4. Quantity limit of one tablet daily. Please see Dosage Consolidation List. 5. Approved if immuno suppressed/ HIV or if the member has failed a 7-day trial of a preferred antifungal therapy. 6. Eraxis will be approved if submitting with documentation that it was initiated during a hospitalization and this request is to finish the hospital course.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. The other criteria are listed on the Antifungal PA form including the required proof of a non-cosmetic fungal infection. DDI: Any Griseofulvin will now be non-preferred and require prior authorization if it is currently being used in combination with either prevacid, pantoprazole, prilosec, or any currently non preferred PPI. DDI: Sporanox is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for current use with enablex 15mg, vesicare 10mg, prandin, prevacid, pantoprazole, prilosec, or any currently non preferred PPI, due to a significant drug-drug interaction. DDI: Vfend is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for concurrent use with warfarin. DDI: Fluconazole (except 150mg strength) will now be non-preferred and require prior authorization if it is currently being used with glimepiride (amaryl), enablex 15mg, or vesicare 10mg. Diflucan is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for concurrent use with either glimepiride (amaryl), enablex 15mg, or vesicare 10mg. DDI: Fluconazole will require prior authorization if being used in combination with plavix or warfarin. DDI: Ketoconazole will now be non-preferred and require prior authorization if they are currently being used in combination with any of the following medications: prevacid, pantoprazole, plavix, onglyza, enablex 15mg, vesicare 10mg, latuda, cometriq, tafinlar or omeprazole. Rezzayo: In patients 18 years of age or older who have limited or no alternative options for the treatment of candidemia and invasive candidiasis.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
HERPES AGENTS	MC/DEL MC/DEL		ACYCLOVIR VALACYCLOVIR HCL	MC/DEL MC MC/DEL MC MC/DEL	8 8 8 8 9	FAMCICLOVIR ¹ SITAVIG ZOVIRAX ¹ VALTREX TABS ¹ FAMVIR TABS ¹	Use PA Form# 20420 1. Must fail Acyclovir and Valacyclovir before non-preferred products in step order.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
INFLUENZA AGENTS	MC MC MC/DEL		AMANTADINE CAPS RELENZA DISKHALER AEPB OSELTAMIVIR ¹	MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL		AMANTADINE TABS FLUMADINE TABS FLUMIST RIMANTADINE HCL TABS TAMIFLU ¹ TAMIFLU SUS XOFLUZA	Use PA Form# 20420 1. Tamiflu and Oseltamivir 10 caps or 60cc's per month. Will be audited for presence of positive influenza tests in patient or family member.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
IMMUNE SERUMS								
IMMUNE SERUMS	MC		HYPERRHO INJ					
HEPATITIS AGENTS								
HEPATITIS C AGENTS	MC MC MC/DEL MC/DEL MC/DEL MC MC/DEL		SOFOSBUVIR/VELPATASVIR ² (Authorized generic labeler 72626 Asegua Therapeutics) MAVYRET ² PEGASYS KIT ¹ PEGASYS SOLN PEG-INTRON KIT ¹ RIBAVIRIN RIBASPHERE	MC/DEL MC/DEL MC MC MC/DEL MC MC MC MC MC/DEL		COPEGUS TABS DAKLINZA EPCLUSA ² HARVONI ² REBETOL CAPS RIBAPAK SOVALDI ² VIEKIRA PAK ² VIEKIRA XR ² VOSEVI ZEPATIER ²	Use PA Form #10700 1. Dosing limits apply, please see dosage consolidation list. 2. Approvals will require clinical PA. Please see the Hepatitis PA form for criteria.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Olysio will require a prior authorization if it is currently being used in combination with drugs known to be significant CYP3A4 inhibitors (ketoconazole, itraconazole, clarithromycin, indinavir, nefazodone, nelfinavir, ritonavir, atazanavir, saquinavir and telithromycin).
HEPATITIS AGENTS - MISC.				MC		ACTIMMUNE	Use PA Form# 20420	Approved for chronic granulomatous disease, osteopetrosis and idiopathic pulmonary fibrosis.
HEPATITIS B ONLY	MC/DEL MC		ENTECAVIR TENOFVIR	MC MC MC MC		BARACLUDE HEPSERA TABS TYZEKA VEMLIDY	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Baraclude is indicated for treatment of chronic Hep B virus (HBV) in adults with: evidence of active viral replication AND either evidence of persistent elevation in serum aminotransferases (ALT or AST) or histologically active disease, Patient is 16 years of age or older. Boxed warning: Use not recommended for those co-infected with HIV and HBV who are not also receiving highly active antiretroviral therapy (HAART). Vemlidy remain non-preferred and require prior authorization and be available to those who have evidence of bone loss or renal insufficiency or who are unable to tolerate or who have failed on preferred medications.
RSV PROPHYLAXIS								
RSV PROPHYLAXIS				MC		SYNAGIS ¹	Use PA Form# 30120 1. PA requests may be approved starting at the onset of RSV season for a maximum of 5 doses and a dosing interval not less than 30 days between injections. PA requests will be reviewed starting November of the current calendar year. Synagis dosing authorizations will extend for the recommended number of doses or until the end of epidemic RSV season as defined by CDC - whichever occurs first. Monthly prophylaxis should be discontinued for any infant or young child who experiences a breakthrough RSV hospitalization or if a child receives Nirsevimab (Beyfortus).	Please see the criteria listed on the Synagis PA form.
MS TREATMENTS								
MULTIPLE SCLEROSIS - INTERFERONS	MC MC/DEL MC		AVONEX KIT ¹ BETASERON SOLR ¹ REBIF SOLN ¹	MC MC/DEL		PLEGRIDY ¹ EXTAVIA	Use PA Form# 20430 1.Clinical PA is required to establish diagnosis and medical necessity.	Non-Preferred drugs must be tried in step-order and failed due to lack of efficacy or intolerable side effects before lower ranked non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
MULTIPLE SCLEROSIS - NON-INTERFERONS	MC MC/DEL MC/DEL MC/DEL MC MC MC		COPAXONE DALFAMPRIDINE ER DIMETHYL FUMARATE CAP FINGOLIMOD CAP ² KESIMPTA ^{2,5} TERIFLUNOMIDE TAB ² TYSABRI ^{1,2}	MC MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC MC/DEL MC MC MC MC	8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	AMPYRA AUBAGIO BAFIERTAM BRIUMVI GILENYA GLATOPA MAVENCLAD ³ MAYZENT OCREVUS ² OCREVUS ZUNOVO ² PONVORY ² TASCENSO ODT ^{2,4} TECFIDERA VUMERITY ZEPOSIA	Use PA Form# 20430 1. Provider must be enrolled in the TOUCH Prescribing program, a restricted distribution program. Clinical PA is required to establish diagnosis and medical necessity. 2. Clinical PA is required to establish diagnosis and medical necessity. 3. Due to safety profile, use of Mavenclad is generally recommended for patients who have had an inadequate response to, or are unable to tolerate, an alternate drug indicated for the treatment of MS. 4. For the treatment of patients 10 years of age and older. 5. Approved after single step through preferred drugs.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Mavenclad will require multiple trials of preferred agents including mayzent for secondary progressive disease. DDI: Due to significant increases in exposure to siponimod, concomitant use of Mayzent and drugs that cause moderate CYP2C9 and moderate or strong CYP3A4 inhibition is not recommended. Ponvory: Before initiation of Ponvory treatment, assess the following: <ul style="list-style-type: none">Complete Blood Count (CBC) - Obtain a recent (i.e. within the last 6 months) CBC, including lymphocyte count.Cardiac Evaluation - Obtain an electrocardiogram (ECG) to determine whether pre-existing conduction abnormalities are present. In patients with certain pre-existing conditions, advice from a cardiologist should be sought and first-dose monitoring is recommended. Determine whether patients are taking drugs that could slow heart rate of atrioventricular (AV) conduction.Liver Function Tests - Obtain recent (i.e. within the last 6 months) transaminase and bilirubin levels.Ophthalmic Evaluation - Obtain an evaluation of the fundus, including the macula.Current or prior medications with immune system effects - If patients are taking anti-neoplastic, immunosuppressive, or immuno-modulating therapies, or if there is a history of prior use of these drugs, consider possible unintended additive immunosuppressive effects before starting treatment with Ponvory.Vaccinations - Test for antibodies to varicella zoster virus (VZV) before starting Ponvory; VZV vaccination of antibody-negative patients is recommended prior to commencing treatment with Ponvory. If live attenuated vaccine immunizations are required, administer at least 1 month prior to initiation of Ponvory. Mayzent for Relapsing forms of MS: multiple trials of preferred agents, including an intravenous MS product. Mayzent for Active secondary progressive disease: prior trials of two preferred agents are required.
MULTIPLE SCLEROSIS - MISC				MC		ZINBRYTA ¹	Use PA Form# 20430 1. The safety and efficacy of use in children under the age of 17 years have not been established.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ASSORTED NEUROLOGICS								
NEUROLOGICS - MISC.	MC MC		BOTOX ^{2,4} DYSPORT ⁴	MC MC/DEL MC MC MC/DEL		DAXXIFY FIRDAPSE ⁵ MYOBLOC ¹ SKYSONA ^{4,6} XEOMIN ²	Use PA Form# 10210 1. Approval will be limited to Cervical Dystonia. 2. Please see botulinum PA form for additional criteria. 4. Clinical PA required. 5. For adult patients who are anti-acetylcholine receptor (AChR) antibody positive. 6. For the treatment of patients between ages 4-17 years of age.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Failed/did not tolerate therapeutic trials of muscle relaxants, unless contraindicated, including but not limited to baclofen, cyclobenzaprine, orphenadrine, skelaxin, and tizanidine. Migraine: Consideration for Botox approvals will only be made after failures of required trials of the following preferred medications: tricyclic or venlafaxine, beta blocker, valproic acid ,topiramate. Firdapse is recommended for the treatment of Lambert-Eaton myasthenic syndrome (LEMS) in adults.
NEUROLOGICS- hATTR AGENTS				MC MC/DEL MC/DEL MC/DEL MC/DEL		AMVUTTRA ¹ ATTRUBY ONPATTRO ¹ TEGSEDI ¹ VYNDAMAX ¹ VYNDAQEL ¹ WAINUA ¹	Use PA Form# 20420 1. PA required for appropriate diagnosis.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval. Tegsedi should be non-preferred and approved for patients for whom other treatments, including Onpattro, have been ineffective. Vyndamax will be considered for the treatment of the cardiomyopathy of wild type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) in adults to reduce cardiovascular mortality and cardiovascular-related hospitalization.
NEUROLOGICS- SMA	MC MC MC		GENE			GENE	Use PA Form# 20420 1. Clinical PA is required to establish diagnosis and medical necessity. 2. For patients 2 months of age and older.	Zolgensma: The patient is less than 2 years of age AND The diagnosis is spinal muscular atrophy (SMA) AND the patient has bi-allelic mutations of the SMN1 gene AND the patient does not have advanced SMA (e.g. complete paralysis of limbs or permanent ventilator dependence) AND medication is prescribed per the dosing. Spinraza: The diagnosis is spinal muscular atrophy (SMA) type 1, 2, or 3 (results of genetic testing must be submitted) AND The patient has at least 2 copies of the SMN2 gene AND The prescriber is a neurologist, pulmonologist, or other physician with expertise in treating SMA AND Baseline motor ability has been established using one of the following exams: Hammersmith Infant Neurological Exam (HINE) Hammersmith Functional Motor Scale Expanded (HF MSE) Upper Limb Module Test (non-ambulatory) Children's Hospital of Philadelphia Infant Test of Neuromuscular Disorders (CHOP INTEND) AND Prior to starting therapy, and prior to each dose, the following laboratory tests will be conducted: Treating provider attests the member has a platelet count > 50,000/ml or greater Treating provider agrees to do platelet count and coagulation test before each dose Treating provider agrees to do a quantitative spot urine protein test before each dose Concomitant use of Spinraza and Zolgensma is investigational and will not be approved AND Use of Spinraza after gene replacement therapy, including Zolgensma is investigational and will not be approved
			ZOLGENSMA ¹					
			NON-GENE			NON-GENE		
			EVRYSDI ^{1,2} SPINRAZA ¹					

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
								Note: Initial approval will be granted for 4 loading doses (the first 3 loading doses should be administered at 14-day intervals; the 4th loading dose should be administered 30 days after the 3rd dose). Renewal may be granted for up to 12 months with a maximum of 3 doses approved per year (12mg (5ml) every 4 months). For therapy continuation, clinical documentation must be submitted documenting improvement or maintenance of motor ability OR slower progression of disease than would otherwise be expected.
NEUROLOGICS- RETT SUNDROME				MC		DAYBUE ^{1,2}	Use PA Form# 20420 1.Clinical PA required for appropriate diagnosis 2. For the treatment of patients 2 years of age and older.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ALS DRUGS	MC/DEL		RILUZOLE	MC MC MC MC MC MC		EXSERVAN QALSODY RILUTEK TABS RADICAVA ¹ RELYVRIO ¹ TIGLUTIK	Use PA Form# 20420 1. Clinical PA for indication required.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Qalsody: For the treatment of amyotrophic lateral sclerosis (ALS) in adults who have a mutation in the superoxide dismutase 1 (SOD1) gene. Continued approval for this indication may be contingent upon verification of clinical benefit in confirmatory trial(s).
MOVEMENT DISORDERS	MC MC MC MC		AUSTEDO ¹ AUSTEDO XR ¹ INGREZZA ¹ TETRABENAZINE ¹	MC/DEL		XENAZINE	Use PA Form# 20420 1. Clinical PA required for appropriate diagnosis	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Avoid concomitant use of VMAT2 inhibitors with MAO inhibitors (e.g. isocarboxazid, phenelzine, or selegiline).Concomitant use with strong CYP3A4 inducers (e.g. rifampin, carbamazepine, phenytoin, St. John's wort) is not recommended
MUSCULAR DYSTROPHY AGENTS	MC		EMFLAZA ²	MC MC MC MC MC MC MC MC		AGAMREE ⁴ AMONDYS 45 ¹ DEFLAZACORT ELEVIDYS ³ DUVYZAT EXONDYS 51 ¹ VILTEPSO ³ VYONDYS 53	Use PA Form# 20420 1. Clinical PA to verify diagnosis and use of stable dose of corticosteroid for at least 6 months. 2. For the treatment of Duchenne muscular dystrophy (DMD) in patients 2 years of age and older and a documented intolerance of oral corticosteroid. 3. Clinical prior authorization to verify diagnosis and use of stable dose of corticosteroid. 4. For the treatment of Duchenne muscular dystrophy (DMD) in patients 2 years of age and older.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Amondys 45, Exondys 51 and Vyondys 53: • The prescriber is, or has consulted with, a neuromuscular disorder specialist AND • The dose does not exceed 30mg/kg once weekly AND • The patient is currently on a stable corticosteroid dose for at least 6 months (at least 3 months for Elevidy). Amondys 45, Exondys 51, Vyondys 53 Note: Initial approval will be granted for 6 months. For re-approval after 6 months, the patient must demonstrate a response to therapy. Duvyzat: The patient must meet the FDA approved age AND have a diagnosis of Duchenne Muscular Dystrophy confirmed with a confirmed mutation of the DMD gene AND • The prescriber is, or has consulted with, a neuromuscular disorder specialist • The patient is ambulatory AND • The patient is currently on a stable corticosteroid dose for at least 6 months AND • Baseline platelet counts are > 150 x 109/L and baseline triglycerides are < 300 mg/dL Elevidys and Viltepsso: The prescriber is, or has consulted with, a neuromuscular disorder specialist AND • The dose does not exceed dosing AND • The patient is currently on a stable corticosteroid dose for at least 3 months. Viltepsso: For Duchenne muscular dystrophy (DMD) in patients who have a confirmed mutation of the DMD gene that is amenable to exon 53 skipping. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.
MYASTHENIA GRAVIS	MC		PYRIDOSTIGMINE	MC MC MC MC MC	8	IMAAVY MESTINON VYVGART ¹ VYVGART HYTRULO ¹ ZILBRYSQ ¹	Use PA Form# 20420 1. For the treatment of generalized myasthenia gravis (gMG) in adult patients who are anti-acetylcholine receptor (AChR) antibody positive.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Zilbrysq recommended to vaccinate patients for meningococcal infection per current Advisory Committee on Immunization Practices (ACIP) recommendations at least 2 weeks prior to administering the first dose.
FRIEDREICH'S ATAXIA AGENTS				MC		SKYCLARYS ^{1,2}	Use PA Form# 20420 1. Clinical PA required for appropriate diagnosis. 2. For the treatment of patients 16 years of age and older.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
STERIODS								
GLUCOCORTICOIDS/ MINERALOCORTICOIDS	MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL		BUDESONIDE EC 3mg DR CAPS CELESTONE SUSP CORTEF 5 CORTISONE ACETATE TABS DELTASONE TABS DEPO-MEDROL SUSP DEXAMETHASONE DEXPAK FLUDROCORTISONE ACETATE TABS	MC MC MC/DEL MC MC MC/DEL MC MC MC	8	ALKINDI SPRINKLE CORTEF 10 and 20 TABS FLORINEF TABS HEMADY KHINDIVI¹ MEDROL TABS MEDROL DOSEPAK TABS MILLIPRED ORTIKOS	Use PA Form# 20420 1. Trial and failure, contra-indication or intolerance to Alkindi Sprinkle is required.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: All preferred steroids will require clinical PA for patients over 60 that are currently on fluoroquinolone therapy.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
	MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL		HYDROCORTISONE KENALOG METHYLPREDNISOLONE TABS PREDNISOLONE PREDNISONE SOLU-CORTEF SOLR SOLU-MEDROL SOLR	MC MC MC MC MC		ORAPRED SOLN PEDIAPRED LIQD PREDNISONE INTENSOL CONC STERAPRED TABS ZILRETTA		
HORMONE REPLACEMENT THERAPIES								
ANDROGENS / ANABOLICS	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		ANDRODERM PT24 ANDROGEL 1% ANDROGEL PUMP 1.62% DANAZOL CAPS TESTOSTERONE CYP	MC MC MC/DEL MC MC MC MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC MC/DEL MC/DEL		ANADROL-50 ANDRO LA 200 OIL ANDROGEL PACKETS 1.62% ANDROID CAPS AXIRON AZMIRO DELATESTRYL OIL DEPO-TESTOSTERONE OIL FORTESTA HALOTESTIN TABS JATENZO METHITEST TAB METHYLTESTOSTERONE CAP OXANDROLONE STRIANT MUC ER TESTIM TESTOSTERONE GEL PACKETS TESTOSTERONE SOL TESTRED CAPS TLANDO VOGELXO XYOSTED	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Additionally, laboratory evidence of a testosterone deficiency must be supplied. One of each dosage form should be tried (tablet, injection, and topical). Oxandrolone: Weight gain (adjunctive therapy): Adjunctive therapy to promote weight gain after weight loss following extensive surgery, chronic infections, or severe trauma, and in some patients who, without definite pathophysiologic reasons, fail to gain or to maintain normal weight. Other indications included in manufacturer labeling: Adjunctive therapy to offset protein catabolism with prolonged corticosteroid administration. Requirement for documentation of weight loss over two readings- Patient has involuntary weight loss of more than 10% of total body weight in less than four months) and, BMI < 18.5 (Normal BMI = 18.5 to 24.9).
ESTROGENS - PATCHES / TOPICAL	MC MC/DEL MC/DEL		EVAMIST MINIVELLE PATCH VIVELLE-DOT PTTW	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	5 8 8 8 8	ESTRADIOL PTWK DIVIGEL ¹ CLIMARA PTWK ELESTRIN ¹ MENOSTAR PATCH	Use PA Form# 20420 1. Step order drugs must be used in specified step order.	Approved for failures on multiple oral estrogen agents after 90 day trials or if unable to swallow any oral medication.
ESTROGENS - TABS	MC/DEL MC/DEL		ESTRADIOL PREMARIN TABS	MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC		ENJUVIA ESTRADIOL-NORETHINDRONE ESTRACE TABS ESTRATAB TABS MENEST TABS NORETHINDRON-ETHINYL ORTHO-EST TABS	Use PA Form# 20420 Must fail preferred products before non-preferred products.	Preferred drugs must be tried for at least 90 days and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ESTROGEN COMBO'S	MC/DEL MC/DEL MC/DEL MC/DEL		ANGELIQ COMBIPATCH PTTW PREMPHASE TABS PREMPRO TABS	MC/DEL MC/DEL MC MC/DEL MC/DEL		FEMHRT 1/5 TABS ¹ FYAVOLV LOPREEZA TAB ORTHO-PREFEST TABS ¹ SYNTEST H.S. TABS ¹	Use PA Form# 20420 1. Must fail Premphase and Prempro products before non preferred products.	Preferred drugs must be tried for at least 90 days and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
PROGESTINS	MC/DEL MC/DEL MC MC		MEDROXYPROGESTERONE ACETA ¹ NORETHINDRONE ACETATE TABS ¹ 17-ALPH HYDROXYPROGESTERONE PWDR PROGESTERONE CAPS	MC/DEL MC MC MC/DEL MC/DEL		AYGESTIN TABS CYCRIN TABS PROGESTERONE POWD PROMETRIUM CAPS PROVERA TABS	Use PA Form# 20420 1. Must fail Medroxyprogesterone and Norethindrone products before non-preferred products.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ENDOMETRIOSIS								
CENTRAL PRECOCIOUS PUBERTY AGENTS	MC		FENSOLVI ¹				Use PA Form# 20420 1. For pediatric patients 2 years of age and older with central precocious puberty (CPP).	
ENDOMETRIOSIS- NASAL	MC/DEL		SYNAREL (NASAL) SPRAY				Use PA Form# 20420	Synarel is also indicated for central precocious puberty.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
ENDOMETRIOSIS/ UTERINE FIBROIDS- ORAL	MC/DEL MC		ORILISSA ¹ MYFEMBREE ^{1,2}	MC		ORIAHNN ¹	Use PA Form# 20420 1. Prior treatment of NSAID and hormonal contraceptives required. 2. Limited to 24 months due to the risk of continued bone loss, which may not be reversible.	
ENDOMETRIOSIS- INJECTABLE	MC/DEL		DEPO-SUBQ PROVERA 104				Use PA Form# 20420	
CONTRACEPTIVES								
CONTRACEPTIVES - PROGESTIN ONLY	MC/DEL MC/DEL MC MC MC/DEL MC/DEL		CAMILA TABS ERRIN INCASSIA TAB HEATHER TAB NORETHINDRONE ACETATE 0.35MG TABS SLYND	MC/DEL MC/DEL MC		JOLIVETTE NORA-BE TABS ORTHO MICRONOR TABS	Use PA Form# 20420 Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. If member experienced adverse reactions, consider using Oral Contraceptives from other groups. DDI: Preferred Oral Contraceptives will now be non-preferred and require prior authorization if it is currently being used in combination with Tracleer.	
CONTRACEPTIVES - INJECTABLE	MC/DEL		MEDROXYPROGESTERONE ACETATE 150mg IM	MC/DEL		DEPO-PROVERA 150 mg SUSP	Use PA Form# 20420	The preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
CONTRACEPTIVE - EMERGENCY	MC/DEL MC MC MC MC/DEL MC MC/DEL MC MC/DEL		ELLA ENCONTRA ONE STEP ECONTRA EZ NEW DAY OPCION OPTION 2 MY CHOICE MY WAY LEVONORGESTREL NEXT CHOICE ¹				Use PA Form# 20420 1. Allowed 2 tablets per 30 days without PA.	Due to the extensive list of products, any covered emergency contraceptive product preferred is and available without a PA.
CONTRACEPTIVES - PATCHES/ VAGINAL PRODUCTS	MC MC MC MC/DEL		ELURYNG ¹ NUVARING RING ¹ TWIRLA XULANE ²	MC MC MC		ANNOVERA PHEXXI ZAFEMY	Use PA Form# 20420 1. Quantity limit allowing 1 every 28 days without PA. 2. Dose limits apply allowing 3 patches per 28 days supply.	Approved if adequate clinical reason given why patient unable to comply with other preferred agents including long acting injectable.
CONTRACEPTIVES- LONG ACTING REVERSIBLE	MC/DEL		MIRENA	MC/DEL MC MC MC/DEL MC/DEL		KYLEENA LILETTA NEXPLANON PARAGARD SKYLA	Use PA Form# 20420	
CONTRACEPTIVES - MONOPHASIC COMBINATION O/C'S	MC/DEL MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		APRI TABS AVIANE TABS BALZIVA CRYSSELLE-28 TABS DESOGEN TABS ESTARYLLA TAB HAILEY FE TAB ISIBLOOM TAB JUNEL FE TAB LARIN FE TAB LESSINA TAB LEVORA-28 TAB MILI TAB NORGESTIMATE-ETHINYL ESTRADIOL TAB MIBELAS 24 FE TAB MICROGESTIN FE TAB RECLIPSEN SAFYRAL TAB SPRINTEC 28 TABS	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		BEYAZ BREVICON-28 TABS LESSINA-28 TABS LEVORA LOESTRIN FE 1/20TABS LOESTRIN 1.5/30-21 TABS MICROGESTIN FE TABS LOESTRIN 1/20-21 TABS LO/OVRAL 21 TABS LO/OVRAL 28 TABS NEXTSTELLIS NORDETTE-28 TABS NORTREL OCELLA OVRAL PORTIA-28 TABS SAFYRAL ZOVIA	Use PA Form# 20420 If member experienced adverse reactions, consider using Oral Contraceptives from other groups. Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. If member experienced adverse reactions, consider using Oral Contraceptives from other groups. DDI: Preferred Oral Contraceptives will now be non-preferred and require prior authorization if it is currently being used in combination with Tracleer.	

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
	MC/DEL MC/DEL		YASMIN 28 TABS YAZ					
CONTRACEPTIVES - BI-PHASIC COMBINATIONS	MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC MC MC/DEL		AZURETTE TAB CAMRESE CAMRESE LO DESOGESTREL/ ETH/ ESTRAD 0.15/30mcg KARIVA TABS LO LOESTRIN FE PIMTREA TAB NORETHINDRONE-ETH ESTRADIOL TAB 0.5-35/1-35 SIMPESE TBDSPK 3MO VIORELE TAB	MC/DEL MC		LOSEASONIQUE	Use PA Form# 20420 If member experienced adverse reactions, consider using Oral Contraceptives from other groups.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. If member experienced adverse reactions, consider using Oral Contraceptives from other groups. DDI: Preferred Oral Contraceptives will now be non-preferred and require prior authorization if it is currently being used in combination with Tracleer.
CONTRACEPTIVES - TRI-PHASIC COMBINATIONS	MC/DEL MC/DEL MC/DEL MC MC MC MC/DEL MC/DEL MC		ENPRESSE NORGESTIMATE-ETHINYL ESTRADIOL TAB TRIPHASIL 28 TABS TRI-LO-MILI TAB TRI-LO-ESTARYLLA TAB TRI-ESTARYLLA TRI-SPRINTEC TAB TRI-LO-SPRINTEC TRINESSA	MC/DEL MC MC MC MC/DEL MC/DEL MC		NORTREL 7/7/7 ORTHO TRI-CYCLEN LO TABS	Use PA Form# 20420 If member experienced adverse reactions, consider using Oral Contraceptives from other groups.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. If member experienced adverse reactions, consider using Oral Contraceptives from other groups. DDI: Preferred Oral Contraceptives will now be non-preferred and require prior authorization if it is currently being used in combination with Tracleer.
CONTRACEPTIVES - MULTI-PHASIC COMBINATIONS				MC		NATAZIA	Use PA Form# 20420	
VASOMOTOR SYMPTOMS AGENTS								
VASOMOTOR SYMPTOMS AGENTS				MC/DEL		VEOZAH	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Veozah: Approval requires at least one preferred Hormone Replacement Therapy (HRT) and two preferred non-hormonal therapies (i.e., SSRIs, SNRIs, gabapentin, pregabalin, clonidine). DDI: Avoid concomitant use of Veozah with drugs that are weak, moderate or strong CYP1A2 inhibitors.
DIABETES SUPPLIES								
DIABETIC- SUPPLIES			CONTINUOUS GLUCOSE MONITORING ¹ DIABETIC- LANCETS DIABETIC- LANCING DEVICES DIABETIC- LANCING DEVICES DIABETIC- PEN NEEDLES DIABETIC- SYRINGES DIABETIC- TEST STRIPS DIABETIC- METERS				Use PA Form# 20420 1. Dosing limits apply. Please refer to Dose Consolidation List.	Please refer to the MaineCare Preferred Diabetic Supply List available at www.mainearepdl.org Continuous Glucose Monitoring Criteria: Patient has a diagnosis of Diabetes Mellitus AND Practitioner feels patient has sufficient training to use CGM. • 2 years of age or older for Dexcom G6 and Dexcom G7, ≥ 14 years for Medtronic Guardian, or ≥ 4 years for Freestyle Libre 2. • At least one of the following are documented: o Hypoglycemic unawareness o Treated with insulin (at least 1X day) o Has history of problematic hypoglycemia with documentation of at least one recurrent level 2 hypoglycemic events, or 1 level 3 hypoglycemic event • Approval of non-preferred products will be limited to cases where the CGM is directly integrated with the patient's insulin pump. The make and model of pump must be documented on the prior authorization.
DIABETES THERAPIES								
DIABETIC - INSULIN	MC/DEL MC MC MC MC MC MC MC MC MC MC/DEL MC/DEL		FIASP HUMALOG KWIKPEN INJ 100/ML HUMALOG JUNIOR KWIKPEN 100/ML HUMALOG MIX 75/25 HUMALOG 50/50 VIAL HUMULIN INJ 70/30 KWIKPEN HUMULIN INJ 70/30 HUMULIN R INJ U-500 INSULIN ASPART PROT MIX 70-30 INSULIN ASPART INSULIN LISPRO LANTUS SOLN LEVEMIR	MC/DEL MC/DEL MC MC MC MC MC MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC		APIDRA ADMELOG AFREZZA ¹ BASAGLAR HUMALOG KWIKPEN U-200 HUMULIN INJ 50/50 HUMULIN N INJ U-100 HUMULIN R U-100 INSULIN DEGLUDEC LYUMJEV MERILOG NOVOLIN NOVOLOG NOVOLOG MIX NOVOLOG MIX 70/30 FLEXPEN RELION	Use PA Form# 20420 1. Not to be as a monotherapy. Obtain lab values of pulmonary function and recent smoking history. 2. For the treatment of patients ≥ 3 years of age.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
DIABETIC - PENFILLS	MC MC MC MC MC/DEL MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL		HUMALOG MIX KWIK 50/50 HUMALOG MIX INJ 75/25 KWP HUMALOG KWIK INJ 100/ML HUMALOG KWIK INJ 200/ML HUMULIN R U-500 KWP INSULIN ASPART PROT MIX 70-30 PEN INSULIN ASPART PEN INSULIN LISPRO KWIKPEN U-100 LANTUS SOLOSTAR LEVEMIR FLEXTOUCH LEVEMIR FLEXPEN TOUJEO MAX SOLOSTAR TOUJEO SOLOSTAR	MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL		APIDRA OPTICLIK PEN MERILOG NOVOLIN 70/30 PEN NOVOLOG MIX PENFILL NOVOLOG PENFILL SOLN NOVOLOG FLEXPEN NOVOLOG MIX 70/30 VIAL REZVOGLAR KWIKPEN TRESIBA	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
DIABETIC - DPP- 4 ENZYME INHIBITOR	MC/DEL MC/DEL		JANUVIA ^{1,2} TRADJENTA ²	MC/DEL MC/DEL MC/DEL MC		NESINA ONGLYZA ² QTERN ZITUVIO	Use PA Form# 20420 1. Preferred if therapeutic doses of metformin are seen in members drug profile for at least 60 days within the past 18 months or if phosphate binder is currently seen in the members drug profile. 2. Dosing limits apply. Please refer to Dose Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Onglyza 5mg will require a prior authorization if it is currently being used in combination with drugs known to be significant CYP3A4 inhibitors (ketoconazole, itraconazole, clarithromycin, indinavir, nefazodone, nelfinavir, ritonavir, atazanavir, saquinavir and telithromycin).
DIABETIC - DPP- 4 ENZYME INHIBITOR-COMBO	MC/DEL MC/DEL MC/DEL		JANUMET ^{1,2} JANUMET XR ^{1,2} JENTADUETO ¹	MC/DEL MC/DEL MC MC/DEL MC MC		JENTADUETO XR KAZANO KOMBIGLYZE XR OSEN ZITUVIMET ZITUVIMET XR	Use PA Form# 20420 1. Preferred if therapeutic doses of metformin are seen in members drug profile for at least 60 days within the past 18 months or if phosphate binder is currently seen in the members drug profile. 2. Dosing limits apply. Please refer to Dose consolidation list.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Zituvimet/ Zituvimet XR: Approvals will require trial of preferred sitagliptin/metformin products or other preferred diabetic agents.
DIABETIC - LANCET-LANCET DEVICE							Use PA Form# 20420	Please refer to the MaineCare Preferred Diabetic Supply List available at www.maineicarepd.org
DIABETIC - SYRINGES-NEEDLES							Use PA Form# 20420	Please refer to the MaineCare Preferred Diabetic Supply List available at www.maineicarepd.org
DIABETIC - OTHER				MC/DEL MC		CYCLOSET SYMLIN	Use PA Form #20420	
SGLT 2 INHIBITORS	MC/DEL MC/DEL		FARXIGA JARDIANCE	MC/DEL MC/DEL		INVOKANA ¹ STEGLATRO	Use PA Form# 20420 1.Dosing limits apply please refer to Dose Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
SGLT 2 INHIBITOR COMBINATIONS	MC/DEL MC/DEL MC/DEL		SYNJARDY SYNJARDY XR XIGDOU XR	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		GLYXAMBI INVOKAMET INVOKAMET XR SEGLUROMET STEGLUJAN TRIJARDY XR	Use PA Form# 20420	Preferred drugs must be tried for at least 3 months at full therapeutic doses and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Glyxambi /Xigduo XR- Verify prior trials and failures or intolerance of preferred treatments from other diabetic categories. Synjardy XR is not recommended for patients with type 1 DM or for the treatment of diabetic ketoacidosis.
DIABETIC MONITOR	MC		RELION TRUEMETRIX AIR BLOOD GLUCOSE MONITORING SYSTEM TRUEMETRIX AIR BLOOD GLUCOSE MONITORING SYSTEM TRUEMETRIX BLOOD GLUCOSE MONITORING SYSTEM	MC MC MC MC MC MC MC MC MC		ACCUCHECK ASCENSIA ASSURE CONTOUR BREEZE Z EXACTECH FREESTYLE INSULINX FREESTYLE LITE SYSTEM KIT PRECISION XTRA METER PRODIGY	Use PA Form# 20420	Effective October 25th 2007, approvals for all non preferred meters/ test strips will require medical necessity documenting clinically significant features that are not available on any of the preferred meters.
DIABETIC TEST STRIPS	MC		RELION TRUEMETRIX TRUEMETRIX	MC MC MC MC MC MC MC		ACCUCHECK ASCENSIA ASSURE CONTOUR BREEZE Z EXACTECH FREESTYLE FREESTYLE LITE	Use PA Form# 20420	Effective October 25th 2007, approvals for all non preferred meters/ test strips will require medical necessity documenting clinically significant features that are not available on any of the preferred meters. Effective October 1, 2023, a maximum of 100 blood glucose test strips every 90 days will be available without Prior Authorization for members currently utilizing continuous glucose monitors (CGM).

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
				MC MC MC		FREESTYLE INSULINX PRECISION XTRA PRODIGY		
INCRETIN MIMETIC	MC/DEL MC MC/DEL		RYBELSUS TRULICITY VICTOZA	MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL	5 8 8 8 8 8	OZEMPIC ADLYXIN BYDUREON BCISE MOUNJARO SOLIQUA XULTOPHY	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Soliqua must try both insulin and a preferred incretin mimetic and have a medical necessity for use that is not based on convenience or simply due to the fact that one injection is needed instead of two.
DIABETIC - ORAL SULFONYLUREAS	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		CHLORPROPAMIDE TABS GLIMEPIRIDE GLIPIZIDE TABS GLIPIZIDE ER TABS GLYBURIDE MICRONIZED TABS GLYBURIDE TABS ¹ TOLAZAMIDE TABS TOLBUTAMIDE TABS	MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL		AMARYL TABS DIABETA TABS GLUCOTROL TABS GLUCOTROL XL TBCR GLYNASE TABS MICRONASE TABS	Use PA Form# 20420 1. PA required for members ≥65. Glyburide has a greater risk of severe prolonged hypoglycemia in older adults.	Preferred drugs must be tried for at least 3 months at full therapeutic doses and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: All sulfonylureas (except glyburide) will now be non-preferred and require prior authorization if it is currently being used with either ranitidine or cimetidine. DDI: Glimepiride will now be non-preferred and require prior authorization if it is currently being used with either fluconazole (except 150mg strength) or fluvoxamine. Amaryl is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for concurrent use with either fluconazole or fluvoxamine.
DIABETIC -ORAL BIGUANIDES	MC/DEL MC/DEL		METFORMIN HCL TABS METFORMIN ER	MC MC MC MC/DEL		GLUCOPHAGE TABS GLUCOPHAGE XR TB24 FORTAMET METFORMIN ER OSMOTIC	Use PA Form# 20420	Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
DIABETIC - THIAZOL / BIGUANIDE COMBO				MC/DEL MC/DEL MC MC		ACTOPLUS MET ¹ ACTOPLUS MET XR AVANDARYL ¹ AVANDAMET TABS ¹	Use PA Form# 20420 1. Requires use of Actos, Metformin, or other preferred anti-diabetics.	DDI: Actos, Avandia , or any combination product with Actos or Avandia will now be non-preferred and require prior authorization if it is currently being used with gemfibrozil.
DIABETIC - / THIAZOL	MC/DEL		PIOGLITAZONE HCL ¹	MC/DEL MC		ACTOS TABS ³ AVANDIA TABS ²	Use PA Form# 20420 1. Pioglitazone HCL is non-preferred as monotherapy. Pioglitazone HCL is preferred if therapeutic doses of metformin, sulfonylurea or insulin are seen in members drug profile for at least 60 days within the past 18 months. 2. Current users of Avandia who have tried Actos will be able to continue use of Avandia. 3. Dosing limits apply. See Dose Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Actos, Avandia , or any combination product with Actos or Avandia will now be non-preferred and require prior authorization if it is currently being used with gemfibrozil.
DIABETIC - ALPHAGLUCOSIDASE				MC		PRECOSE TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
DIABETIC - SULFONYLUREA / BIGUANIDE	MC/DEL		GLYBURIDE/METFORMIN	MC MC MC/DEL		GLUCOVANCE TABS ¹ METAGLIP TABS ¹ DUETACT ²	Use PA Form# 20420 1. Use individual ingredients. 2. Use Actos with generic glimepiride.	Approved for patients failing to achieve good diabetic control with maximal doses of individual components.
DIABETIC - MEGLITINIDES	MC		NATEGLINIDE	MC/DEL MC/DEL		PRANDIN TABS STARLIX TABS	Use PA Form# 20420	Preferred drugs from other diabetic sub-categories must be tried and failed due to lack of inadequate diabetic control or intolerable side effects before non-preferred drug will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Prandin is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for current use with both sporanox and gemfibrozil, due to a significant drug-drug interaction.
GLUCOSE ELEVATING AGENTS								
GLUCOSE ELEVATING AGENTS	MC/DEL MC/DEL		BAQSIMI ¹ GVOKE ²	MC MC		GLUCAGON DIAGNOSTIC KIT ZEGALOGUE ³	Use PA Form# 20420 1. For the treatment of patients ≥ 4 years of age. 2. For the treatment of patients ≥ 2 years of age. 3. For the treatment of patients ≥ 6 years of age.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
THYROID								
THYROID EYE DISEASE				MC		TEPEZZA	Use PA Form# 20420	

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
				MC MC/DEL MC/DEL	8 8 8	SAIZEN SOLR SOGROYA TEV-TROPIN		
ACHONDROPLASIA TREATMENT				MC		VOXZOGO ¹	Use PA Form# 20420 1. Pediatric patients with achondroplasia who are 5 years of age and older with open epiphyses.	Voxzogo: To increase linear growth in pediatric patients with achondroplasia who are 5 years of age and older with open epiphyses. This indication is approved under accelerated approval based on an improvement in annualized growth velocity. Continued approval for this indication may be contingent upon verification and description of clinical benefit in confirmatory trial(s).
SOMATOSTATIC AGENTS				MC/DEL MC MC MC/DEL MC	7 8 8 8 8	OCTREOTIDE INJ ¹ BYNFEZIA ¹ MYCAPSSA ¹ SANDOSTATIN ¹ SOMATULINE ¹	Use PA Form# 10710 1. Non-preferred products must be used in specified step order.	
GROWTH HORMONE ANTAGONISTS								
GH ANTAGONISTS				MC		SOMAVERT	Use PA Form# 10710	Approved for acromegaly patients failing surgery/radiation/drug therapy including bromocriptine and sandostatin.
VASOPRESSIN RECEPTOR ANTAGONIST								
VASOPRESSIN RECEPTOR ANTAGONIST				MC MC/DEL		JYNARQUE ¹ SAMSCA	Use PA Form# 20420 1. Clinical PA required for appropriate diagnosis	Samsca Drug Warning- Avoid use in patients with underlying liver disease, including cirrhosis, because the ability to recover from liver injury may be impaired. Limit duration of therapy to 30 days to minimize the risk of liver injury. DDI: Jynarque- Concomitant use with strong CYP3A inhibitors is contraindicated. Avoid concomitant use of Jynarque with OATP1B1/B3 and OAT3 substrates (e.g. statins, bosentan, glyburide, nateglinide, repaglinide, methotrexate, furosemide).
URINARY INCONTINENCE								
VASOPRESSINS	MC/DEL MC/DEL		DESMOPRESSIN TABS DDAVP SOLN	MC/DEL MC/DEL MC MC/DEL MC	5 6 8 8 8	DDAVP TABS DESMOPRESSIN SPRAY¹ DESMOPRESSIN ACETATE SOLN ¹ NOCDURNA ¹ NOCTIVA ¹	Use PA Form# 20420 1. Products must be used in specified step order. Nocturnal enuresis patients will be encouraged to periodically attempt stopping DDAVP.	Approved for central diabetes insipidus and for nocturnal enuresis. For nocturnal enuresis- must be over 6 years old, must fail an adequate trial of alarm training (higher success rate, lower relapse rate) and must periodically attempt weaning (at 6 month intervals).
ANTISPASMODICS	MC/DEL MC/DEL		OXYBUTYNIN TOLTERODINE	MC/DEL MC/DEL MC/DEL	8 8 8	DARIFENACIN ER TAB DITROPAN FLAVOXATE HCL TAB	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTISPASMODICS - LONG ACTING	MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		FESOTERODINE GELNIQUE GEL PACKET MYRBETRIQ OXYBUTYNIN ER TABS OXYTROL SOLIFENACIN SUCCINATE TAB TROSPIUM	MC MC/DEL MC MC/DEL MC/DEL MC MC	8 8 8 8 8 8 8	DITROPAN XL TBCR ENABLEX ^{1,2} GEMTESA ² TOLTERODINE TAB TOVIAZ VESICARE ¹ VESICARE ³ LS	Use PA Form# 20420 1. See Criteria Section. 2. Use a preferred long acting antispasmodic. 3. For the treatment of patients ≥ 2 years of age.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. 1. Vesicare 5mg and Enablex 7.5mg maximum doses if given with drugs known to be significant CYP3A4 inhibitors (ketoconazole, sporanox, erythromycin, fluconazole, nefazodone, nelfinavir, and ritonavir). DDI: Enablex 15mg and Vesicare 10mg will now be non-preferred and require prior authorization if they are currently being used in combination with any of the following medications: clarithromycin, erythromycin, ketek, crixivan, norvir, ketoconazole, fluconazole (except 150mg strength), sporanox, nefazodone, or diltiazem.
CHOLINERGIC	MC/DEL		BETHANECHOL	MC/DEL		URECHOLINE	Use PA Form# 20420	
HYPERAMMONIA TREATMENTS	MC		CARGLUMIC ACID TABS	MC		CARBAGLU TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
UREA CYCLE DISORDER	MC MC		BUPHENYL TABLET PHEBURANE GRANULES	MC MC MC MC/DEL MC/DEL		BUPHENYL POWDER RAVICTI LIQUID OLPRUVA SODIUM PHENYL BUTYRATE POWDER SODIUM PHENYL BUTYRATE TAB	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Olpruva: As adjunctive therapy to standard of care, which includes dietary management, for the chronic management of adult and pediatric patients weighing 20kg or greater and with a body surface area (BSA) of 1.2m2 or greater, with urea cycle disorders (UCDs) involving deficiencies of carbamylphosphate synthetase (CPS), ornithine transcarbamylase (OTC), or argininosuccinic acid synthetase (AS).
METABOLIC MODIFIER								
HERED. TYROSINEMIA				MC MC	6 6 8	ORFADIN NITYR HARLIKU¹	Use PA Form# 20420 1.Clinical PA is required to establish diagnosis and medical necessity.	Approved for Type 1 hereditary tyrosinemia patients. Must include laboratory evidence of dx at first PA. Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
FABRY DISEASE AGENTS				MC MC MC/DEL		ELFABRIO ¹ FABRAZYME ² GALAFOLD ¹	Use PA Form# 20420 1.Clinical PA to verify appropriate diagnosis. 2.For the treatment of patients 2 years of age and older.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Elfabrio and Galfold: For the treatment of adults with confirmed Fabry disease.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
ANTIHYPERTENSIVES / CARDIAC								
CARDIAC GLYCOSIDES	MC/DEL MC/DEL MC/DEL		DIGITEK TABS DIGOXIN LANOXIN				Use PA Form# 20420	
CARDIAC MYOSIN INHIBITORS				MC		CAMZYOS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Camzyos: For the treatment of adults with symptomatic New York Heart Association (NYHA) class II-III obstructive hypertrophic cardiomyopathy (HCM) to improve functional capacity and symptoms. DDI: Concomitant use of Camzyos with a moderate to strong CYP2C19 inhibitor or a strong CYP3A4 inhibitor is contraindicated.
CARDIAC - SINUS NODE INHIBITORS				MC		CORLANOR	Use PA Form#20420	In patients with stable, symptomatic chronic heart failure with left ventricular ejection fraction ≤35%, who are in sinus rhythm with resting heart rate ≥70 beats per minute (bpm) and either are on maximally tolerated doses of beta-blockers or have a contraindication to beta-blocker use
CARDIAC- ERAs				MC		TRYVIO	Use PA Form#20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Tryvio: In combination with other antihypertensive drugs, is indicated for the treatment of resistant hypertension, to lower blood pressure (BP) in adult patients who are not adequately controlled on other drugs. Resistant HTN is defined as a patient who takes at least 3 different class antihypertensive medications with complementary mechanisms including thiazide, ACE inhibitor, ARB, long-acting calcium channel blocker, with a trial of spironolactone, unless contra-indicated.
CARDIAC- SOLUBLE GUANYLATE CYCLASE STIMULATORS				MC/DEL		VERQUVO	Use PA Form# 20420	
CARDIAC RISK REDUCTION- SGLT2/GLP-1				MC MC MC/DEL		INPEFA ¹ LODOCO WEGOVY	Use PA Form #23976 1. To reduce the risk of cardiovascular death, hospitalization for heart failure, and urgent heart failure visit in adults with: Heart failure or Type 2 diabetes mellitus, chronic kidney disease, and other cardiovascular risk factors.	Other Preferred SGLT inhibitors must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Lodoco: Patient must have tried and failed generic colchicine due to lack of efficacy or intolerable side effects Wegovy: Patient does not have diagnosis of diabetes, end stage renal disease/dialysis, or HFrEF (EF < 45%) • Patient has BMI > 27 kg/m2, and is not being used for weight loss only • Patient has history of at least one of the following: o Stroke o Myocardial Infarction o Symptomatic peripheral arterial disease
ANTIANGINALS--Isosorbide Di-nitrate/ Mono-Nitrates	MC/DEL MC/DEL		ISOSORBIDE MONONITRATE TABS ISOSORBIDE MONONITRATE ER	MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC		DILATRATE SR CPR ISORDIL TABS ISORDIL TITRADOSE TABS ISOSORBIDE DINITRATE SUBL ISOSORBIDE DINITRATE TABS ISOSORBIDE DINITRATE CR TBCR ISOSORBIDE DINITRATE ER TBCR ISOSORBIDE DINITRATE TD TBCR IMDUR TB24 ISMO TABS MONOKET TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
NITRO - OINTMENT/CAP/CR	MC/DEL MC/DEL MC MC		NITROBID OINT NITROGLYCERIN CPR NITROL OINT NITRO-TIME CPR				Use PA Form# 20420	
NITRO - PATCHES	MC/DEL MC/DEL	1 1	NITROGLYCERIN PT24 NITRO-DUR PT 24 0.8MG	MC MC/DEL		NITRODISC PT24 NITRO-DUR PT24	Use PA Form# 20420	All preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
NITRO - SUBLINGUAL/ SPRAY	MC/DEL		NITROSTAT SUBL	MC/DEL MC MC		NITROQUICK SUBL NITROLINGUAL SOLN NITROLINGUAL TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
BETA BLOCKERS - NON SELECTIVE	MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		CARVEDILOL LEVATOL TABS NADOLOL TABS PINDOLOL TABS PROPRANOLOL HCL SOLN ¹ PROPRANOLOL HCL TABS ¹ PROPRANOLOL HCL 60MG TABS PROPRANOLOL LA CAPS	MC MC/DEL MC MC MC MC/DEL MC/DEL MC/DEL		ASPRUZYO BETAPACE TABS BETAPACE AF TABS COREG CR ³ COREG TABS CORGARD TABS INDERAL TABS HEMANGEOL SOL	Use PA Form# 20420 1. Recommend using BID since its effects do not last 24 hours. 2. Please use other strengths in combination to obtain this dose. 3. Dosing limits still apply. Please see dose consolidation list.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Concomitant use of Ranolazine products with strong CYP3A inhibitors, including ketoconazole, itraconazole, clarithromycin, nefazodone, nelfinavir, ritonavir, indinavir, and saquinavir, is contraindicated.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
	MC MC/DEL MC/DEL MC/DEL		RANOLAZINE ER TABS SOTALOL AF SOTALOL HCL TABS TIMOLOL MALEATE TABS	MC MC MC MC		INDERAL XL CAP INDERAL LA CPR INNOPRAN XL RANEXA		
BETA BLOCKERS - CARDIO SELECTIVE	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		ACEBUTOLOL HCL CAPS ATENOLOL TABS ¹ BETAXOLOL HCL TABS BISOPROLOL FUMARATE TABS BYSTOLIC METOPROLOL TARTRATE TABS ¹ METOPROLOL ER NEBIVOLOL HCL TAB	MC MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL		KERLONE TABS LOPRESSOR TABS SECTRAL CAPS TENORMIN TABS TOPROL XL TB24 ZEBETA TABS	Use PA Form# 20420 1. Recommend using Atenolol (and Metoprolol) BID since its effects do not last 24 hours.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
BETA BLOCKERS - ALPHA / BETA	MC/DEL		LABETALOL HCL TABS	MC		TRANDATE TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
BETA BLOCKERS & DURECTIC COMBOS	MC/DEL		METOPROLOL-HYDROCHLOROTHIAZIDE TAB	MC/DEL		DUTOPROL	Use PA Form# 20420	
CALCIUM CHANNEL BLOCKERS-- Amlodipine, Diltiazem, Felodipines, Nifedipines, Nisoldipine, and Verapamil	MC/DEL		AMLODIPINE ¹	MC/DEL MC MC/DEL		KATERZIA NORLIQVA NORVASC TABS ¹	Use PA Form# 20420 1. Dosing limits apply, see Dose Consolidation List.	
	MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL		DILTIA XT CP24 DILTIAZEM HCL ER CP24 DILTIAZEM HCL XR CP24 DILTIAZEM CD 300MG CP24 DILTIAZEM CD 360MG CP24 CARTIA XT CP24 ¹ DILTIAZEM CD CP24 ¹ DILTIAZEM HCL ER CP24 ¹ DILTIAZEM XR CP24 ¹ TIAZAC CP24 ¹	MC/DEL MC/DEL MC MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL	5 6 8 8 8 8 8 8 8 8	DILACOR XR CP24 ¹ TAZTIA ¹ CARDIZEM TABS ¹ CARDIZEM CD CP24 ¹ CARDIZEM LA TB24 ¹ CARDIZEM SR CP12 ¹ DILTIAZEM HCL TABS ¹ DILTIAZEM HCL ER CP12 ¹ DILTIAZEM HCL ER CP12 ¹	Use PA Form# 20420 1. Products must be used in specified order or PA will be required. Just write "Diltiazem 24-hour"and the pharmacy will use a preferred long acting Diltiazem that does not require PA.	Preferred drugs must be tried and failed (in step-order) due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: All preferred Diltiazem will now be non-preferred and require prior authorization if they are currently being used in combination with either enablex 15mg or vesicare 10mg. All non-preferred Diltiazem require prior authorization, but with any prior authorization request, the member's drug profile will also be monitored for current use with enablex 15mg or vesicare 10mg.
				MC/DEL MC/DEL		PLENDIL TB24 FELODIPINE	Use PA Form# 20420	Other Preferred calcium channel blockers must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
				MC MC		CARDENE SR CPR NICARDIPINE HCL CAPS	Use PA Form# 20420	Other Preferred calcium channel blockers must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		AFEDITAB CR NIFEDIAC CC NIFEDICAL XL TBCR NIFEDIPINE TBCR NIFEDIPINE ER TBCR	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		ADALAT CC TBCR ¹ NIFEDIPINE CAPS PROCARDIA CAPS PROCARDIA XL TBCR	Use PA Form# 20420 1. Established users of Adalat CC are grandfathered.	Preferred drug must be tried and failed in step order due to lack of efficacy or intolerable side effects before non-preferred drugs in step order will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
				MC MC		SULAR TB24 SULAR CR ¹	Use PA Form# 20420 1. Established users of 10MG and 20MG strengths are grandfathered.	
	MC/DEL MC/DEL MC/DEL	1 1 1	VERAPAMIL HCL CR TBCR VERAPAMIL HCL ER TBCR VERAPAMIL HCL SR TBCR	MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		CALAN TABS CALAN SR TBCR COVERA-HS TBCR ISOPTIN-SR VERAPAMIL HCL ER CP24 VERAPAMIL HCL SR CP24 VERAPAMIL HCL TABS VERELAN CP24 VERELAN PM CP24	Use PA Form# 20420 Products must be used in specified order or PA will be required. Just write "Verapamil 24-hour" and the pharmacy will use a preferred long acting generic that does not require PA.	Preferred drugs must be tried and failed (in step-order) due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIARRHYTHMICS	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		AMIODARONE HCL DISOPYRAMIDE FLECAINIDE MEXILETINE HCL PROCAINAMIDE PROPAFENONE	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC		CORDARONE DISOPYRAMIDE MULTAQ NORPACE PACERONE QUINIDEX	Use PA Form# 20420 1. Prescription must be written by Cardiologist.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Amiodarone will now be non-preferred and require prior authorization if it is currently being used in combination with either lovastatin (doses greater than 40mg/day) or lipitor (doses greater than 20mg/day) or levofloxacin or gemifloxacin, or moxifloxacin, or ofloxacin. DDI: Multaq will be preferred unless the following medications are seen in the member's drug profile within the last 35 days for brand name medications or 90 days

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
	MC MC/DEL MC/DEL		QUINAGLUTE QUINIDINE GLUCONATE QUINIDINE SULFATE	MC/DEL MC/DEL MC MC/DEL		TAMBOCOR TIKOSYN ¹ RYTHMOL SR RYTHMOL		
ACE INHIBITORS	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		BENAZEPRIL HCL CAPTOPRIL TABS ENALAPRIL MALEATE TABS FOSINOPRIL SODIUM LISINOPRIL TABS RAMIPRIL QUINAPRIL HCL	MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC/DEL	5 5 8 8 8 8 8 8 8 8 8 8 8	MAVIK TABS ACCUPRIL TABS ACEON TABS ¹ ALTACE CAPS ¹ EPANED LOTENSIN TABS ¹ MOEXIPRIL HCL ¹ MONOPRIL HCT TABS ¹ PRINIVIL TABS ¹ QBRELIS UNIVASC ¹ VASOTEC TABS ¹ ZESTRIL TABS ¹	Use PA Form# 20420 1. Non-preferred products must be used in specified order.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Non-preferred products are subject to step-order requirements unless clinical circumstances warrant exception.
ANGIOTENSIN RECEPTOR BLOCKER	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		AMLODIPINE-OLMESARTAN TAB ³ IRBESARTAN ¹ LOSARTAN ¹ MICARDIS TABS ³ OLMESARTAN ¹ TELMISARTAN ¹	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC	8 8 8 8 8 8	ATACAND TABS AVAPRO BENICAR TABS COZAAR DIOVAN EDARBI TEVETEN TABS	Use PA Form# 20420 1. Dosing limits apply, please see Dose Consolidation List. 2. Use preferred active ingredients which are available without PA. 3. Preferred without a PA only if patient on a diabetic therapy or prior ACE therapy.	Per best practices patient should have trialed prior therapy of ACE inhibitor or currently on a diabetic therapy.
DIRECT RENIN INHIBITOR				MC/DEL MC/DEL MC/DEL		AMTURNIDE TEKTURN ¹ TEKAMLO	Use PA Form# 20420 1. Must show failure of single and combination therapy from all preferred antihypertensive categories.	
ANTIHYPERTENSIVES - CENTRAL	MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL		CLONIDINE HCL TABS GUANFACINE HCL TABS HYDRALAZINE HCL TABS HYLOREL TABS METHYLDOPA TABS MINOXIDIL TABS PRAZOSIN HCL CAPS RESERPINE TABS	MC/DEL MC/DEL MC MC MC/DEL MC MC/DEL MC/DEL		CLONIDINE PATCH CLONIDINE TTS GUANABENZ ACETATE TABS ISMELIN TABS MINIPRESS CAPS NEXICLON TENEX TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ACE INHIBITORS AND CA CHANNEL BLOCKERS				MC/DEL MC MC MC/DEL	8 8 8 9	AMLODIPINE/BENAZEPRIL PRESTALIA ¹ TARKA TBCR LOTREL CAPS	Use PA Form# 20420 1. Prestalia will only be approved for patients ≥ 18 years of age. Use individual preferred generic medications.	
ACE AND THIAZIDE COMBO'S	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		BENAZEPRIL HCL/HYDROCHLOR CAPTOPRIL/HYDROCHLOROTHIA ENALAPRIL MALEATE/HCTZ TABS LISINOPRIL-HCTZ TABS LOTENSIN HCT TABS	MC/DEL MC MC/DEL MC/DEL MC MC/DEL		ACCURETIC TABS MONOPRIL HCT TABS PRINZIDE TABS UNIRETIC TABS VASERETIC TABS ZESTORETIC TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
BETA BLOCKERS AND DIURETIC COMBO'S	MC/DEL MC/DEL MC/DEL		ATENOLOL/CHLORTHALIDONE BISOPROLOL FUMARATE/HCTZ PROPRANOLOL/HCTZ	MC/DEL MC/DEL MC MC MC/DEL		CORZIDE TABS LOPRESSOR HCT TABS TENORETIC TIMOLIDE 10/25 TABS ZIAC TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ARB'S AND CA CHANNEL BLOCKERS	MC/DEL MC/DEL MC/DEL		AMLODIPINE/VALSARTAN AMLODIPINE/VALSARTAN HCT TRIBENZOR	MC/DEL MC MC/DEL MC/DEL		AZOR BYVALSON EXFORGE EXFORGE HCT	Use PA Form# 20420	DDI: Byvalson will be non-preferred and require a prior authorization if it is currently being used in combination with drugs known to be significant CYP2D6 inhibitors (e.g. quinidine, propafenone, fluoxetine, paroxetine). Per best practices, patient should have trialed prior therapy of ACE inhibitor or currently on a diabetic therapy.
ARB'S AND DIURETICS	MC/DEL MC/DEL MC/DEL MC/DEL		BENICAR HCT ¹ LOSARTAN HCT ¹ MICARDIS HCTTABS ¹ VALSARTAN-HCT ¹	MC/DEL MC/DEL MC MC/DEL	7 8 8 8	IRBESARTAN HYDROCHLOROTHIAZIDE ATACAND HCT TABS AVALIDE TABS ¹ DIOVAN HCT TABS ¹	Use PA Form# 20420 1. Dosing limits apply, see Dose Consolidation List.	Per best practices, patient should have trialed prior therapy of ACE inhibitor or currently on a diabetic therapy.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
				MC/DEL MC	8 8	HYZAAR TABS TEVETEN HCT TABS		
ANGIOTENSIN MODULATORS- ARB COMBINATION	MC		ENTRESTO	MC/DEL MC		EDARBYCLOR ENTRESTO SPRINKLES	Use PA Form# 20420	
ARB'S AND DIRECT RENIN INHIBITOR COMBINATION				MC/DEL		VALTURNA	Use PA Form# 20420	
DIURETICS	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC		ACETAZOLAMIDE TABS AMILORIDE HCL BUMETANIDE CHLOROTHIAZIDE TABS CHLORTHALIDONE TABS EDECIN TABS EDECIN TABS HYDROCHLOROTHIAZIDE INDAPAMIDE TABS METHAZOLAMIDE TABS METHYLCLOTHIAZIDE TABS SPIRONOLACTONE SPIRONOLACTONE/HYDRO TORSEMIDE TABS TRIAMTERENE/HCTZ ZAROXOLYN TABS	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC	8	ALDACTAZIDE TABS ALDACTONE TABS BUMEX TABS DEMADEX TABS DIAMOX DIURIL DYAZIDE CAPS CAROSPIR ENDURON TABS FUROSCIX HEMICLOR INSPRA INZIRQO KERENDIA KEVEYIS LASIX TABS MAXZIDE MICROZIDE CAPS MIDAMOR TABS NAQUA TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Furoscix: The indication for use is the treatment of congestion due to fluid overload in adults with NYHA Class II or Class III chronic heart failure AND the medication is being prescribed by or in consultation with a cardiologist AND the patient is experiencing symptoms despite compliance with oral loop diuretic therapy AND oral loop diuretic therapy will be resumed as soon as practical AND medical reasoning beyond convenience is provided for not pursuing therapy in an outpatient infusion setting. PA approval will be authorized for 1 month. Kerendia: Patient must be on max tolerated preferred ACE-I/ARB and SGLT-2. DDI: The concomitant use of Keveyis with high dose aspirin is contraindicated.
CCB / LIPID				MC/DEL		CADUET	Use PA Form# 20420	
NEUROGENIC ORTHOSTATIC HYPOTENSION								
NEUROGENIC ORTHOSTATIC HYPOTENSION				MC		NORTHERA	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
LIPID DRUGS								
CHOLESTEROL - BILE SEQUESTRANTS	MC/DEL MC/DEL		CHOLESTYRAMINE COLESTIPOL HCl	MC/DEL MC/DEL MC MC/DEL		COLESTID PREVALITE QUESTRAN WELCHOL TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
CHOLESTEROL - FIBRIC ACID DERIVATIVES	MC/DEL MC/DEL MC/DEL		FENOFIBRATE TAB GEMFIBROZIL TABS NIACIN ER	MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC		ANTARA LOPID FENOFIBRATE 120mg TAB FENOFIBRATE CAP FIBRICOR LIPOFEN LOFIBRA NIASPAN ER TRICOR TRIGLIDE	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Fenofibrate is preferred but will require a prior authorization requests if used concurrent with warfarin. DDI: Gemfibrozil will now be non-preferred and require prior authorization if it is currently being used with any of the following medications: prandin, actos, avandia, any avandia/actos combination product, any HMG-COA Reductase Inhibitors (statins), or warfarin.
CHOLESTEROL - HMG COA + ABSORB INHIBITORS MORE POTENT DRUGS/- COMBINATIONS	MC/DEL MC/DEL MC MC/DEL		ATORVASTATIN EZETIM/SIMVA TAB ROSUVASTATIN SIMVASTATIN ¹	MC MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC/DEL MC		ATORVALIQ CRESTOR EZALLOR SPRINKLES ³ FLOLIPID LIPITOR LIPTRUZET ZOCOR SIMVASTATIN 80MG ^{1,2} VYTORIN	Use PA Form# 20420 1. Dosing limits apply, see Dosage Consolidation List. 2. Current users grandfathered. 3. For the treatment of patients ≥ 18 years of age.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Lipitor (doses greater than 20mg/day) will now be non-preferred and require prior authorization if they are currently being used in combination cyclosporine. DDI: Lipitor (doses greater than 20mg/day) will now be non-preferred and require prior authorization if it is currently being used in combination with amiodarone. DDI: All preferred statins will now be non-preferred and require prior authorization if it is currently being used in combination with gemfibrozil.
CHOLESTEROL - HMG COA + ABSORB INHIBITORS LESS POTENT DRUGS/- COMBINATIONS	MC/DEL MC/DEL MC/DEL		EZETIMIBE TABS LOVASTATIN TABS ² PRAVASTATIN ²	MC MC/DEL MC/DEL MC	8 8 8 8	ALTOPREV TB24 FLUVASTATIN TAB ER LESCOL XL TB24 LIVALO	Use PA Form# 20420 2. Dosing limits apply, please see Dosage Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Zetia will be approved for patients unable to tolerate all other therapies or unable to achieve cholesterol goal with maximally tolerated dose of most potent statins.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
				MC/DEL MC MC MC/DEL MC/DEL MC	8 8 8 8 8 8	MEVACOR TABS NEXLETOL NEXLIZET PRAVACHOL TABS PRAVIGARD ZETIA TABS		DDI: Lescol will now be non-preferred and require prior authorization if it is currently being used in combination with diclofenac. DDI: Lovastatin (doses greater than 40mg/day) will now be non-preferred and require prior authorization if it is currently being used in combination with amiodarone. DDI: Lovastatin (doses greater than 20mg per day) will now be non-preferred and require prior authorization if it is currently being used in combination cyclosporine. DDI: All preferred statins will now be non-preferred and require prior authorization if it is currently being used in combination with gemfibrozil.
CHOLESTEROL - HMG COA + ABSORB INHIBITORS STATIN/ NIACIN COMBO	MC		SIMCOR	MC		ADVICOR TBCR	Use PA Form# 20420	
FAMILIAL HYPERCHOLESTEROLEMIA	MC MC		PRALUENT (LABLER 72733) PEN ^{1,2,3,5} REPATHA ^{1,2,3}	MC MC MC MC		EVKEEZA ^{1,4} JUXTAPID KYNAMRO ¹ LEQVIO	Use PA Form# 20420 1. Clinical PA required for appropriate diagnosis. 2. Quantity limits apply. 3. Documented adherence to lipid lowering medications and abstinence from tobacco for previous 90 days. 4. For treatment of patients ≥ 12 years of age. 5. Approval of Praluent NDC's with labeler code 00024 will be considered only if labeler code 72733 NDC's are on a long-term backorder and unavailable from the manufacturer.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Juxtapid is contraindicated with strong CYP3A4 inhibitors. Juxtapid dosage should not exceed 30mg daily when it is used concomitantly with weak CYP3A4 inhibitors. Kynamro requires an appropriate lab testing prior to starting (ALT<AST), alkaline phosphatase and total bilirubin, monthly liver-related tests for the first year, then every three months. Repatha and Praluent Criteria for approval: The patient's age is FDA approved for the given indication AND • Concurrent use with statin therapy AND • Documented adherence to prescribed lipid lowering medications for the previous 90 days AND • Recommended or prescribed by a lipidologist or cardiologist AND • Inability to reach goal LDL-C despite a trial of 2 or more maximum tolerated dose of statins (one of which must be atorvastatin or rosuvastatin) and ezetimibe 10mg daily. Additional criteria for the diagnosis of heterozygous familial hypercholesterolemia (HeFH): (both are required): Total cholesterol > 290 mg/dL OR LDL-C > 190 mg/dL AND one of the following • Presence of tendon xanthomas OR • In 1st or 2nd degree relative-documented tendon xanthomas, MI at age ≤ 60 years or TC > 290 mg/dL. Additional criteria for the diagnosis of clinical atherosclerotic cardiovascular disease: History of MI, angina, coronary or other arterial revascularization, stroke, TIA, or PVD of atherosclerotic origin. Additional criteria for the diagnosis of homozygous familial hypercholesterolemia (Repatha only): Total cholesterol levels > 290mg/dL or LDL-C > 190mg/dL (adults) OR Total cholesterol levels > 260mg/dL or LDL-C > 155mg/dL (children < 16 years) and TG within reference range OR Confirmation of diagnosis by gene testing.
FAMILIAL HYPERCHOLESTEROLEMIA AND HYPERTRIGLYCERIDEMIA						TRYNGOLZA	Use PA Form# 20420	Tryngolza requires fasting triglycerides of ≥ 880 mg/dL and confirmed genetically identified familial chylomicronemia syndrome (FCS).
HYPERPHAGIA - MISC								
HYPERPHAGIA - MISC				MC	8	VYKAT XR		FDA approved for the treatment of hyperphagia in adults and pediatric patients 4 years of age and older with Prader-Willi syndrome (PWS).
PULMONARY ANTI-HYPERTENSIVES								
PULMONARY ANTI-HYPERTENSIVES	MC MC/DEL MC/DEL MC		EPOPROSTENOL INJ ³ SILDENAFIL TADALAFIL VENTAVIS ³	MC/DEL MC MC/DEL MC MC MC MC MC MC/DEL MC MC MC MC MC/DEL		ADEMPAS ^{1,3} ADCIRCA ⁴ ALYQ TAB FLOLAN ³ LIQREV OPSUMIT ^{1,2} OPSYNVI ⁴ ORENITRAM REMODULIN ³ REVATIO ⁴ TADLIQ ⁴ TYVASO UPTRAVI VELVETRI ³ WINREVAIR ⁴	Use PA Form# 20420 1. Requires previous trials/failure of multiple preferred medications. 2. Dosing limits apply, see the Dose Consolidation List. 3. Require WHO Group 1 diagnosis of primary PAH (Primary Pulmonary Hypertension) and NYHA functional class 3 or 4. 4. Require WHO Group 1 diagnosis of primary PAH (Primary Pulmonary Hypertension) and NYHA functional class 2 or 3.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Sildenafil will be preferred with clinical PA for treatment of pulmonary arterial hypotension (WHO Group 1) in adults to improve exercise ability and delay clinical worsening. Avoid concomitant use of sildenafil with moderate or strong Cyp3A inhibitors. DDI: Uptravi will require a prior authorization if it is currently being used in combination with strong inhibitors of CYP2C8 (gemfibrozil). DDI: Opsumit will require a prior authorization if it is currently being used in combination with drugs known to be significant CYP3A inhibitors (ketoconazole, itraconazole, clarithromycin, indinavir, nefazodone, nelfinavir, ritonavir, atazanavir, saquinavir and telithromycin). DDI: Adempas will require a prior authorization if it is currently being used in combination with drugs known to be PDE inhibitors should be avoided (including dipyridamole, adcira and tadalafil) with adempas. Liqrev: treatment of pulmonary arterial hypertension (WHO Group 1) in adults to improve exercise ability and delay clinical worsening. Avoid concomitant use of liqrev with moderate or strong CYP3A inhibitors.
ERA / ENDOTHELIN RECEPTOR ANTAGONIST	MC MC		LETAIRIS ^{1,2} TRACLEER				Use PA Form# 20420 1. Providers must be registered with LEAP Prescribing program, a restricted distribution program. 2. Clinical PA is required to establish diagnosis and medical necessity.	Letairis approvals will require WHO Group 1 diagnosis of primary PAH (Primary Pulmonary Hypertension) and functional class 2 or 3 symptoms. Tracleer approvals will require WHO Group 1 diagnosis of primary PAH (Primary Pulmonary Hypertension) and NYHA functional class 2 thru 4. DDI: Preferred Oral Contraceptives will now be non-preferred and require prior authorization if it is currently being used in combination with tracleer.
IMPOTENCE AGENTS								
IMPOTENCE AGENTS							As of January 1, 2006, per CMS (federal govt.), impotence agents are no longer covered.	As of January 1, 2006, per CMS (federal govt.), impotence agents are no longer covered.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
ANTI-EMETOGENICS								
ANTIEMETIC - ANTICHOLINERGIC / DOPAMINERGIC	MC MC MC/D MC/D MC		DOXYLAMINE SUCC-PYRIDOXINE DOXYLAMINE SUCC-PYRIDOXINE HCL MECLIZINE HCL TABS PROMETHAZINE SUPP PROMETHAZINE TRANSDERM-SCOP PT72	MC MC MC MC MC MC MC		ANTIVERT TABS BARHEMSYS BONJESTA DICLEGIS PHENERGAN SOLN PROMETHAZINE 50MG SUPP PROMETHEGAN SUPP TORECAN TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Concomitant use of MAOIs and Bonjesta is contraindicated.
ANTIEMETIC - 5-HT3 RECEPTOR ANTAGONISTS/ SUBSTANCE P NEUROKININ	MC/D MC/D MC/D MC/D MC/D		DRONABINOLCAPS GRANISETRON TAB ONDANSETRON TAB ONDANSETRON ODT TBDP ONDANSETRON SOL	MC MC MC MC MC MC MC MC MC/D MC/D MC MC MC MC/D MC/D MC/D MC	8 8	AKYNZEO ¹ APREPITANT ALOXI ANZEMET TABS APONVIE ⁴ CESAMET ¹ CINVANTI ⁴ EMEND ² FOCINVEZ ^{1,2} KYTRIL MARINOL CAPS SANCUSO SUSTOL SYNDROS TRIMETHOBENZAMIDE CAP VARUBI ZOFRAN ODT TBDP ³ ZOFRAN TABS ³ ZOFRAN INJ ³ ZUPLENZ	Use PA Form# 20420 1. Approvals will require diagnosis of chemo induced nausea/vomiting and failed trials of all preferred anti-emetics, including 5-HT3 class (Ondansetron) and Marinol . 2. Clinical PA is required for members on highly emetic anti-neoplastic agents. 3. Dosing limits apply, see Dosage Consolidation List. 4. Clinical PA required for appropriate diagnosis.	Preferred drugs and step therapy must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. * Ondansetron limits still apply as listed on the Ondansetron PA form for covered indications including chemotherapy, radiotherapy, post operative nausea & vomiting and hyperemesis gravidarum. Other medical indications will be approved or denied on a case-by-case basis. Hyperemesis and other medical indications approved are still subject to failure of multiple preferred antiemesis drugs. Akynzeo - Concomitant use should be avoided in patients who are chronically using a strong CYP3A inducer such as rifampin. Aponvie is for the prevention of postoperative nausea and vomiting (PONV) in adults. Varubi – Available to the few who are unable to tolerate or who have failed on preferred medications.
NON-SEDATING ANTIHISTAMINES / DECONGESTANTS								
ANTIHISTIMINES - NON-SEDATING	MC MC/D MC/D MC		ALAVERT TABS CETIRIZINE TABS LORATADINE TAVIST ND (OTC)	MC MC MC/D MC/D MC/D MC MC/D MC/D MC/D MC/D	5 5 5 5 5 8 8 8 8 8 8 8 9	CLARINEX TABS ^{1,5} CLARINEX SYR ^{1,2} FEXOFENADINE ¹ ZYTEC ¹ ZYTEC SYR ^{1,2} ALLEGRA ³ CLARITIN ³ DESLORATADIN LORATADINE ODT ⁴ LEVOCETIRIZINE ⁴ XYZAL ³	Use PA Form# 20530 1. Must fail preferred drugs, OTC loratadine and cetirizine before moving to non-preferred step order drugs. 2. Clarinex and Zyrtec syrup <6 yr w/o PA. 3. Must fail all step 5 drugs (clarinex, fexofenadine and zyrtec) before moving to next step product. 4. All OTC versions of Loratadine ODT are now non-preferred. 5. PA's for Clarinex RediTabs will only be approved if between the ages of 6-11 years old.	Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. No combination product with decongestant will be approved since pseudoephedrine available without PA. Pseudoephedrine is available with prescription.
ANTIHISTIMINES - OTHER	MC/D MC/D MC/D		CLEMASTINE CHLORPHENIRAMINE DIPHENHYDRAMINE				Use PA Form# 20530	
ALLERGY / ASTHMA THERAPIES								
ANAPHYLACTIC DEVICES	MC/D MC/D MC/D		EPINEPHRINE EPIPEN EPIPEN JR	MC MC MC		AUVI- Q NEFFY TWINJECT	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ALLERGEN IMMUNOTHERAPY				MC MC MC MC MC		ODACTRA ORALAIR ¹ PALFORZIA RAGWITEK GRASTEK	Use PA Form# 20420 1. See criteria section	Prescriber must provide the testing to show that the patient is allergic to the components in the prescribed therapy and must provide a clinically valid rationale why single agent sublingual therapy is being chosen over subcutaneous therapy. Palforzia is approved for use in patients with a confirmed diagnosis of peanut allergy. Initial dose escalation may be administered to patients aged 4 through 17 years. Up-dosing and maintenance may be continued in patients 4 years of age and older. Odactra is approved for use in persons 12 through 65 years of age. Note that Odactra is not indicated for the immediate relief of allergic symptoms. Treatment must start 12 weeks before expected onset of pollen season and only after confirmed by positive skin test or in vitro testing for pollen-specific IgE antibodies for any of the 5 grass species contained in Oralair. Oralair: Patient age ≥10 years and ≤65 years. Have an auto-injectable epinephrine on-hand.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
ANTIASTHMATIC - ANTICHOLINERGICS - INHALER	MC MC/DEL MC/DEL		INCRUSE ELLIPTA ³ SPIRIVA HANDIHALER ^{1,2} SPIRIVA RESPIMAT	MC MC/DEL		LONHALA MAGNAIR TUDORZA	Use PA Form# 20420 1. Quantity limit of 1 inhalation daily (1 capsule for inhalation daily). Spiriva will require PA if Combivent or Atrovent nebulizer solution is in member's current drug profile. 2. We ask physicians to write "asthma" on the prescription whenever Spiriva is primarily being used for that condition. 3. Quantity limit of 1 inhalation daily.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIASTHMATIC - PHOSPHODIESTERASE 4 INHIBITORS	MC/DEL		ROFLUMILAST	MC/DEL MC		DALIRESP OHTUVAYRE ¹	Use PA Form# 20420 1. For the maintenance treatment of chronic obstructive pulmonary disease (COPD) in adult patients.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIASTHMATIC - ANTICHOLINERGICS - NEBULIZER	MC/DEL		IPRATROPIUM BROMIDE SOLN	MC MC/DEL		ATROVENT SOLN YUPELRI	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIASTHMATIC - ANTIINFLAMMATORY AGENTS	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		CROMOLYN SODIUM NEBU DUPIXENT ^{2,4} FASENRA ² FASENRA ² AUTO INJECT XOLAIR ¹	MC MC MC MC		CINQAIR ³ NUCALA ² TEZSPIRE ⁵	Use PA Form# 20420 1. Need max inhaled steroids and written by pulmonary or allergy specialist. Must have elevated IgE and ≥ age 6. 2. For patients with severe asthma aged 12 years or older and eosinophilia. 3. For patients ≥ 18 years of age with eosinophilia. 4. Clinical PA required. 5. For adult and pediatric patients aged 12 years and older with severe asthma.	All will require suboptimal response to maximal doses of inhaled steroid as evidenced by asthmatic ER/Hospital admissions and Allergy/Pulmonary specialist management. Dupixent limited to patient with asthma not controlled on high dose ICS-LABA who have eosinophil greater than or equal to 150 cells or the patient is depend on an oral corticosteroid. Fasenra, Nucala and Cinqair are not indicated for treatment of other eosinophilic conditions and are not indicated for the relief of acute bronchospasm or status asthmaticus.
ANTIASTHMATIC - NASAL STEROIDS	MC/DEL MC/DEL MC MC/DEL MC/DEL MC		BUDESONIDE SPRAY FLUTICASONE SPR ³ OLOPATADINE SPRAY OMNARIS SPR ³ TRIAMCINOLONE NS QNASL	MC/DEL MC/DEL MC/DEL MC MC/DEL MC MC MC MC/DEL MC MC/DEL MC	8 8 8 8 8 8 8 8 8 8 8 8 8	DYMISTA FLONASE SUSP ^{2,3} FLUNISOLIDE SOLN ^{1,3} NASONEX SUSP RHINOCORT AERO ^{2,3} RHINOCORT AQUA SUSP ^{2,3} RYALTRIS ⁴ TRI-NASAL SOLN ^{2,3} VANCENASE POKKETHALER AERS ^{2,3} VERAMYST ^{2,3} XHANCE ² ZETONNA ³	Use PA Form# 20420 1. All preferred drugs must be tried before moving to non preferred steps. 2. All step 5 medications need to be tried before moving to step 8's. 3. Dosing limits apply to whole category, see Dosage Consolidation List. 4. Use of individual ingredients or other preferred agents.	Preferred drugs and step therapy must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Xhance will be considered for the treatment of nasal polyps in patients 18 years of age or older. The patient has had a documented side effect, allergy, or treatment failure of two preferred nasal glucocorticoids, one of which must be fluticasone.
ANTIASTHMATIC - NASAL MISC.	MC/DEL MC/DEL MC		AZELASTINE CROMOLYN NASAL 4% IPRATROPIUM NASAL SOL ¹	MC/DEL MC/DEL	8 8	ASTEPRO ² PATANASE	Use PA Form# 20420 1. Ipratropium will be approved if submitted with documentation supporting use of CPAP machine. 2. Utilize Multiple preferred, as well as step therapy Azelastine.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Approved if patient fails on nonsedating antihistamines and steroid nasal sprays.
ANTIASTHMATIC - BETA - ADRENERGICS	MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC MC/DEL MC/DEL MC MC/DEL		ALBUTEROL NEB ALBUTEROL HFA (Teva labeler 00093 AND Sandoz 00781) LEVALBUTEROL TARTRATE METAPROTERENOL PROAIR RESPICLICK PROVENTIL HFA SEREVENT STRIVERDI TERBUTALINE SULFATE TABS ALBUTEROL 0.63mg/3ml VENTOLIN HFA AERS PROAIR DIGIHALER ⁴	MC/DEL MC/DEL MC/DEL MC MC MC MC/DEL MC/DEL MC MC/DEL		ACCUNEB NEBU ALBUTEROL HFA BRETHINE VOLMAX TBCR VOSPIRE ER TB12 XOPENEX HFA ³ XOPENEX NEBU ^{1,2}	Use PA Form# 20420 1. Xopenex users w/ prior asthma hospitalization due to albuterol nebulizer failure will be grandfathered. 2. Quantity Limit: 12 cc/day.. 3. Dosing limits apply, see Dosage Consolidation List. 4. For the treatment of patients ≥ 4 years of age.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
ANTIASTHMATIC - ADRENERGIC COMBINATIONS	MC MC MC MC MC/DEL MC/DEL MC/DEL		ADVAIR DISKUS ¹ ADVAIR HFA ¹ AIRDUO RESPICLICK ² BREO ELLIPTA ¹ DULERA FLUTICASONE-SALMETEROL SYMBICORT	MC MC/DEL MC/DEL MC		AIRDUO DIGIHALER ² AIRSUPRA BREZTRI AEROSPHERE TRELEGY ELLIPTA ¹	Use PA Form# 20420 1. Dosing limits apply, see Dosage Consolidation List. 2. For patients ≥ 12 years and older.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. AirDuo® Respiclick be non-preferred and require prior authorization and be available to those who are unable to tolerate or who have failed on preferred medications. DDI: Avoid concomitant use of strong CYP3A4 inhibitors (e.g. ritonavir, atazanavir, clarithromycin, indinavir, itraconazole, nefazodone, nelfinavir, saquinavir, ketoconazole, telithromycin) with AirDuo® Respiclick is not recommended due to increased systemic corticosteroid and increased cardiovascular adverse effects.
ANTIASTHMATIC - ADRENERGIC ANTICHOLINERGIC	MC/DEL MC MC/DEL MC/DEL		ALBUTEROL/IPRATROPIUM NEB. SOLN ANORO ELLIPTA COMBIVENT RESPIMAT STIOLTO	MC/DEL MC/DEL MC/DEL		BEVESPI AEROSPHERE ^{2,3} DUAKLIR PRESSAIR DUONEB SOLN ¹	Use PA Form# 20420 1. Please use preferred individual ingredients Albuterol and Ipratropium. 2. Dosing limits apply, see Dosing Consolidation List. 3. The safety and efficacy of use in children under the age of 18 years have not been established.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DuoNeb components are available separately without PA. Bevespi should be used with extreme caution in patients being treated with MAO inhibitors, TCAs, or other drugs known to prolong the QTc interval. DDI: Avoid concomitant use of Bevespi with other anticholinergic-containing drugs, due to an increased risk of anticholinergic adverse events. Bevespi should be used with extreme caution in patients being treated with MAO inhibitors, TCAs, or other drugs known to prolong the QTc interval.
ANTIASTHMATIC - XANTHINES	MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL		AMINOPHYLLINE TABS THEOCHRON TB12 THEOLAIR-SR TB12 THEOPHYLLINE CR TB12 THEOPHYLLINE ELIX THEOPHYLLINE SOLN THEOPHYLLINE ER CP12 THEOPHYLLINE ER TB12	MC/DEL MC MC/DEL		THEO-24 CP24 THEOLAIR TABS UNIPHYL TBCR	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIASTHMATIC - STEROID INHALANTS	MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC		ARNUITY ELLIPTA ASMANEX TWISTHALER ^{3,4} ASMANEX HFA ⁵ BUDESONIDE NEB 0.25MG & 0.5MG ¹ PULMICORT FLEXHALER ³ QVAR AERS ³	MC MC/DEL MC MC/DEL MC/DEL	8 8 8 8 8	AEROSPAN ALVESCO ³ ARMONAIR DIGIHALER BUDESONIDE NEB 1MG PULMICORT SUSP	Use PA Form# 20420 1. Budesonide Neb 0.25mg & 0.5mg will be preferred for members under the age of 8 years old. PA will be required for members 8 years of age and older, please consider other preferred options. 2. All preferred must be tried before moving to non preferred steps. 3. Dosing limits apply, see Dosage Consolidation List. 4. Asmanex 110mcg will be limited to member between the ages of 4-11 years old. 5. Asmanex HFA will be preferred for members under the age of 6 years old. PA will be required for members 6 years of age and older, please consider other preferred options.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIASTHMATIC - 5-Lipoxygenase Inhibitors				MC		ZYFLO CR TABS	Use PA Form# 20420	Other Preferred asthma controller drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIASTHMATIC - LEUKOTRIENE RECEPTOR ANTAGONISTS	MC/DEL MC/DEL MC/DEL		MONTELUKAST GRANULE ¹ MONTELUKAST SODIUM TAB MONTELUKAST SODIUM CHEW TAB	MC/DEL MC/DEL MC/DEL	8 8 8	ACCOLATE TABS SINGULAIR ² SINGULAIR GRANULES	Use PA Form# 20420 1. Montelukast Granules will only be approved if between ages of 6 months-24 months. 2. Singulair Chewable 4mg from 2 years- 5 years and Singulair Chewable 5mgs from 6 years- 14 years old.	
ANTIASTHMATIC - ALPHA-PROTEINASE INHIBITOR				MC MC/DEL MC MC	8 8 8 8	ARALAST ZEMAIRA GLASSIA PROLASTIN SUSR	Use PA Form# 20420	Prolastin and Azemaira will be approved for members with A1AT deficiency and clinically demonstrable panacinar emphysema.
ANTIASTHMATIC - HYDRO-LYTIC ENZYMES				MC/DEL		PULMOZYME SOLN	Use PA Form# 20420	Will be approved for cystic fibrosis patients.
ANTIASTHMATIC - MUCOLYTICS	MC/DEL		ACETYLCYSTEINE ¹	MC		MUCOMYST	Use PA Form# 20420 1. Acetylcysteine is covered with diagnosis of CF.	

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
ANTIASTHMATIC-CFTR POTENTIATOR AND COMBINATIONS				MC MC MC MC MC/DEL		ALYFTREK BRONCHITOL ¹ KALYDECO ORKAMBI SYMDEKO TRIKAFTA	Use PA Form# 20420 1. For the treatment of patients ≥18 years of age with CF.	<p>Alftytrek will be considered for the treatment of patients 6 years and older with at least one responsive mutation, including 31 additional mutations not responsive to other CFTR modulator therapies.</p> <p>Bronchitol will be considered as add-on maintenance therapy to improve pulmonary function in adult patients 18 years and older with cystic fibrosis (CF). Use Bronchitol only for adults who have passed the Bronchitol Tolerance Test (BTT). (see Recommended Dosage section for further information).</p> <p>Kalydeco will be considered for patients with cystic fibrosis (CF) aged 1 month and older who have at least one mutation in the CFTR gene that is responsive to ivacaftor potentiation based on clinical and/or in vitro assay data. If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to detect the presence of a CFTR mutation followed by verification with bi-directional sequencing when recommended by the mutation test instructions for use.</p> <p>Orkambi will be considered for patients with cystic fibrosis (CF) aged 1 year and older who are homozygous for the F508del mutation in the CFTR gene. If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to detect the presence of the F508del mutation on both alleles of the CFTR gene. The efficacy and safety of Orkambi have not been established in patients with CF other than those homozygous for the F508del mutation.</p> <p>Symdeko will be considered for patients with cystic fibrosis (CF) aged 6 years and older who are homozygous for the F508del mutation or who have at least one mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to tezacaftor/ivacaftor based on in vitro data and/or clinical evidence. If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to detect the presence of a CFTR mutation followed by verification with bi-directional sequencing when recommended by the mutation test instructions for use.</p> <p>Trikafta will be considered for the treatment of cystic fibrosis (CF) in patients aged 2 years and older who have at least one F508del mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene or mutation in the CFTE gene that is responsive based on in vitro data. If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to confirm the presence of at least one F508del mutation or a mutation that is responsive based on in vitro data.</p>
IDIOPATHIC PULMONARY FIBROSIS	MC/DEL		OFEV ¹	MC MC		ESBRIET ¹ PIRFENIDONE	Use PA Form# 20420 1. Diagnosis required	<p>Ofev- Avoid concomitant use with P-gp and CYP4A inducers (e.g. carbamazepine, phenytoin, and St. John's wort.</p> <p>Esbriet- The concomitant use with strong CYP1A2 inhibitors (e.g. fluvoxamine, enoxacin) is not recommended.</p>
COUGH/COLD								
COUGH/COLD	MC/DEL MC/DEL MC/DEL MC/DEL MC MC		DEXTROMETHORPHAN CAPS ¹ DEXTRO-GUAIF SYRP ¹ GUAIFENESIN SYRP ¹ PSEUDOEPHEDRINE ¹ ROBITUSSIN DM SYRP ¹ ROBITUSSIN SUGAR FREE SYRP ¹				Use PA Form# 20420 1. All of cough cold preparations are not covered except these preferred products.	All non-preferred products are not covered as permitted by Federal Medicaid regulations and MaineCare Policy.
DIGESTIVE AIDS / ASSORTED GI								
GI - ANTIPERISTALTIC AGENTS	MC/DEL MC/DEL MC/DEL MC/DEL MC		DIPHENOXYLATE DIPHENOXYLATE/ATROPINE LOPERAMIDE HCL CAPS/LIQ OPIUM TINCTURE TINC PAREGORIC TINC	MC/DEL MC MC		LOFENE TABS LONOX TABS MOTOFEN TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval.
GI - ANTI-DIARRHEAL/ ANTACID - MISC.	MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		ATROPINE SULFATE SOLN BISMATROL BISMUTH SUBSALICYLATE CALCIUM CARBONATE (ANTACID) CHEW DICYCLOMINE HCL GLYCOPYRROLATE TABS HYOSCYAMINE CAPS & TABS HYOSCYAMINE SULFATE KAOPECTATE MAGNESIUM OXIDE TABS MAG-OX 400 TABS PAMINE TABS PROPANTHELINE BROMIDE TABS SODIUM BICARBONATE TABS TUMS	MC/DEL MC/DEL MC/DEL MC MC MC MC/DEL MC MC MC MC MC MC MC		BELLADONNA ALKALOIDS & OP BENTYL TABS BENTYL SYRP CUVPOSA DARTISLA ODT ² ED-SPAZ MYTESI ¹ GLYCOPYRROLATE INJ LEVSIN TABS LEVSIN/SL SUBL NULEV TBDP OSCIMIN ROBINUL INJ ROBINUL TABS	Use PA Form# 20420 1. Dosing limits apply, see Dose Consolidation List. 2. It is not indicated as monotherapy for treatment of peptic ulcer because effectiveness in peptic ulcer healing has not been established.	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval.</p> <p>Preferred products that used to require diag codes still require diag codes unless indicated otherwise.</p> <p>Mytesi requires a diagnosis of non-infectious diarrhea in patients with HIV/AIDS on anti-retroviral therapy, prior trials of preferred, more cost effective anti-diarrheal.</p>
GI- BILE ACID				MC MC		CHOLBAM ¹ CTEXLI¹	Use PA Form# 20420 1. Clinical PA is required to establish diagnosis and medical necessity.	Indication of bile acid synthesis disorders due to single enzyme defects (SEDs) AND for adjunctive treatment of peroxisomal disorders (PDs).
GI- EOSINOPHILIC ESOPHAGITIS	MC		EOHILIA ¹				Use PA Form# 20420 1. Approvals will not be longer than 12 weeks of treatment in adult and pediatric patients 11 years of age and older.	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p> <p>Eohilia: Dietary modification, PPIs, and topical glucocorticoids are required as initial therapy.</p>
GI - H2-ANTAGONISTS	MC MC/DEL MC/DEL		ACID REDUCER TABS CIMETIDINE FAMOTIDINE	MC MC MC/DEL MC		AXID CAPS AXID AR TABS NIZATIDINE CAPS PEPCID PEPCID AC	Use PA Form# 20420	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p> <p>DDI: Cimetidine will now be non-preferred and require prior authorization if it is currently being used with any sulfonylurea (except for glyburide).</p> <p>DDI: Cimetidine will require prior authorization if being used in combination with plavix.</p>

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
GI- IBAT INHIBITORS				MC MC		BYLVAY ^{1,2} LIVMARLI ^{1,2}	Use PA Form# 20420 1. For the treatment of patients ≥ 3 months of age. 2. Clinical PA required for appropriate diagnosis.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval.
GI - PROTON PUMP INHIBITOR	MC/DEL MC/DEL MC/DEL		OMEPRAZOLE CAPS ² PANTOPRAZOLE ² LANSOPRAZOLE CAPS ²	MC/DEL MC/DEL MC MC MC/DEL MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	6 6 7 7 8 8 8 8 8 8 8 8 8 8	NEXIUM CPDR ³ NEXIUM SUS ⁵ PRILOSEC OTC ³ ACIPHEX TBEC ³ DEXILANT (KAPIDEX) ² KONVOME ² OMEPRAZOLE-SODIUM BICARBONATE CAPS OMEPRAZOLE MAGNESIUM PREVACID CPDR ³ PREVACID SOLUTABS ^{1,4} PRILOSEC CPDR PROTONIX INJ PROTONIX ² VOQUEZNA TABS	Use PA Form# 20720 1. Prevacid Solutabs available without PA for child less than 9 years old. 2. Dosing limits apply, please see Dosage Consolidation List. 3. All preferred and step therapy must be tried and failed. 4. Payment for Prevacid SoluTabs for patients 9 and older will be considered for those patients who cannot tolerate a preferred solid oral dosage form. 5. Nexium sus available without PA if member is < 12 yrs of age and ≤ 1 pack per day.	All preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Please refer to the PPI PA form for additional criteria on Non-Preferred PPIs. DDI: Omeprazole will require prior authorization if being used in combination with plavix. DDI: Lansoprazole will require prior authorization if being used in combination with plavix. DDI: Prevacid, Omeprazole and Pantoprazole will now be non-preferred and require prior authorization if they are currently being used in combination with any of the following medications: ampicillin, B-12, fe salts, griseofulvin, sporanox, ketoconazole, reyataz, or vantin. DDI: All non-preferred PPIs require prior authorization, but with any prior authorization request, the member's drug profile will also be monitored for current use with ampicillin, B-12, Fe salts, griseofulvin, itraconazole, ketoconazole, reyataz or vantin due to a significant drug-drug interaction.
GI - ULCER ANTI-INFECTIVE	MC MC		PYLERA TALICIA			VOQUEZNA DUAL PAK VOQUEZNA TRIPLE PAK	Use PA Form# 20420	
GI - PROSTAGLANDINS	MC		MISOPROSTOL TABS	MC/DEL		CYTOTEC TABS	Use PA Form# 20420	Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drug will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
GI - DIGESTIVE ENZYMES	MC/DEL MC		CREON ¹ ZENPEP ¹	MC/DEL MC/DEL MC/DEL		PERTZYE ULTRESA VIOKACE	Use PA Form# 20420 1. Clinical PA is required to establish CF diagnosis and medical necessity. In all cases except cystic fibrosis patients, objective evidence of pancreatic insufficiency (fat malabsorption test etc) must be supplied.	Non -Preferred drugs must be tried and failed in step-order due to lack of efficacy or intolerable side effects before other non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
GI - ANTI - FLATULENTS / GI STIMULANTS	MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL		AMITIZA CALULOSE SYRP CONSTULOSE SYRP ENULOSE SYRP GASTROCROM CONC GENERLAC SYRP LACTULOSE SYRP METOCLOPRAMIDE HCL	MC MC/DEL MC MC/DEL		CEPHULAC SYRP INFANTS GAS RELIEF SUSP GIMOTI SPRAY REGLAN TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval.
GI - INFLAMMATORY BOWEL AGENTS	MC MC/DEL MC MC MC/DEL MC/DEL		APRISO BALSALAZIDE MESALAMINE ENMA KIT PENTASA SULFAZINE EC TBEC SULFASALAZINE TABS	MC/DEL MC/DEL MC MC MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC MC MC		ASACOL 800MG HD AZULFIDINE EN-TABS TBEC AZULFIDINE TABS COLAZAL CAPS DELZICOL DIPENTUM CAPS GIAZO LIALDA TABS ¹ MESALAMINE TAB ROWASA ENEM SFROWASA UCERIS RECTAL FOAM ² UCERIS TABS ²	Use PA Form# 20420 1. Current users grandfathered. 2. Diagnosis required.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Giazo is only indicated for males, as the safety efficacy for use in females has not been established. Prior trials of preferred products. Uceris Rectal Foam or Tab- Concomitant use with CYP3A inhibitors (e.g. ketoconazole, itraconazole, ritonavir, indinavir, saquinavir, erythromycin, cyclosporine, and grapefruit juice) should be avoided. Verify prior trials and failures or intolerance of preferred treatments.
GI - IRRITABLE BOWEL SYNDROME AGENTS	MC/DEL		LOTRONEX TABS	MC		VIBERZI	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
GI- SHORT BOWL SYNDROME				MC		GATTEx	Use PA Form# 20420	Gattex requires a diagnosis of adult SBS who are dependent on parenteral support. Appropriate colonoscopy and lab assessments 6months prior to starting.
GI- NASH				MC		REZDIFFRA	Use PA Form# 20420	Rezdiffra: The patient must have a diagnosis of NASH with fibrosis Stage 2 or 3 and utilizing imaging and scanning test such as fibro scan, MRI or ultra sound AND the patient does not have evidence of decompensated cirrhosis.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
MISCELLANEOUS GI								
GI - MISC.	MC/DEL		BISAC-EVAC SUPP	MC/DEL		ACTIGALL CAPS	Use PA Form# 20420 2. For the treatment of carcinoid syndrome diarrhea in combination with somatostatin analog (SSA) therapy in adults inadequately controlled by SSA therapy. 3. For the treatment of Opioid Induced Constipation (OIC). 5. Dosing limits apply, see Dose Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval. Trulance should be avoided in pediatric patients less than 18 years of age. Iqirvo: For the treatment of primary biliary cholangitis (PBC) in combination with ursodeoxycholic acid (UDCA) in adults who have had an inadequate response to UDCA, or as monotherapy in patients unable to tolerate UDCA. Continued approval for this indication may be contingent upon verification and description of clinical benefit in confirmatory trial(s). Livdelzi: Clinical PA is required for the treatment of primary biliary cholangitis (PBC) in combination with ursodeoxycholic acid (UDCA) in adults who have had an inadequate response to UDCA, or as monotherapy in patients unable to tolerate UDCA. Patients who do not have a diagnosis of decompensated cirrhosis.
	MC/DEL		BISACODYL	MC		BENEFIBER		
	MC		BISCOLAX SUPP	MC/DEL		CARAFATE		
	MC		CINOBAC CAPS	MC/DEL		CLEARLAX POW		
	MC/DEL		CITRATE OF MAGNESIA SOLN	MC/DEL		COLACE CAPS		
	MC/DEL		CITRUCEL	MC		DIOCTO-C SYRP		
	MC/DEL		CLENPIQ SOL	MC		DOC SOD /CAS CAP		
	MC/DEL		COLYTE	MC		DOC-Q-LAX CAPS		
	MC/DEL		DIOCTO SYRP	MC/DEL		DOCUSATE SODIUM/CAS CAPS		
	MC		DOCUSATE CALCIUM CAPS	MC/DEL		DOK PLUS		
	MC/DEL		DOCUSATE SODIUM	MC/DEL		DULCOLAX SUPP		
	MC/DEL		FIBER LAXATIVE TABS	MC		ENEMEEZ		
	MC		FLEET	MC		FIBER CON TABS		
	MC/DEL		GENFIBER POWD	MC/DEL		FIBER-LAX TABS		
	MC/DEL		GLYCERIN	MC/DEL		GAVILYTE-H		
	MC		HIPREX TABS	MC		GOLYTELY SOLR		
	MC/DEL		KRISTALOSE PACK	MC		IBSRELA		
	MC/DEL		LINZESS ⁵	MC		IQIRVO		
	MC		MAALOX	MC		LIVDELZI		
	MC/DEL		MILK OF MAGNESIA SUSP	MC		MALTSUPEX		
	MC		MINERAL OIL	MC		MIRALAX PACKETS		
	MC		MIRALAX BULK POWD (BRAND)	MC/DEL		MOTEGRITY		
	MC/DEL		MOVANTIK	MC		PEG-ELECTROLYTES SOLR		
	MC/DEL		MOVIPREP POWD PACK	MC		PEG 3350 PACKETS		
	MC		PEG 3350- ELECTROLYTE SOL	MC		PREPOPIK PAK		
	MC		PEG 3350 POWDER	MC		RELISTOR TABS		
	MC/DEL		SENNA	MC/DEL		SENEXON TABS		
	MC/DEL		SENOKOT GRAN	MC/DEL		SENOKOT TABS		
	MC/DEL		SENOKOT SYRP	MC		SENOKOT S TABS		
	MC/DEL		SENOKOT CHILDRENS SYRP	MC/DEL		SORBITOL		
	MC		SENOKOT XTRA TABS	MC		STOOL SOFTENER PLUS CAPS		
	MC/DEL		STOOL SOFTENER CAPS	MC		SUFLAVE		
	MC/DEL		SUCRALFATE TABS	MC		SUTAB		
	MC/DEL		SUPREP SOL	MC/DEL		SYMPROIC ³		
	MC		TRULANCE ²	MC/DEL		UNI-CENNA TABS		
	MC		UNI-EASE CAPS	MC		UNI-EASE PLUS CAPS		
	MC		URSO FORTE	MC		V-R NATURAL SENNA LAXATIV TABS		
	MC/DEL		URSODIOL	MC		URSO 250		
				MC		XERMELO ²		
MISC. UROLOGICAL								
UROLOGICAL - MISC.	MC		ACETIC ACID 0.25% SOLN	MC		CITRIC ACID/SODIUM CITRAT SOLN	Use PA Form# 20420 1. Elmiron requires adequate proof of Dx with supportive testing.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC		CYTRA-K SOLN	MC/DEL		CYTRA-2 SOLN		
	MC		FOSFOMYCIN (NDC 82036427401 ONLY)	MC/DEL		ELMIRON CAPS ¹		
	MC		K-PHOS MF TABS	MC		FURADANTIN SUSP		
	MC/DEL		METHENAMINE MANDELATE TABS	MC/DEL		MACROBID CAPS		
	MC/DEL		NEOSPORIN GU IRRIGANT SOLN	MC/DEL		MACRODANTIN CAPS		
	MC/DEL		NITROFURANTOIN MONO CAPS	MC/DEL		NITROFURANTOIN MACR SUSP		
	MC/DEL		PHENAZOPYRIDINE HCL TABS	MC		POTASSIUM CITRATE/CITRIC SOLN		
	MC/DEL		PHENAZOPYRIDINE PLUS	MC/DEL		PYRIDIUM PLUS TABS		
	MC		POT CITRATE TAB	MC		PYRIDIUM TABS		
	MC/DEL		PROSED/DS TABS	MC/DEL		RENACIDIN SOLN		
	MC		TRICITRATES SYRP	MC		UROCIT-K		
	MC/DEL		URELIEF PLUS					
	MC		UREX TABS					
	MC/DEL		URISED TABS					
	MC/DEL		UROQID #2 TABS					

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
PHOSPHATE BINDERS								
PHOSPHATE BINDERS	MC/DEL MC/DEL MC/DEL MC MC/DEL		CALCIUM ACETATE CAP ¹ FOSRENOL CHEW ¹ MAGNEBIND - 400 ¹ PHOSLYRA ¹ REVELA ¹	MC MC/DEL MC/DEL MC/DEL MC MC		AURYXIA ¹ CALCIUM ACETATE TAB ¹ ELIPHOS ¹ FOSRENOL PWDR ¹ VELPHORO ¹ XPHOZAH	Use PA Form# 20420 1. Diagnosis required.	Preferred drugs must be tried and failed in step-order due to lack of efficacy or intolerable side effects before less preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Xphozah to reduce serum phosphorus in adults with chronic kidney disease (CKD) on dialysis as add-on therapy in patients who have an inadequate response to phosphate binders or who are intolerant of any dose of phosphate binder therapy.
INTRA-VAGINALS								
VAGINAL - ANTIBACTERIALS	MC/DEL MC/DEL MC MC/DEL MC/DEL		CLEOCIN CREA CLEOCIN SUPP CLINDESSE CREA METRONIDAZOLE VAGINAL GEL ¹ NUVESSA	MC/DEL MC/DEL MC 		METROGEL VAGINAL GEL ¹ VANDAZOLE XACIATO	Use PA Form# 20420 1. Dosing limits apply, see Dosage Consolidation List.	Preferred drugs must be tried and failed in step-order due to lack of efficacy or intolerable side effects before less preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
VAGINAL - ANTI FUNGALS	MC/DEL MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC MC/DEL MC MC		CLOTRIMAZOLE CREA CLOTRIMAZOLE-3 CREA GYNE-LOTRIMIN CREA MICONAZOLE CREA MICONAZOLE 3 KIT CREA OTC MICONAZOLE 7 CREA MICONAZOLE NITRATE CREA NYSTATIN TABS TERCONAZOLE CREAM VAGITROL V-R MICONAZOLE-7 CREA	MC MC MC MC MC/DEL MC/DEL MC MC MC/DEL MC/DEL		AVC CREA CLOTRIMAZOLE 3 DAY CREA GYNAZOLE-1 CREA GYNE-LOTRIMIN 3 TABS MICONAZOLE 3 COMBO PACK KIT ¹ MICONAZOLE 3 SUPP TERAZOL 3 CREA TERAZOL 7 CREA TERCONAZOLE SUPP	Use PA Form# 20420 1. Quantity limit: 1/script/2 weeks.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Miconazole will require prior authorization if being used in combination with warfarin.
VAGINAL - CONTRACEPTIVES							Use PA Form# 20420	Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drug will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
VAGINAL - ESTROGENS	MC/DEL MC/DEL		ESTRING RING PREMARIN CREA	MC/DEL MC/DEL		ESTRACE CREA ¹ VAGIFEM TABS ¹	Use PA Form# 20420 1. Must fail all preferred products before non-preferred.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
VAGINAL - OTHER	MC/DEL MC MC		ACID JELLY GEL ACI-JEL GEL CERVICAL AMINO ACID CREA	MC 		AMINO ACID CERVICAL CREA	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
BENIGN PROSTATIC HYPERPLASIA (BPH)								
BPH	MC/DEL MC/DEL MC/DEL MC/DEL		DOXAZOSIN MESYLATE TABS FINASTERIDE ¹ 5mg TERAZOSIN HCL CAPS TAMSULOSIN HCL	MC/DEL MC/DEL MC MC/DEL MC MC MC/DEL MC/DEL MC/DEL	5 8 8 8 8 8 8 8 8	FLOMAX CP24 ALFUZOSIN AVODART ^{2,4} CARDURA TABS ⁴ ENTADFI ^{5,6} JALYN ^{3,4} PROSCAR TABS ⁴ RAPAFLO ⁴ TEZRULY UROXATRAL ⁴	Use PA Form# 20420 1. There will be dosing limits of 1 tab per day with out PA. 2. Prior use of preferred agent prior to any approvals. 3. Use of preferred (Tamsulosin and Finasteride) and (Tamsulosin and non-preferred Avodart). 4. Non-preferred products must be used in specified order. 5. Use of individual ingredients preferred (Finasteride and Tadalafil). 6. Entadfi is not recommended for more than 26 weeks.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Approval of a non-preferred 5-alpha reductase inhibitor requires objective clinical evidence of a very enlarged prostate rather than just the presence of obstructive urinary outflow symptoms along with adequate trial of preferred proscar.
ANXIOLYTICS								
ANXIOLYTICS - BENZODIAZEPINES	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		ALPRAZOLAM TABS CHLORDIAZEPOXIDE HCL CAPS CLORAZEPATE DIPOTASSIUM TABS DIAZEPAM LORAZEPAM OXAZEPAM CAPS	MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	8 8 8 8 8 8 8 9	ALPRAZOLAM ER ATIVAN LOREEV XR NIRAVAM SERAX TRANXENE XANAX TABS XANAX XR	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
	MC		PROTRIPTYLINE HCL TABS ¹	MC		TOFRANIL		
	MC		SURMONTIL CAPS ¹	MC		VIVACTIL TABS		
SEDATIVE / HYPNOTICS								
SEDATIVE/HYPNOTICS - BARBITURATE	MC MC/DEL MC MC/DEL		BUTISOL SODIUM TABS ¹ CHLORAL HYDRATE SYRP ¹ MEBARAL TABS ¹ PHENOBARBITAL ¹	MC MC/DEL MC MC/DEL		LUMINAL SOLN SOMNOTE CAPS	Use PA Form# 20420 1. PA required for new users of preferred products if over 65 years.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
SEDATIVE/HYPNOTICS - BENZODIAZEPINES	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		DORAL TABS ¹ ESTAZOLAM TABS ¹ FLURAZEPAM HCL CAPS ¹ TEMAZEPAM CAPS 15 & 30MG ¹ TRIAZOLAM TABS ¹	MC MC MC/DEL MC/DEL MC/DEL		HALCION TABS ¹ MIDAZOLAM HCL SYRP RESTORIL CAPS ¹ TEMAZEPAM 7.5MG ¹	Use PA Form# 30110 1. Dosing limits apply, see Dosing Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Benzodiazepines do cause dependence with continued use and usage should be limited to 7-10 days at a time. Chronic intermittent use (2-3 days per week max) is the standard of care.
SEDATIVE/HYPNOTICS - Non-Benzodiazepines	MC/DEL MC MC/DEL MC/DEL	1 1 1 2	MIRTAZAPINE TRAZODONE ZOLPIDEM ² ZALEPLON ^{2,3}	MC/DEL MC/DEL MC/DEL MC/DEL MC MCDEL MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL	7 7 7 8 8 8 8 8 8 8 8 8 8 8 8	AMBIEN ¹ ESZOPICLONE ZOLPIDEM ER AMBIEN CR ¹ BELSOMRA ¹ DAYVIGO ¹ EDLUAR HETLIOZ INTERMEZZO LUNESTA ¹ SONATA CAPS ¹ ROZEREM QUVIVIQ ZOLPIMIST	Use PA Form# 30110 1. Quantity Limit of 12 per 34 days. 2. Quantity limits will be allowed up to 30/30, but intermittent therapy is recommended. 3. Only Zolpidem trial/failure will be required to obtain Zaleplon. 4. Must fail all preferred products before non-preferred.	Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Ambien, Ambien CR, Lunesta, Sonata, Zaleplon and Zolpidem may cause dependence with continued use and as with benzodiazepines, usage should be limited to 7-10 days at a time. Chronic intermittent use (2-3 days per week max) is the standard of care. Please refer to Sedative/Hypnotic PA form. DDI: Belsomra with strong CYP3A inhibitors (e.g. ketoconazole, itraconazole, posaconazole, clarithromycin, nefazodone, ritonavir, saquinavir, nelfinavir, indinavir, boceprevir, telaprevir, telithromycin, and conivaptan) is not recommended.
ANTI-PSYCHOTICS								
ANTIPSYCHOTICS - ATYPICALS	MC MC MC/DEL MC MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL		ABILIFY ASIMTUFII ABILIFY MAINTENA ARIPIRAZOLE TAB ³ ARISTADA ARISTADA INITIO OLANZAPINE ^{2,3} OLANZAPINE ^{2,3} ODT INVEGA HAFYERA INVEGA SUSTENNA INVEGA TRINZA INJ LURASIDONE TAB PALIPERIDONE ER PERSERIS RISPERDAL CONSTA RISPERIDONE ODT RISPERIDONE TAB ^{2,3} RISPERIDONE SOLN ² RYKINDO QUETIAPINE ^{2,3} QUETIAPINE XR VRAYLAR ⁴ ZIPRASIDONE ^{2,3}	MC/DEL MC MC/DEL MC/DEL MC MC MC MC MC/DEL MC MC MC MC MC MC MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL	8 9	ABILIFY DISC TAB, INJ and SOL ¹ ABILIFY TABS ² ARIPIRAZOLE SOL ARIPIRAZOLE ODT CAPLYTA COBENFY ERZOFRI FANAPT GEODON INVEGA IGALMI LATUDA LYBALVI NUPLAZID OPIPZA REXULTI RISPERDAL TAB RISPERDAL M TAB ¹ RISPERDAL SOLN SAPHRI ¹ SECUADO SEROQUEL TABS UZEDY ZYPREXA TABS ZYPREXA RELPREVV ZYPREXA ZYDIS TBDP ¹ SEROQUEL XR	Use PA form# 20440 for Multiple Antipsychotic requests Use PA form# 10130 for non-preferred single therapy atypical requests If prescribing 2 or more antipsychotics, PA will be required for both drugs, except if one is Clozapine . This includes combination of Seroquel with Seroquel XR . 1. Established users of single therapy atypicals were grandfathered. 2. Prior Authorization will be required for preferred medications for members under the age of 5. 3. Dosing limits apply, refer to the Dose Consolidation List. 4. Requires step through 1 preferred drug for all indications except AMDD. AMDD requires insufficient response from two antidepressants.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Non preferred atypicals will be approved for patients with FDA-approved indications, and for specific conditions supported by at least two published peer-reviewed double-blinded, placebo-controlled randomized trials that are not contradicted by other studies of similar quality and as long as all first line preferred therapies have been tried and failed at full therapeutic doses for adequate durations (at least two weeks). Quetiapine prescriptions for are limited to a maximum daily dose of 800mg. Uzedy: Establish tolerability with oral risperidone prior to initiating Uzedy. Atypicals: Prior Authorization will be required for preferred medication to assure indication is in accordance with FDA approved or literature supported evidence-based best practices. The approved indications are: <ul style="list-style-type: none">• schizophrenia• bipolar disorder• agitation related to autism• adjunct in major depressive disorder Lybalvi: Step through aripiprazole and latuda. If criteria is met then initial approval for 3 months. Subsequent approvals will be based on evidence of not gaining >= 10% baseline body weight for ongoing approval. If weight gain >= 10% of initial body weight, then criteria for ongoing use not met. Cobenfy: Patient must be 18–65 years old AND meet criteria for the diagnosis of schizophrenia, AND trial of 2 prior preferred second generation antipsychotics showing minimal response in control of symptoms of schizophrenia OR trial of SGA that have yielded side effects of weight gain which has not been responsive to lifestyle & medication augmentation AND patient must have baseline tests including heart rate, liver enzymes, kidney function tests, and bilirubin prior to starting treatment. Invega Hafyera: The patient is started and stabilized on the medication OR the patient has been adequately treated with Invega Sustenna (paliperidone palmitate 1-month) for at least four months or Invega Trinza (paliperidone palmitate 3- month) following at least one 3-month injection cycle. DDI: It is recommended to reduce the Vraylar dose if it is used concomitantly with a strong CYP3A inhibitor (such as itraconazole, ketoconazole). The concomitant use of Vraylar with a CYP3A4 inducer (such as rifampin, carbamazepine) is not recommended. DDI: The concomitant use of Nuplazid with other drugs known to prolong the QT interval (e.g. Class IA antiarrhythmics, Class 3 antiarrhythmics, antipsychotics, and antibiotics such as gatitoxacin and moxifloxacin).
ANTIPSYCHOTICS - SPECIAL ATYPICALS	MC/DEL		CLOZAPINE TABS	MC/DEL MC/DEL MC/DEL		CLOZAPINE ODT CLOZARIL TABS VERSACLOZ SUSP	Use PA Form# 20420	Preferred generic drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred brand will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Patients previously stabilized on brand name drug will be approved.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
ANTIPSYCHOTICS - TYPICAL	MC/DEL MC/DEL MC/DEL MC MC/DEL MC MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL		CHLORPROMAZINE HCL FLUPHENAZINE DECANOATE FLUPHENAZINE HCL HALDOL HALOPERIDOL HALOPERIDOL DECANOATE SOLN HALOPERIDOL LACTATE SOLN LOXAPINE SUCCINATE CAPS LOXITANE-C CONC MOBAN TABS PERPHENAZINE PROCHLORPERAZINE SERENTIL THIORIDAZINE HCL THIOTHIXENE TRIFLUOPERAZINE HCL TABS	MC/DEL MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL		COMPAZINE COMPRO SUPP FLUPHENAZINE HCL CONC HALDOL DECANOATE LOXITANE CAPS MELLARIL NAVANE CAPS PROLIXIN STELAZINE TABS	Use PA Form# 20420 If prescribing 2 or more antipsychotics, PA will be required for both drugs, except if one is Clozapine .	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. If prescribing 2 or more antipsychotics, PA will be required for both drugs, except if one is Clozapine .
LITHIUM								
LITHIUM	MC/DEL MC/DEL		LITHIUM CARBONATE LITHIUM CITRATE SYRP	MC/DEL MC/DEL		ESKALITH CAPS ESKALITH CR TBCR	Use PA Form# 20420	
COMBINATION - PSYCHOTHERAPEUTIC								
PSYCHOTHERPEUTIC COMBINATION				MC/DEL MC/DEL		CHLORDIAZEPOXIDE/AMITRIPT PERPHENAZINE/AMITRIPTYLIN	Use PA Form# 20420	
STIMULANTS								
STIMULANT - AMPHETAMINES - SHORT ACTING	MC/DEL MC/DEL MC		AMPHETAMINE SALT COMBO ^{1,3,4} DEXTROAMPHET SULF TABS ^{1,2,3} PROCENTRA ^{1,3}	MC/DEL MC MC/DEL MC		ADDERALL TABS ^{1,2,3} EVEKEO METHAMPHETAMINE HCL ZENZEDI	Use PA Form# 20420 1. Preferred stimulants will be available without PA if diagnosis of ADHD or Narcolepsy. 2. As per recent FDA alert, Adderall & Dexedrine should not be used in patients with underlying heart defects since they may be at increased risk for sudden death. 3. Dosing limits apply, see Dosing Consolidation List. 4. Max daily dose of 50mg.	
STIMULANT - LONG ACTING AMPHETAMINES SALT	MC/DEL MC MC		AMPHETAMINE/DEXTROAMPHET ER ^{3,4,7} ADDERALL XR CP24 ^{1,3,4,7} VYVANSE ^{2,3,4}	MC MC MC		MYDAYIS ⁵ VYVANSE CHEW ⁴ XELSTRYM ⁸	Use PA Form# 20420 1. As per recent FDA alert, Adderall should not be used in patients with underlying heart defects since they may be at increased risk for sudden death. 2. FDA approval is currently for adults and children 6 or older. Will be available without PA for this age group if within dosing limits. Limit of one capsule daily. Max dose of 70MG daily. 3. Preferred stimulants will be available without PA if diagnosis of ADHD. 4. Dosing limits apply, see Dosing Consolidation List. 5. For the treatment of Attention Deficit Hyperactivity Disorder (ADHD) in patients 13 years and older. 7. FDA approval is currently for adults and children 6 or older. Will be available without PA for this age group if within dosing limits. Max dose of 50MG daily without a PA. 8. For the treatment of patients 6 years of age and older.	DDI: The concomitant use of Mydayis is contraindicated with monoamine oxidase inhibitors (MAOIs) or within 14 days after discontinuing MAOI treatment, as concomitant use can increase hypertensive crisis.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
LONG ACTING AMPHETAMINES	MC MC/DEL MC		DEXTROAMPHET SULF CPSR ^{1,3} DEXTROAMPHETAMINE ER DYANAVEL XR SUS	MC/DEL MC MC MC MC		ADZENYS ER ³ ADZENYS XR- ODT ADZENYS XR ³ DEXEDRINE CAP SR ^{2,3} DYANAVEL XR TAB	Use PA Form# 20420 1. Preferred stimulants will be available without PA if diagnosis of ADHD. 2. As per recent FDA alert, Adderall & Dexedrine should not be used in patients with underlying heart defects since they may be at increased risk for sudden death. 3. Dosing limits apply, see Dosing Consolidation List.	DDI: The concomitant use of Adzenys XR is contraindicated with monoamine oxidase inhibitors (MAOIs) or within 14 days after discontinuing MAOI treatment.
STIMULANT - METHYLPHENIDATE	MC/DEL MC/DEL MC/DEL MC/DEL		DEXMETHYLPHENIDATE IR TABS METHYLPHENIDATE SOL METHYLPHENIDATE TAB METHYLIN TABS ^{1,2}	MC/DEL MC/DEL MC MC MC/DEL MC/DEL		FOCALIN IR TABS METADATE ER METHYLPHENIDATE HCL CHEW METHYLIN CHEWABLES METHYLIN SOL RITALIN	Use PA Form# 20420 1. Preferred stimulants will be available without PA if diagnosis of ADHD. 2. Dosing limits apply, see Dosing Consolidation List. Maximum daily doses are as follows: 72mg daily for Methylphenidate and 36mg daily for Dexmethylphenidate .	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Please refer to General Criteria category E.
STIMULANT - METHYLPHENIDATE - LONG ACTING	MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC MC/DEL		CONCERTA TBCR DEXMETHYLPHENIDATE CAP ER 50/50 FOCALIN XR METHYLPHENIDATE LA CAPS METHYLPHENIDATE ER CAPS 50/50 METHYLPHENIDATE ER CAPS 40/60 METHYLPHENIDATE CD CAPS 30-70 QUILLICHEW ER ^{5,1} QUILLIVANT XR SUS ^{1,5} RITALIN LA ⁴	MC MC/DEL MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL	5 8 8 8 8 8 8 8 8	METADATE CD CPR ADHANSIA XR ^{2,6} APTENSIO XR ² AZSTARYS ⁵ COTEMPLA XR ² COTEMPLA XR ODT ² DAYTRANA ^{2,3} JORNAY PM ^{2,6} METHYLPHENIDATE ER CAPS ^{2,4}	Use PA Form# 20420 1. Preferred stimulants will be available without PA if diagnosis of ADHD. 2. Non-preferred products must be used in specified step order. 3. FDA approval currently only for ages 6-16. Limit of one patch daily. Max dose of 30MG daily. 4. Dosing limits apply, see Dosing Consolidation List. 5. Quillivant XR and Quillichew ER are only indicated for use in patients 6 years of age and older. 6. For the treatment of patients ≥ 6 years of age.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
STIMULANT - STIMULANT LIKE	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC		ATOMOXETINE HCL ARMODAFINIL CLONIDINE ER GUANFACINE ER MODAFINIL TABS QELBREE ^{6,7}	MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC	7 7 8 8 8 8 8 8 8 8 9 9 9	PROVIGIL TABS ³ STRATTERA ^{1, 2} CAFCIT SOLN ³ INTUNIV KAPVAY ONYDA XR ⁶ SUNOSI WAKIX XYREM SOL XYWAV ⁵ NUVIGIL ³ DESOXYN TABS ³ DESOXYN CR ³	Use PA Form# 20710 for Provigil, Nuvigil and Xyrem Use PA Form# 20420 for all others 1. Failure of both an amphetamine and methylphenidate is required for consideration for approval of Strattera , unless history of substance abuse without current use of abusable medication(s). Additionally, for patients <17 years of age, a trial of Guanfacine required before approval of Strattera . 2. Strattera currently has dosing limitations allowing one tablet per day for all strengths if obtain approval. Max daily dose of Strattera is 100mg. Please see Dosing Consolidation List. 3. Non-preferred products must be used in specified step order. 4. Please use generic Guanfacine. 5. For patients 7 years of age and older with narcolepsy. 6. For pediatric patients 6 years of age or older. 7. Preferred with a trial and fail either Atomoxetine OR any 2 preferred ADHD agents.	Provigil requests require diagnosis of Narcolepsy, ADHD, or Obstructive Sleep Apnea. Previous failures of methylphenidate and amphetamine is required for Narcolepsy and ADHD diagnosis, with additional Strattera trial needed with ADHD diagnosis. Please refer to detailed criteria on Provigil PA form. Sunsosi is non-preferred and is indicated for to improve wakefulness in adult patients with excessive daytime sleepiness associated with narcolepsy or obstructive sleep apnea (OSA). Wakix is non-preferred and is indicated for the treatment of excessive daytime sleepiness (EDS) in adults with narcolepsy. Xywav: Diagnosis of cataplexy associated with narcolepsy OR excessive daytime sleepiness associated with narcolepsy. Diagnosis must be confirmed by submission of supporting documentation to include the specialist's interpretation of the Polysomnography (PSG) and Multiple Sleep Latency Test (MSLT) results. FDA reminded healthcare professionals and patients that the combined use of Xyrem (sodium oxalate) with alcohol or central nervous system (CNS) depressant drugs can markedly impair consciousness and may lead to severe breathing problems (respiratory depression). DDI: Sunosi is contraindicated with MAO inhibitors or within 14 days after discontinuing the MAO inhibitor. DDI: Concomitant use of Qelbree with an MAO inhibitor or within 2 weeks after discontinuing an MAO inhibitor is contraindicated. DDI: Concomitant use of Qelbree significantly increases the total exposure, but not peak exposure, of sensitive CYP1A2 substates, which may increase the risk of adverse reactions associated with these CYP1A2 substrates. Coadministration of Qelbree with sensitive CYP1A2 substrates or CYP1A2 substrates with a narrow therapeutic range (e.g. alosetron, duloxetine, ramelteon, tasimelteon, tizanidine, theophylline), is contraindicated.
ANTI-CATAPLECTIC AGENTS								
PSYCHOTHERAPEUTIC AGENTS - MISC.				MC MC		NUDEXTA XENAZINE	Use PA Form# 20710 for Xenazine	
WEIGHT LOSS								
WEIGHT LOSS							No longer covered: Phentermine, Xenical, Didrex, and Meridia	Weight loss drugs are not covered as permitted by Federal Medicaid regulations and Maine Medicaid (MaineCare) Policy.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
ALZHEIMER DISEASE								
ALZHEIMER - Cholinomimetics/Others	MC/DEL		DONEPEZIL HYDROCHLORIDE TABS ¹	MC	6	ARICEPT TABS ²	Use PA Form# 20420 1. PA is required to establish dementia diagnosis and baseline mental status score. 2. Must fail all preferred products before moving to non-preferred. 3. Approvals will require trials and failure or clinical rationale why preferred patches can't be used.	Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Kisunla and Leqembi: Testing to rule out reversible causes of dementia (CBC, CMP, TSH, B12, urine drug screen, RPR/VDRL, (folate if alcohol abuse is present), HIV (if risk present) and an assessment including a review of current medications as a cause of intellectual decline - Prescribed by or in consultation with a neurologist or geriatrician or geriatric psychiatrist. Diagnosis of Alzheimer's disease defined as: • Confirmed presence of amyloid pathology and mild cognitive impairment or mild dementia stage of disease, consistent with Stage 3 and Stage 4 Alzheimer's disease OR • Confirmed presence of amyloid pathology and prodromal or mild dementia stage of disease, consistent with Stage 3 and Stage 4 Alzheimer's disease Testing: • Clinical Dementia Rating (CDR) global score of 0.5 or 1.0 OR • Repeatable Battery for Assessment of Neuropsychological Status (RBANS) delayed memory index score ≤ 85 OR • Mini-Mental State Examination (MMSE) score of 20-30 OR • Montreal Cognitive Assessment (MoCA) score ≤ 22 – Member is age 50 or older – Obtain recent (within one year) brain magnetic resonance imaging (MRI) prior to initiating treatment – Provider attestation to obtain MRIs prior to the 7th infusion (first dose of 10 mg/kg) and 12th infusion (sixth dose of 10 mg/kg) – Member does NOT have history or increased risk of amyloid related imaging abnormalities-edema (ARIA-E), which includes brain edema or sulcal effusions and amyloid related imaging abnormalities hemosiderin deposition (ARIA-H), which includes microhemorrhage and superficial siderosis – Member does NOT have hypersensitivity to any components of these drugs – Failure of or inability to tolerate at least two other preferred Alzheimer therapies for at least four months each, one of which should include a combination of a cholinesterase inhibitor with memantine • If the initial drug utilized is the combination of a cholinesterase inhibitor and memantine, then only that single trial of two drugs is required
	MC/DEL		DONEPEZIL HYDROCHLORIDE ODT ¹	MC	6	ARICEPT ODT ²		
	MC/DEL		EXELON DIS ¹	MC/DEL	7	DONEPEZIL HYDROCHLORIDE TABS 23MG		
	MC/DEL		GALANTAMINE CAPS ¹					
	MC/DEL		GALANTAMINE TAB ¹	MC	8	ADLARITY ³		
	MC/DEL		MEMANTINE ¹	MC/DEL	8	EXELON CAP		
	MC/DEL		RIVASTIGMINE TARTRATE CAPS ¹	MC/DEL	8	GALANTAMINE HYDROBROMIDE SOL		
				MC	8	KISUNLA		
				MC	8	LEQEMBI ^{1,2}		
				MC/DEL	8	MEMANTINE HCL SOL		
				MC/DEL	8	NAMENDA		
				MC/DEL	8	NAMENDA XR CAPS		
				MC/DEL	8	NAMZARIC		
				MC	8	RAZADYNE ²		
				MC	9	COGNEX CAPS ² ZUNVEYL		
SMOKING CESSATION								
NICOTINE PATCHES / TABLETS	MC/DEL MC/DEL MC/DEL MC/DEL		CHANTIX TAB ¹ CHANTIX STARTER PACK NICOTINE DIS PT24 ¹ VARENICLINE TAB	MC/DEL		NICODERM CQ PT24 ¹	Use PA Form# 20420 1. See criteria section for exemptions	As of July 1, 2014 per MaineCare policy, smoking cessation products will be covered without a copay(including MEDEL). No annual or lifetime limits, must follow FDA approved indications and therapy guidelines. Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Note: MaineCare policy, smoking cessation product were "not covered" except for during pregnancy between 9/1/12 and 1/1/14, between 1/1/2014 and 7/1/14 smoking cessation products were covered with limitations. Patients may qualify for the medication through The Maine Tobacco Helpline if they do not have MaineCare or MEDEL. Patients are encouraged to call The Maine Tobacco helpline at 1-800-207-1230.
NICOTINE REPLACEMENT - OTHER	MC/DEL MC/DEL MC/DEL		NICOTINE POLACRILEX GUM ¹ NICOTINE LOZENGE MINI NICOTINE LOZENGE	MC/DEL MC/DEL MC/DEL MC	8 8 8 8	NICOTROL INHALER ^{1,2} NICOTROL NASAL SPRAY ^{1,2} NICORETTE GUM ^{1,2} NICORETTE LOZENGES	Use PA Form# 20420 1. See criteria section for exemptions. 2. Must use non-preferred products in specified step order.	As of July 1, 2014 per MaineCare policy, smoking cessation products will be covered without a copay(including MEDEL). No annual or lifetime limits, must follow FDA approved indications and therapy guidelines. Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Note: MaineCare policy, smoking cessation product were "not covered" except for during pregnancy between 9/1/12 and 1/1/14, between 1/1/2014 and 7/1/14 smoking cessation products were covered with limitations. Patients may qualify for the medication through The Maine Tobacco Helpline if they do not have MaineCare or MEDEL. Patients are encouraged to call The Maine Tobacco helpline at 1-800-207-1230.
ALCOHOL DETERRENTS								
ALCOHOL DETERRENTS	MC/DEL MC MC MC/DEL		ACAMPROSATE ANTABUSE TABS DISULFIRAM TABS NALTREXONE HCL TABS	MC/DEL		ACAMPRO ¹	Use PA Form# 20420 1. Should only be used in conjunction with formal structured outpatient detoxification program.	Preferred generic drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
MISCELLANEOUS ANALGESICS								
ANALGESICS - MISC.	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC		ACETAMINOPHEN ASPIRIN ASPRIN/ APAP/ CAFF TAB BUTAL/ASA/CAFF BUTALBITAL COMPOUND BUTALBITAL/ACET TABS BUTALBITAL/APAP CAPS BUTALBITAL/APAP/CAFFEINE TABS CHOLINE MAGNESIUM TRISALI DIFLUNISAL TABS EXCEDRIN	MC MC/DEL MC/DEL MC MC MC/DEL MC MC MC MC MC		AXOCET CAPS ESGIC-PLUS FIORICET TABS FIORINAL CAPS FIORTAL CAPS FORTABS TABS JOURNAVX ¹ PHRENILIN TABS PHRENILIN FORTE CAPS TRILISATE LIQD TRILISATE TABS	Use PA Form# 20420 1. QL: 1. QL: No greater than 14-day supply within 90 days.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Journavx requires patient must have documented clinical reason as to why they are unable to use acetaminophen and NSAIDS (which can include Cox-II inhibitors). Journavx is FDA approved for moderate to severe ACUTE pain in adults.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
	MC/DEL		SALSALATE TABS	MC MC		ZEBUTAL CAPS ZORPRIN TBCR		
LONG ACTING NARCOTICS								
NARCOTICS - LONG ACTING	MC/DEL		FENTANYL PATCH ⁴	MC	8	ARYMO ER	Use PA Form# 20510	Preferred drugs (Fentanyl Patch, Morphine Sulfate ER tab, and Butrans) must be tried for at least 2 weeks each & failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug & the preferred drug(s) exists. Adequate trials include prevention/treatment of common adverse effects associated w/ narcotics (antinausea, antipruritic, etc.) as well as adequate equianalgesic dosing when converting from one narcotic to another. Also, adequate documentation of attempts to titrate dose of preferred agents to achieve adequate pain relief & desired clinical response must be provided. Member's drug regimen for additions &/or discontinuations of medications that may affect absorption &/or metabolism of preferred agents must be monitored. Approvals will not be granted if patient had access to either non-preferred products or high doses of short acting narcotics during the trial period. Non-preferred drugs will not be approved for patients showing evidence of usage patterns consistent w/ controlled substance abuse such as: 1.Frequent or persistent early refills of controlled drugs; 2.Multiple instances of early refill overrides due to reports of misplacement, stolen, dropped in toilet or sink, distant travel, etc.; 3.Breaches of narcotic contracts with any provider; 4.Failure to comply with patient responsibilities in attached opioid documentation (see PA form) including but not limited to failing to submit to and pass pill counts; 5.Failing to take or pass random drug testing; 6.Failing to provide old records regarding prior use of narcotics; 7.Receiving controlled substances from other prescribers that the provider submitting the PA is unaware of 8.Documented history of substance abuse. Substance abuse evaluations may be required for patients with medical records displaying documented substance abuse or potential signs of narcotic misuse and abuse such as chronic early refills, short dosing intervals, frequent dose increases, multiple lost/stolen etc scripts and intolerance or "allergy" to all products but Oxycontin. 9. Circumventing MaineCare prior authorization requirements for narcotics by paying cash for affected narcotics (prescribers failed to submit prior authorization prior to cash narcotic scripts being filled by member). 10. Requests for any Brand name controlled substance, considered by authorities to be highly abused and diverted (Oxycontin, Percocet, Tylox, Vicodin, Dilaudid, Ultracet...) with an available AB rated generic equivalent will be denied unless it will be provided in a setting that virtually eliminates the risk of diversion. 11. Allergic reactions to any product within a specific narcotic class will justify and preclude use of any other product in the same class due to the risk of cross-hypersensitivity. Hysingla ER - Concomitant use should be avoided with mixed agonist/antagonist analgesics, partial agonist analgesics, and MAOIs. Verify prior trials and failures or intolerance of preferred treatments. Methadone – Established users must have a trial and failure of at least 2preferred drugs for least 2 weeks. Otherwise they will be allowed 180 days to transition to a preferred product.
	MC/DEL		BUTRANS ⁴	MC	8	AVINZA	Use PA Form #10300 for PAs over the opiate limit	
	MC/DEL		MORPHINE SULFATE ER TB12	MC	8	BELBUCA		
				MC	8	EXALGO		
				MC/DEL	8	HYSINGLA ER	1. Oxycontin will be available without PA for patients treated for or dying from cancer or hospice patients. CA (cancer) or HO (hospice) diag code may be used but store must verify since all scripts will be audited and stores will be liable.	
				MC	8	KADIAN		
				MC/DEL	8	METHADONE ⁶	3. Oxycodone ER allowed only 2 per day for all strengths except 80 mg, where 4 are allowed to achieve max total daily dose of 320mg.	
				MC/DEL	8	METHADOSE ⁶		
				MC/DEL	8	MORPHABOND ER		
				MC/DEL	8	MORPHINE SULFATE ER CAP		
				MC/DEL	8	MORPHINE SULFATE SUPP		
				MC/DEL	8	MS CONTIN TB12		
				MC	8	OPANA ER		
				MC/DEL	8	ORAMORPH SR TB12	4. 25mcg, 50mcg, 75mcg, 100mcg. Dosing limits apply, see Dose Consolidation List.	
				MC/DEL	8	OXYCONTIN TB12 ¹		
				MC	8	XARTEMIS ER	5. Non-preferred products must be used in specific order.	
				MC	8	ZOHYDRO ER		
				MC	8	OXYCODONECONC	6. Methadone will be available without PA for patients treated for or dying from cancer or hospice patients or similar conditions as supported by clinical documentation. CA (cancer) or HO (hospice) diag code may be used but store must verify since all scripts will be audited and stores will be liable.	
				MC/DEL	9	OXYCODONE ER ^{3,5}		
NARCOTICS - SELECTED	MC/DEL		TRAMADOL HCL TABS 50 mg ²	MC/DEL	7	RYZOLT	Use PA Form# 20420	Preferred drugs from this and other narcotic classes must be tried for at least 2 weeks each and failed due to lack of efficacy or intolerable side effects before non-preferred drugs from this class will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Approvals will not be granted if patient had access to either non-preferred products or high doses of short acting narcotics during the trial period. Substance abuse evaluations may be required for patients with medical records displaying potential signs of narcotic misuse and abuse such as chronic early refills, short dosing intervals, frequent dose increases, multiple lost/stolen etc scripts and intolerance or "allergy" to all products but desired product. Allergic reactions to any product within a specific narcotic class will justify and preclude use of any other product in the same class due to the risk of cross-hypersensitivity. Non-preferred drugs will not be approved for patients showing evidence of usage patterns consistent with controlled substance abuse such as: 1. frequent or persistent early refills of controlled drugs; 2. multiple instances of early refill overrides due to reports of misplacement, stolen, dropped in toilet or sink, distant travel; 3. breaches of narcotic contracts with any provider; 4. failure to comply with patient responsibilities in attached opioid documentation (see PA form) including but not limited to failing to submit to and pass pill 5. failing to take or pass random drug testing; 6. failing to provide old records regarding prior use of narcotics; 7. receiving controlled substances from other prescribers that the provider submitting the PA is unaware of. Substance abuse evaluations may be required for patients with medical records displaying potential signs of narcotic misuse and abuse such as chronic early refills, short dosing intervals, frequent dose increases, multiple lost/stolen etc scripts and intolerance or "allergy" to all products but Oxycontin. Allergic reactions to any product within a specific narcotic class will justify and preclude use of any other product in the same class due to the risk of cross-hypersensitivity. Beginning January 2017, all current opiate users who are above the maximum combined daily dose of 100 MME must titrate their total daily dose of opioid medications below 30 MME. Also, the maximum daily supply of an opiate prescription for acute pain will be limited to 7-day supplies. The maximum day supply of an opiate prescription for chronic pain will be limited to 30-day supplies. As of July 1, 2017 all users of opioid medications must comply with the maximum combined daily dose of 100 MME. However, for MaineCare members, effective January 1, 2017, opioid prescription(s) for more than a 7-day supply and/or more than 30 MME/ day will require a prior authorization. Please note that MaineCare implemented a 30 MME limit January 1, 2013 that is still effective. Post-surgical members may receive prior authorizations for opiates up to 60 days in length if medical necessity is provided by the surgical provider. An MME conversion chart is available at www.mainearepd.org. Click on "General Pharmacy Info." Please see the Pain Management Policy tab for the complete criteria.
	MC/DEL		TRAMADOL/APAP TABS	MC	8	BUPRENEX SOLN	Use PA form #10300 for PAs over the opiate limit	
				MC/DEL	8	BUTORPHANOL		
				MC	8	NALBUPHINE HCL SOLN	1. Only available if component ingredients are unavailable.	
				MC	8	QDOLO SOLN		
				MC	8	SEGLENTIS ¹	2. Dosing limits apply, please see Dosing Consolidation List.	
				MC	8	STADOL NS SOLN		
				MC	8	TRAMADOL ER		
				MC	8	ULTRACET TABS ¹		
				MC	9	ULTRAM ER		

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
MISCELLANEOUS NARCOTICS								
NARCOTICS - MISC.	MC/DEL		ACETAMINOPHEN/CODEINE	MC/DEL	8	ABSTRAL	Use PA Form# 20420	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Please refer to General Criteria category E.</p> <p>Beginning January 2017, all current opiate users who are above the maximum combined daily dose of 100 MME must titrate their total daily dose of opioid medications below 30 MME. Also, the maximum daily supply of an opiate prescription for acute pain will be limited to 7-day supplies. The maximum day supply of an opiate prescription for chronic pain will be limited to 30-day supplies. As of July 1, 2017 all users of opioid medications must comply with the maximum combined daily dose of 100 MME.</p> <p>However, for MaineCare members, effective January 1, 2017, opioid prescription(s) for more than a 7-day supply and/or more than 30 MME/ day will require a prior authorization. Please note that MaineCare implemented a 30 MME limit January 1, 2013 that is still effective.</p> <p>Post-surgical members may receive prior authorizations for opiates up to a 60 days in length if medical necessity is provided by the surgical provider.</p> <p>An MME conversion chart is available at www.mainearepdl.org. Click on "General Pharmacy Info."</p> <p>Please see the Pain Management Policy for the complete criteria</p>
	MC/DEL		ASPIRIN/CODEINE TABS	MC/DEL	8	APADAZ	Use PA form #10300 for PAs over the opiate limit	
	MC/DEL		BUTAL/ASA/CAFF/COD CAPS	MC/DEL	8	ASCOMP/CODEINE CAPS		
	MC		BUTALBITAL/ASPIRIN/CAFFEI CAPS	MC/DEL	8	BUTALBITAL/APAP/CAFFEINE/ CAPS	1. Fentanyl OT loz (Barr) and Capital and Codeine Suspension products require PA for users over 18 years of age. PA is not required if under 18 years of age.	
	MC		CAPITAL AND CODEINE SUSP ¹	MC/DEL	8	BUTALBITAL COMPOUND- CODEINE CAP	2. Oxycodone/Acet 10/650 is 8 times more expensive. Use twice as many of Oxycodone/Acet 5/325 instead. You can mix and match preferred strengths of Oxycodone and Oxycodone/Acet to minimize Acet dose similar to certain non-preferred drugs.	
	MC/DEL		CODEINE PHOSPHATE SOLN	MC	8	DEMEROL	3. Only preferred manufacturer's products will be available without prior authorization.	
	MC/DEL		CODEINE SULFATE TABS	MC/DEL	8	DILAUDID		
	MC/DEL		ENDOCET TABS ³	MC	8	DILAUDID-HP SOLN		
	MC/DEL		ENDODAN TABS	MC	8	FENTANYL CITRATE SOLN		
	MC/DEL		FENTANYL OT LOZ ¹	MC/DEL	8	FENTORA		
	MC/DEL		HYDROCODONE/ACETAMINOPHEN	MC/DEL	8	FIORICET/CODEINE CAPS		
	MC/DEL		HYDROMORPHONE HCL ³	MC	8	FIORINAL/CODEINE #3 CAPS		
	MC		LORTAB ELX	MC	8	FIORTAL/CODEINE CAPS		
	MC/DEL		MEPERIDINE SOL	MC/DEL	8	HYDROCODONE/IBUPROFEN		
	MC/DEL		OXYCODONE TAB	MC/DEL	8	HYDROMORPHONE ER		
	MC/DEL		OXYCODONE/ACETAMINOPHEN ^{2,3}	MC/DEL	8	HYDROMORPHONE RECTAL SUPP		
	MC/DEL		ROXICET	MC	8	IBUDONE		
	MC		ROXIPRIN TABS	MC/DEL	8	LEVORPHANOL TARTRATE TAB		
				MC/DEL	8	LORCET		
				MC	8	LORTAB		
				MC	8	MAXIDONE TABS		
				MC/DEL	8	MEPERIDINE TABS		
				MC/DEL	8	NORCO TABS		
				MC/DEL	8	ONSOLIS		
				MC/DEL	8	OXECTA		
				MC/DEL	8	OXYCODONE CAP		
				MC/DEL	8	OXYCODONE/APAP 10/650		
				MC/DEL	8	OXYCODONE/APAP 7.5/500		
				MC/DEL	8	PENTAZOCINE/ACET TABS		
				MC/DEL	8	PENTAZOCINE/NALOXONE TABS		
				MC	8	PERCOCET TABS		
				MC	8	PHRENILIN W/CAFFEINE/CODE CAPS		
				MC/DEL	8	ROXICET 5/500 TABS		
				MC	8	ROXICODONE TABS		
				MC/DEL	8	ROXYBOND		
				MC	8	SYNALGOS-DC CAPS		
				MC	8	TALACEN TABS		
				MC	8	TREZIX		
				MC	8	TYLENOL/CODEINE #3 TABS		
				MC	8	TYLOX CAPS		
				MC	8	XOLOX		
				MC	8	VICODIN		
				MC	8	VICOPROFEN TABS		
				MC	8	ZYDONE TABS		
				MC	9	ACTIQ LPOP		
				MC	9	CONZIP		
				MC	9	OPANA		
OPIOID DEPENDENCE TREATMENTS	MC		SUBOXONE FILM ²	MC/DEL		BUPRENORPHINE ¹	Use PA form #20200 for Extended Release Buprenorphine	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p> <p>Members will continue to be required to follow the criteria listed below:</p> <p>1-Induction period for 30 days</p> <p>2-Max dose of 32 mg for induction</p> <p>3-Max dose of 24 mg for maintenance</p> <p>4-There is not more than one opioid fill in member's drug profile between current fill of Buprenorphine and a prior Buprenorphine fill within the past 90 days</p> <p>5- Should provide evidence of monthly monitoring including random pill counts, urine drug tests and use of Maine Prescription Monitoring Program reports.</p> <p>6- Buprenorphine monotherapy is preferred if member is pregnant and dose not > 24 mg day and pregnancy diagnosis is noted on the prescription.</p>
	MC/DEL		BUPRENORPHINE/NALOXONE TABS ²	MC		ZUBSOLV	Use PA Form #20100 for all others 1. Buprenorphine will only be approved for use during pregnancy. 2. See Criteria Section.	

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
EXTENDED RELEASE BUPRENORPHINE	MC MC		BRIXADI ¹ SUBLOCADE ¹				Use PA form #20200 for Extended Release Buprenorphine 1. Clinical PA required.	Brixadi and Sublocade: The prescriber can attest (and medical record should document) that: -member has a documented history of opioid use disorder (OUD), -XRB is being used for the treatment of OUD (rather than pain or any other non-FDA approved indication) and -member's total daily dose of sublingual buprenorphine is less than or equal to 24 mg daily. AND at least one of the following is true: -The member's previous use of sublingual buprenorphine has included misuse, overuse, or diversion. -The member is at high risk of overdose (e.g., individuals leaving incarceration or abstinence-based treatment programs; individuals who are unhoused; or those facing potential gaps in care due to delays in care or geographically limited treatment access). -The member has experienced significant medical complications of OUD and/or of injection drug use. Occurrence should be in the last 5 years, or it should be clearly documented that the risk indicated by this infection or complication is ongoing (Examples of medical complications of OUD include: threatened the function of organs or life or limb threatening and required medical and/or surgical therapy. Examples of medical complications of injection drug use include osteomyelitis, endocarditis, renal failure, joint infection or other serious medical complications directly related to OUD.) -The member has treatment-resistant OUD, including those with ongoing illicit substance use in the context of sublingual buprenorphine treatment as documented by positive urine drug screens or other clear objective evidence, and/or further functional decline with explicit documentation of the functional decline. -The member has a significant intolerance of, or documented allergy to, sublingual buprenorphine (either buprenorphine monotherapy or buprenorphine/naloxone combination therapy) that has resulted in the patient's inability to comply with continued treatment using the sublingual product. (A true allergy is usually accompanied by rash, respiratory symptoms, or anaphylaxis. Other complaints such as bad taste, mouth tingling, etc. do not constitute evidence of allergy or significant intolerance. Formulation preference or convenience are not, in and of themselves, indications for using XRB.) -The member is in ongoing treatment with XRB and would like to continue the medication.
OPIOID WITHDRAWAL AGENTS				MC		LUCEMYRA ¹	Use PA Form#20420 1. Clinical PA for appropriate approved use and patient has documented contraindication to Clonidine .	
NARCOTIC ANTAGONISTS								
NARCOTIC - ANTAGONISTS	MC/DEL MC MC MC MC MC		NALTREXONE HCL TABS NALOXONE INJ NARCAN NS NALOXONE SPRAY OTC VIVITROL INJ ZIMHI	MC MC MC MC/DEL		EVZIO OPVEE ² KLOXXADO REVIA TABS ¹	Use PA Form# 20420 1. Will only be approved for side effects experienced with generic that are not described in the literature as occurring with the brand version. 2. For the treatment of adult and pediatric patients 12 years of age and older.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
COX 2 / NSAIDS								
COX 2 INHIBITORS - SELECTIVE / HIGHLY SELECTIVE	MC/DEL MC/DEL MC/DEL MC/DEL		CELECOXIB ^{4,5} KETOROLAC TROMETHAMINE ^{2,3,5} NABUMETONE TABS ⁵ MELOXICAM TABS ^{1,5}	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC		CELEBREX CAPS ^{4,5} MELOXICAM CAPS ⁵ MOBIC ⁵ MOBIC SUSP ⁵ RELAFEN TABS ⁵ QMIIZ ODT VIVLODEX XIFYRM ⁵	Use PA Form# 20420 1. Meloxicam and Xifyrm have dosing limits allowing one tablet daily of all strengths without PA. 2. Ketorolac Tromethamine is indicated for the short term (up to 5 days) management of moderately severe acute pain that requires analgesic at the opioid level in adults. Not indicated for minor or chronic pain conditions. 3. Ketorolac has dosing limits allowing 24 tablets for a 5 day supply every 30 days. 4. Dosing limits will be set at a maximum of 400mg daily. 5. The FDA has issued a Public Health Advisory warning of the potential for increased cardiovascular risk & GI bleeding with NSAID use.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
NSAIDS	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		CHILDRENS IBUPROFEN DICLOFENAC POTASSIUM TABS DICLOFENAC SODIUM TABS DICLOFENAC SODIUM 1% GEL ¹ ETODOLAC	MC MC MC MC MC/DEL		ADVIL TABS ANAPROX TABS ANAPROX DS TABS CAMBIA CATAFLAM TABS	Use PA Form# 20420 The FDA has issued a Public Health Advisory warning of the potential for increased cardiovascular risk & GI bleeding with NSAID use.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Approvals will be granted for other requests based on failure of at least one generic NSAID from at least 3 different NSAID classes as described in the COX-II PA form.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
				MC MC MC/DEL MC MC MC MC MC MC MC		REDITREX REMICADE RENFLEXIS SIMLANDI TOFIDENCE VELSIPITY YUFLYMA YUSIMRY XATMEP ⁵ ZYMFENTRA		
ALOPECIA AREATA AGENTS								
ALOPECIA AREATA AGENTS				MC MC/DEL MC	7 8 8	OLUMIANT LITFULO LEQSELVI ¹	Use PA Form# 20420 1. Clinical PA is required to establish diagnosis and medical necessity.	Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
MISCELLANEOUS ARTHRITIS								
ARTHRITIS - MISC.	MC MC		RIDAURA CAPS MYOCHRYSLINE SOLN	MC/DEL		ARTHROTEC ¹	Use PA Form# 20420 1. The individual components of Arthrotec are available without PA.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. The individual components of Arthrotec are available without PA.
LUPUS-SLE								
LUPUS-SLE				MC MC MC		BENLYSTA ¹ LUPKYNIS SAPHNELO	Use PA Form# 20420 1. Approvals will require previous trial of corticosteroids, antimalarials, NSAIDS and immunosuppressives.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Lupkynis is a sensitive CYP3A4 substrate. Co-administration with strong or moderate CYP3A4 inhibitors increases voclosporin exposure, which may increase the risk of Lupkynis adverse reactions. Co-administration of Lupkynis with strong CYP3A4 inhibitors (e.g. ketoconazole, itraconazole, clarithromycin) is contraindicated. Reduce Lupkynis dosage when co-administered with moderate CYP3A4 inhibitors (e.g. verapamil, fluconazole, diltiazem).
PIK3CA-Related Overgrowth Spectrum (PROS)								
PIK3CA-Related Overgrowth Spectrum (PROS)				MC		VIJOICE ¹	Use PA Form# 20420 1. PA required to confirm FDA approved indication.	Preferred drugs must be tried and failed, in step-order, due to lack of efficacy (failure to reach target IOP reduction) or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
MIGRAINE THERAPIES								
MIGRAINE - ERGOTAMINE DERIVATIVES				MC/DEL MC		D.H.E. 45 SOLN TRUDHESA	Use PA Form# 10110	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
MIGRAINE - CARBOXYLIC ACID DERIVATIVES	MC		DIVALPROEX ER TB24	MC		DEPAKOTE ER TB24	Use PA Form# 10110	
MIGRAINE - SELECTIVE SEROTONIN AGONISTS (5HT)-- Tabs/Nasal	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	1 1 1 1 1 1 2	MIGRANAL NASAL SPRAY RELPAX ¹ RIZATRIPTAN ODT RIZATRIPTAN TABS SUMATRIPTAN TABS ¹ ZOLMITRIPTAN TAB ¹ NARATRIPTAN HCI TABS ¹	MC MC MC/DEL MC MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		AMERGE TABS ^{1,2} AXERT TABS ^{1,2} FROVA TABS ^{1,2} IMITREX NASAL SPRAY ¹ IMITREX TABS ^{1,2} MAXALT ^{1,2,3} MAXALT MLT ^{1,2,3} ONZETRA XSAIL ² SUMATRIPTAN NASAL SPRAY ¹ ZOLMITRIPTAN ODT ZOLMITRIPTAN SPRAY ZOMIG TABS ^{1,2} ZOMIG NASAL SPARY ^{1,2} ZOMIG ZMT TBDP ^{1,2}	Use PA Form# 10110 1. All drugs in this category have dosing limits. Refer to Dose Consolidation Table. 2. Must fail all preferred products before non-preferred. 3. Established users will be grandfathered.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Quantity limit exceptions will require ongoing therapy with therapeutic doses of highly effective prophylactic medication as listed on the Triptan PA form.
MIGRAINE - SELECTIVE SEROTONIN AGONISTS (5HT)-- Injectables	MC MC/DEL MC/DEL		IMITREX CARTRIDGE ¹ SUMATRIPTAN SYRINGE ¹ SUMATRIPTAN PEN INJCTR ¹	MC/DEL MC MC		TOSYMRA ZEMBRACE ¹ IMITREX PEN INJCTR ¹	Use PA Form# 10110 1. Dosing limits apply, see Dosage Consolidation List.	
MIGRAINE - SELECTIVE SEROTONIN AGONISTS (5HT)-- Combinations				MC/DEL		TREXIMET ^{1,2}	Use PA Form# 10110 1. Dosing limits apply, see Dosage Consolidation List.	

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
							2. Use preferred Sumatriptan and Naproxen separately. Treximet only available if component ingredients of Sumatriptan and Naproxen are unavailable.	
MIGRAINE - SELECTIVE SEROTONIN AGONISTS--Combinations				MC	8	SYMBRAVO ¹	1. Dosing limits apply, see Dosage Consolidation List.	
MIGRAINE - PREVENTATIVE TREATMENT	MC MC/DEL MC/DEL MC/DEL MC/DEL		AIMOVIG ¹ AJOVY ¹ AJOVY AUTO INJECT ¹ EMGALITY SYRINGE ¹ 120mg/ml EMGALITY PEN ¹ 120mg/ml	MC MC MC		NURTEC ODT ² QULIPTA VYEPTI ²	Use PA Form# 10110 1. See criteria section. 2. Dosing limits apply, see Dosage Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Aimovig, Ajovy and Emgality: The patient is 18 years of age or older AND patient has a diagnosis of episodic migraine (4-14 headache days per month with migraine lasting 4 hours or more) or chronic migraine (≥ 15 headache days per month, of which ≥ 8 are migraine days, for at least 3 months) AND patient has failed or has a contraindication to an adequate trial (≥ 60 days) of at least 2 medications for migraine prophylaxis from at least 2 different classes. Nurtec ODT will be preferred after 2 adequate trials of at least two preferred triptans.
MIGRAINE - ACUTE TREATMENT	MC MC/DEL		NURTEC ODT ¹ SPASTRIN TABS	MC MC MC/DEL MC/DEL MC MC MC/DEL		BELCOMP-PB SUPP ELYXYB MIGRAZONE CAPS MIGERGOT SUP REYVOW UBRELVY ZAVZPRET	Use PA Form# 10110 1. Dosing limits apply, see Dosage Consolidation List.	Nurtec ODT will be preferred after 2 adequate trials of at least two preferred triptans. Reyvow is non-preferred and is indicated for the acute treatment of migraine with or without aura in adults. Reyvow is not indicated for the preventive treatment of migraine. Ubrelyvy is non-preferred and is indicated for the acute treatment of migraine with or without aura in adults. This is not indicated for the preventive treatment of migraine. Zavzpret: The patient must have a documented side effect, allergy, or treatment failure to preferred oral CGRP Inhibitor and two non-preferred oral CGRP Inhibitors.
GOUT								
GOUT	MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL		ALLOPURINOL TABS COLCHICINE TAB FEBUXOSTAT TAB MITIGARE PROBENECID TABS PROBENECID/COLCHICINE TABS	MC/DEL MC MC MC/DEL MC		COLCHICINE CAP COLCRYS GLOPERBA ULORIC ¹ ZYLOPRIM TABS	Use PA Form# 20420 1. Failure of therapeutic (300mg) dose of Allopurinol (failure define as not being able to get uric acid levels below 6mg/dl) or severe renal disease.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: The concomitant use of Gloperba and CYP3A4 inhibitors (e.g. clarithromycin, ketoconazole, grapefruit juice, erythromycin, verapamil, etc.) should be avoided due to the potential for serious and life-threatening toxicity.
MISC.								
ACID SPHINGOMYELINASE DEFICIENCY (ASMD)				MC		XENPOZYME ^{1,2}	Use PA Form# 20420 1. For treatment of non-central nervous system manifestations of acid sphingomyelinase deficiency (ASMD) in adult and pediatric patients. 2. Clinical PA required for appropriate diagnosis and clinical parameters.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANESTHETICS - MISC.	MC MC MC		BUPIVACAINE HCL SOLN LIDOCAINE HCL SOLN MARCAINE SOLN	MC MC/DEL MC		SENSORCAINE-MPF SOLN SYNVISC INJ XYLOCAINE SOLN	Use PA Form# 30130	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
COLD AGGLUTININ DISEASE (CAD)				MC		ENJAYMO ¹	Use PA Form# 20420 1. Indicated to decrease the need for red blood cell transfusion due to hemolysis in adults with cold agglutinin disease (CAD).	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
CONGENITAL ADRENAL HYPERPLASIA				MC		CRENESSITY	Use PA Form# 30130	Crenessity - As adjunctive treatment to glucocorticoid replacement to control androgens in adults and pediatric patients 4 years of age and older with classic congenital adrenal hyperplasia (CAH).
PRIMARY HYPEROXALURIA TYPE 1 (PH1)						OXLUMO ¹ RIVFLOZA	Use PA Form# 20420 1. PA is required to establish diagnosis and medical necessity.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Rivfloza: The patient has a diagnosis of Primary Hyperoxaluria Type 1 (PH1) confirmed via genetic testing (identification of alanine: glyoxylate aminotransferase gene (AGXT) mutation) AND urinary oxalate excretion > 0.5mmol/1.73 m2 or urinary oxalate: creatinine ratio is above the upper limit of normal for age AND is at least 9 years of age AND medication is being prescribed by, or in consultation, with a nephrologist or urologist.
SICKLE CELL DISEASE	MC MC MC/DEL MC		DROXIA CASGEVY ^{2,3} HYDROXYUREA LYFGENIA ^{2,3}	MC MC MC/DEL		ADAKVEO ENDARI ¹ SIKLOS XROMI	Use PA Form# 20420 1. Evidence of other preferred L-glutamine products utilization and reason for failure. 2. For the treatment of patients ≥ 12 years of age. 3. PA required to confirm FDA approved indication.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

[illegible]

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria			
	MC/DEL		PRIMIDONE TABS	MC	8	ROWEEPRA TAB					
	MC/DEL		QUDEXY XR	MC	8	SABRIL					
	MC/DEL		TEGRETOL SUS	MC	8	SEZABY					
	MC/DEL		TOPIRAMATE	MC	8	SPRITAM					
	MC/DEL		TOPIRAMATE SPRINKLE IR CAPS	MC	8	SYMPAZAN					
	MC/DEL		TRILEPTAL SUS	MC/DEL	8	TEGRETOL TAB					
	MC/DEL		VALPROIC ACID TABS	MC/DEL	8	TIAGABINE					
	MC/DEL		VALPROIC ACID SOL	MC	8	TOPAMAX					
	MC		VALTOCO ²	MC/DEL	8	TOPIRAMATE ER CAPS					
	MC/DEL		ZONISAMIDE	MC	8	TOPAMAX SPRINKLE ER CAPS ²					
				MC	8	TOPAMAX SPRINKLE IR CAPS ²					
				MC/DEL	8	TOPIRAMATE SPRINKLE ER CAPS ²					
				MC	8	TROKENDI ^{2,6}					
				MC	8	VIGAFYDE					
				MC/DEL	8	VIMPAT ⁴					
				MC/DEL	8	VIMPAT SOL ⁴					
				MC	8	XCOPRI					
				MC/DEL	8	ZARONTIN SYRP					
				MC/DEL	8	ZARONTIN CAP					
				MC/DEL	8	ZARONTIN SOL					
				MC	8	ZONISADE					
				MC	8	ZTALMY					
				MC/DEL	9	KEPPRA XR					
				MC/DEL	9	NEURONTIN					
				MC/DEL	9	TEGRETOL-XR TB12					
										<u>BIPOLAR DISORDER: STEP ORDER</u>	SEE ANTICONVULSANT INDICATION CHART AT THE END OF THIS DOCUMENT
									M ~ A		
									4 ~ 4	LAMICTAL	
									4 ~ 4	LITHIUM	M= Monotherapy
									4 ~ 4	CARBAMAZEPINE	A= Adjunctive
									4 ~ 4	VALPROATE	9= No Evidence
									4 ~ 4	ATYPICAL ANTIPSYCHOTICS EXC. CLOZAPINE	The step orders show the relative strength of evidence for use in bi-polar and will guide prior authorization determinations.
									5 ~ 5	TRILEPTAL	Step 4 drugs-no PA required.
									9 ~ 6	TOPAMAX	
									9 ~ 7	KEPPRA TABS	
									9 ~ 8	GABITRIL TABS	
									9 ~ 9	NEURONTIN	
										<u>PEDIATRIC BIPOLAR1 DISORDER: STEP ORDER</u>	
									M ~ A	(6-18 YEARS WITH OR WITHOUT PSYCHOSIS)	Two-step 1 preferred drugs must be tried before Trileptal .
									4 ~ 4	LITHIUM	The step orders show the relative strength of evidence for use in bi-polar and will guide prior authorization determinations.
									4 ~ 4	CARBAMAZEPINE	Step 4 drugs-no PA required.
									4 ~ 4	VALPROATE	
									4 ~ 4	ATYPICAL ANTIPSYCHOTICS EXC.CLOZAPINE	
									4 ~ 4	LAMICTAL	
					5 ~ 5	TRILEPTA					
ANTI-PARKINSON DRUGS											
PARKINSONS - ANTICHOLINERGICS	MC/DEL MC MC/DEL		BENZTROPINE MESYLATE TABS COGENTIN SOLN TRIHXYPHENIDYL				Use PA Form# 20420				
PARKINSONS - ADENOSINE RECEPTOR ANTAGONIST				MC/DEL		NOURIANZ	Use PA Form# 20420	Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Avoid use of Nourianz with strong CYP3A4 inducers (e.g. carbamazepine, rifampin, phenytoin, St. John's wort).			

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
PARKINSONS - COMT INHIBITORS				MC/DEL MC MC/DEL		COMTAN TABS ONGENTYS TASMAR TABS	Use PA Form# 20420	Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
PARKINSONS - SELECTED DOPAMIN AGONISTS	MC/DEL MC/DEL		PRAMIPEXOLE ROPINIROLE NEUPRO PATCH	MC/DEL MC MC/DEL	5 8 8	MIRAPEX TABS ¹ REQUIP TABS MIRAPEX ER	Use PA Form# 20420 1. As of 12/08 users of Mirapex will be grandfathered if diagnosis is Parkinson's.	Preferred drug must be tried and failed in step-order due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
PARKINSONS- MAOIS				MC		XADAGO	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
PARKINSONS - DOPAMINERGICS/ CARBII/ LEVO	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL		AMANTADINE HCL CAPS AMANTADINE HCL TABS BROMOCRIPTINE MESYLATE TABS BROMOCRIPTINE MESYLATE CAPS CARBIDOPA/LEVODOPA TABS ³ CARBIDOPA/LEVODOPA ER CARBIDOPA/LEVO/ENTACAPONE TAB LARODOPA TABS SELEGILINE CAPS HCL SELEGILINE TABS HCL	MC/DEL MC MC/DEL MC MC MC MC/DEL MC MC MC MC MC MC MC MC MC MC		APOKYN AZILECT ² CARBIDOPA/LEVODOPA RAPDIS CREXONT ⁴ ELDEPRYL CAPS GOCOVRI INBRIJA KYNMOBI LODOSYN TABS ONAPGO OSMOLEX ER PARLODEL CAPS PARLODEL TABS RYTARY SINEMET TABS SINEMET TBCR VYALEV ZELAPAR ¹	Use PA Form# 20420 1. Approvals will require concurrent therapy with Levodopa and failed trials of Selegiline, Comtan, and Stalevo . 2. Approvals will require trials of Carbidopa/-Levodopa, Selegiline, Comtan, and Stalevo. 3. Only preferred manufacturer's products will be available without prior authorization. 4. Approvals will require trials of preferred medications including extended-release levodopa/carbidopa tablets.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Inbrija is recommended for the intermittent treatment of OFF episodes in patients with Parkinson's disease treated with carbidopa/levodopa.
PARKINSONS - COMBO.				MC/DEL MC		STALEVO ¹ CARBIDOPA/LEVODOPA/ENTACA ¹	Use PA Form# 20420 1. Clinical PA is required to establish diagnosis and medical necessity.	
MUSCLE RELAXANTS								
MUSCLE RELAXANTS	MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL		BACLOFEN TABS CHLORZOXAZONE TABS CYCLOBENZAPRINE HCL 5mg & 10mg TABS LIORESAL INTRATHECAL KIT METHOCARBAMOL TABS TIZANIDINE HCL TABS	MC/DEL MC/DEL MC/DEL MC/DEL MC MC MC MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC	7 8 8 8 8 8 8 8 8 8 8 8 9 9 9 9 9	ORPHENADRINE CITRATE CARISOPRODOL 350MG TABS AMRIX DANTRIUM CAPS FLEQSUVY LIORESAL TABS LORZONE LYVISPAH METAXALONE NORFLEX TBCR OZOBAX ROBAXIN-750 TABS VECUROMIUM INJ ZANAFLEX TABS CARISOPRODOL 250MG TABS CHLORZOXAZONE 250mg TABS SKELAXIN TAB SOMA TABS TANLOR	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. At least 4 preferred drugs (including tizanidine) must be tried for at least 2 weeks and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Elderly patients, over 65, will require written notice of the increased sedative risks and impaired driving. Prior Authorization will not be given for: 1. frequent or persistent early refills of controlled drugs; 2. multiple instances of early refill overrides due to reports of misplacement, stolen, dropped in toilet or sink, distant travel, etc. Non-preferred products must be used in specified step order. Non-preferred drugs will not be approved if members circumventing MaineCare prior authorization requirements by paying (prescribers failed to submit prior authorization prior to cash narcotic scripts being filled by member). Lorzone is non preferred and requires at least 4 preferred drugs (including tizanidine) and step care therapy (orphenadrine), as well as reasons for why chlorzoxazone is not acceptable.
MUSCLE RELAXANT - COMBO.				MC/DEL MC/DEL MC MC/DEL MC/DEL MC		CARISOPRODOL/ASPIRIN TABS CARISOPRODOL/ASPIRIN/CODE NORGESIC TABS ORPHENADRINE COMPOUND ORPHENADRINE/ASA/CAFF ORPHENGESIC	Use PA Form# 20420	Individual components are available with PA described in the section above. 1. frequent or persistent early refills of non-controlled drugs; 2. multiple instances of early refill overrides due to reports of misplacement stolen, dropped in toilet or sink, distant travel, etc.

[illegible]

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
	MC MC/DEL MC MC MC/DEL MC/DEL MC MC MC MC MC MC MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC MC/DEL MC MC MC/DEL MC MC MC		FERATAB TABS FER-GEN-SOL SOLN FER-IRON SOLN FERRONATE TABS FERROUS SULFATE FLUOR-A-DAY CHEW FLUORIDE CHEW FLUORIDE SODIUM CHEW FLUORITAB CHEW HM CALCIUM TABS K+ POTASSIUM PACK KAON ELIX KAON-CL-10 TBCR KCL 0.075% / D5W / NACL 0.2% SOLN K-EFFERVESCENT TBEF KLOR-CON KLOTRIX TBCR K-PHOS TABS K-VESCENT TBEF LURIDE CHEW MAGNESIUM GLUCONATE TABS MAGNESIUM SULFATE SOLN MAGTABS MICRO-K 8 MEG OS-CAL TABS OS-CAL 500 + D TABS OYSCO OYST-CAL TABS OYST-CAL D TABS OYST-CAL/VITAMIN D TABS OYSTER CALCIUM TABS OYSTER SHELL PHARMA FLUR PHOSPHA 250 NEUTRAL TABS POTASSIUM BICARBONATE TBEF POTASSIUM CHLORIDE 8MEQ POTASSIUM EFFERVESCENT SELENIUM TABS SLOW-MAG TBCR SODIUM FLUORIDE V-R CALCIUM V-R OYSTER SHELL CALCIUM ZINC SULFATE CAPS	MC MC/DEL MC MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC MC MC/DEL MC MC MC		K-LYTE K-PHOS TABS NEUTRAL K-TABS TBCR K-VESCENT PACK MICRO-K 10 MEG CPCR NU-IRON 150 CAPS OYSTER SHELL CALCIUM/VITA TABS POLY-IRON 150 CAPS POLYSACCHARIDE IRON CAPS POTASSIUM BICARB/CHLORIDE POTASSIUM CHLORIDE 10MEQ CAPS POTASSIUM CHLORIDE 8MEQ CAPS TUMS 500 CHEW VIACTIV CHEW		
PHENYLKETONURIA (PKU) TREATMENT AGENTS								
PHENYLKETONURIA (PKU) TREATMENT AGENTS- INJECTABLES				MC		PALYNZIQ ¹	Use PA Form# 20420 1. For the treatment of patients ≥ 18 years of age.	Palynziq is not to be used in combination with kuvan.
PHENYLKETONURIA (PKU) TREATMENT AGENTS- ORAL				MC		KUVAN JAVYGTOR (ORAL) TABLET SOL 100 MG JAVYGTOR (ORAL) POWD PACK 100 MG JAVYGTOR (ORAL) POWD PACK 500 MG SAPROPTERIN DIHYDROCHLORIDE (ORAL) TABLET SOL 100 MG SAPROPTERIN DIHYDROCHLORIDE (ORAL) POWD PACK 100 MG SAPROPTERIN DIHYDROCHLORIDE (ORAL) POWD PACK 500 MG CYSTADANE (ORAL) POWDER 1G/SCOOP	Use PA Form# 20420	

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
MISC. ELECTROLYTES/NUTRITIONALS								
ELECTROLYTES/ NUTRITIONALS	MC		INTRALIPID EMUL ¹	MC		BOOST ¹	Use PA Form# 20420 & SGA Form 1. This list of nutritionals is incomplete. All nutritionals still require a PA except for the miscellaneous products listed as preferred. SGA form required for nutritionals unless member has a G/I tube. 2. Formerly known as Omacor .	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval. Medical foods are not to be authorized solely for the purpose of enhancing nutrient intake or managing body weight if the participant is able to eat conventional foods adequately. Medical foods may be approved if the member has a medical condition which precludes or restricts the use of conventional foods and necessitates the use of a formula. Concurrent Stimulant therapy is not an acceptable medical reason/condition for use of medical foods for enhancing nutrient intake or managing body weight. For children under the age of 5, MaineCare will not provide milk- or soy-based standard infant formulas. Regular formulas may be sought through your nearest WIC office. MaineCare will continue to cover medical food for all participants in MaineCare when medical necessity is met. Vascepa requires adjunct therapy for specific indication to reduce TG in those with severe hypertriglyceridemia (500mg per deciliter or more). Proper indication per lab values is required before approval.
	MC		P.T.E. -5 SOLN ¹	MC		CASEC POWD ¹		
	MC		SEA-OMEGA CAPS ¹	MC		CHOICE DM LIQD ¹		
				MC		DELIVER 2.0 LIQD ¹		
				MC		DOJOLVI		
				MC		ENFAMIL ¹		
				MC		ENSURE ¹		
				MC		GLUCERNA ¹		
				MC		ISOCAL LIQD ¹		
				MC		KINDERCAL TF LIQD ¹		
				MC		KINDERCAL TF/FIBER LIQD ¹		
				MC		L-CARNITINE CAPS ¹		
				MC		LIPISORB LIQD ¹		
				MC		LOVAZA ^{1,2}		
				MC		MODULEN IBD POWD ¹		
				MC		NUTRAMIGEN POWD ¹		
				MC		NUTREN ¹		
				MC		NUTRITIONAL SUPPLEMENT LIQD ¹		
				MC		NUTRIVENT 1.5 LIQD ¹		
				MC		PEPTAMEN ¹		
				MC		PHENYLADE ¹		
				MC		PHENYL-FREE ¹		
				MC		PKU 3 POWD ¹		
				MC		PREGESTIMIL POWD ¹		
				MC		PROBALANCE LIQD ¹		
				MC		PROSOBEE ¹		
				MC		SCANDISHAKE PACK ¹		
				MC		VASCEPA		
ERYTHROPOEITINS	MC		EPOGEN SOLN	MC	8	ARANESP SOLN ¹	Use PA Form# 10520 1. Clinical PA is required to establish medical necessity and that appropriate lab monitoring is being done.	Non-Preferred drugs must be tried and failed in step-order, due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Please see the EPO PA form for other approval and renewal criteria.
	MC		MIRCERA SYRINGE	MC	8	PROCRIT SOLN ¹		
	MC		RETACRIT					
GRANULOCYTE CSF								
GRANULOCYTE CSF	MC		FULPHILA	MC	8	FYLNETRA	Use PA Form# 20520 1. Must be used in specified step order.	See approval criteria detailed on Granulocyte Colony Stimulating Factor PA form.
	MC		NEUPOGEN SYRINGE	MC	8	GRANIX SYRINGE		
	MC		NEUPOGEN VIAL	MC	8	GRANIX VIAL		
	MC/DEL		NYVEPRIA SYRINGE	MC	8	LEUKINE		
				MC/DEL	8	NIVESTYM		
				MC	8	ROLVEDON		
				MC	8	RYZNEUTA		
				MC	8	STIMUFEND		
				MC/DEL	8	ZARXIO		
				MC/DEL	8	ZIEXTENZO		
				MC	9	NEULASTA ¹		
GAUCHER DISEASE								
GAUCHER DISEASE				MC		CERDELGA ¹	Use PA Form# 20420 1. Clinical PA for indication required.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Exceeding days supply limits for LMWH class requires PA. Yargesa : As monotherapy for the treatment of adult patients with mild to moderate type 1 Gaucher disease for whom enzyme replacement therapy is not a therapeutic option (e.g., due to allergy, hypersensitivity, or poor venous access).
				MC		YARGESA ¹		
NIEMANN-PICK DISEASE AGENTS								
NIEMANN-PICK DISEASE AGENTS				MC		AQNEURSA ¹	Use PA Form# 20420 1. Clinical PA required for appropriate diagnosis.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
				MC		MIPLYFFA ¹		

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
ANTICOAGULANTS / PLATELET AGENTS								
ANTICOAGULANTS	MC MC/DEL MC MC MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		COUMADIN TABS ENOXAPARIN ¹ ELIQUIS ELIQUIS STARTER PACK HEPARIN SODIUM/NACL 0.9% SOLN HEP-LOCK SOLN INNOHEP HEPARIN LOCK SOLN HEPARIN LOCK FLUSH SOLN HEPARIN SODIUM SOLN HEPARIN SODIUM LOCK FLUSH SOLN PRADAXA JANTOVEN WARFARIN SODIUM TABS XARELTO XARELTO STARTER PACK	MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		ARIXTRA SOLN FONDAPARINUX FRAGMIN INJ FRAGMIN VIAL LOVENOX SOLN LOVENOX 300 ² LOVENOX SUBQ SYRINGE PRADAXA ORAL PELLETS ⁴ IPRIVASK SAVAYSAS ³	Use PA form# 20420 1. Enoxaparin therapy durations greater than 7 days every 30 days require PA. 2. Use other strengths available to obtain desired dose. 3. Diagnosis required 4. For the treatment of patients aged 3 months to less than 12 years of age.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Exceeding days supply limits for LMWH class requires PA. DDI: Warfarin will require prior authorization if being used in combination with fluconazole, miconazole, or voriconazole. DDI: Warfarin will require prior authorization if being used in conjunction with gemfibrozil or fenofibrate. DDI: Rifampin will require prior authorization if being used in combination with savaysa.
ANTIHEMOPHILIC AGENTS	MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC MC MC/DEL MC MC MC MC/DEL MC MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC/DEL MC MC/DEL		ALPHANATE ALPHANINE SD ALPROLIX VIAL BEBULIN VIAL BENEFIX SOLR HELIXATE FS KIT HEMOFIL - M HUMATE-P SOLR IXINITY VIAL JIVI ³ KOATE-DVI KONYNE - 80 KOVALTRY REBINYN MONARC - M MONOCLATE - P MONONINE NOVOEIGHT NOVOSEVEN SOLR NUWIQ PROFILNINE RECOMBINATE SOLR REFACTO RIXUBIS VIAL WILATE INJ XYNTHA	MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC MC/DEL MC MC MC MC/DEL MC MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC/DEL		ADYNOVATE VIAL ADVATE ^{1,2,5} ALTUVIIIO ⁴ AFSTYLA BEQVEZ ESPEROCT ELOCTATE HEMGENIX IDELVION KOGENATE FS ⁵ RECOMBINATE VIAL ⁵ ROCTAVIAN ⁴ SEVENFACT	Use PA Form# 20420 1. Only if other products unavailable. 2. Advate may be available with PA in cases of large volume dosing in patients with poor venous access. 3. Not indicated for use in children <12 years of age due to greater risk for hypersensitivity reactions and is not indicated for use in previously untreated patients. 4. Clinical PA required for appropriate diagnosis. 5. Established users will be grandfathered.	Non-preferred will only be approved if other preferred products are unavailable. Beqvez - FDA Approved Indication: An adeno-associated virus vector-based gene therapy indicated for the treatment of adults with moderate to severe hemophilia B (congenital factor IX deficiency) who: • Currently use factor IX prophylaxis therapy, or • Have current or historical life-threatening hemorrhage, or • Have repeated, serious spontaneous bleeding episodes, and, • Do not have neutralizing antibodies to adeno-associated virus serotype Rh74var (AAVRh74var) capsid as detected by an FDA-approve test. Hemgenix is an adeno-associated viral vector-based gene therapy for IV infusion after dilution. For treatment of adults with Hemophilia B (congenital Factor IX deficiency) who: Currently use Factor IX prophylaxis therapy, or have current or historical life-threatening hemorrhage, or have repeated, serious spontaneous bleeding episodes. Altuviio is a von Willebrand Factor (VWF) independent recombinant DNA-derived, Factor VIII concentrate indicated for use in adults and children with hemophilia A (congenital factor VIII deficiency) for: Routine prophylaxis to reduce the frequency of bleeding episodes, On-demand treatment and control of bleeding episodes, Perioperative management of bleeding. Roctavian : For the treatment of adults with severe hemophilia A (congenital factor VIII deficiency with factor VIII activity <1 IU/dL) without antibodies to adeno-associated virus serotype 5 (AAV5) detected by an FDA-approved test. Inclusion: • Severe factor VIII deficiency (less than 1% native factor VIII). Exclusion Criteria: • Antibodies to the virus AAV5 • Factor VIII inhibitors (or history of) • Known significant fibrosis of cirrhosis of the liver, or unexplained elevated LFTs • History of inadequate compliance with prophylaxis, or regular bleeds despite adequate prophylaxis • Conditions in which high-dose steroids are contraindicated. • Inability to abstain from alcohol for one year • Plan to impregnate a partner within 6 months of infusion • Hypersensitivity to mannitol • Active infections, either acute or uncontrolled chronic • HIV infection (limited information on use in this population)
NON-FACTOR REPLACEMENT THERAPY	MC		HEMLIBRA	MC/DEL MC/DEL		ALHEMO HYMPAVZI QFITLIA QFITLIA PEN	Use PA Form# 20420	Subsequent changes made to Antihemophilic Agents: Factor Therapy to move Hemlibra to Non-Factor Therapy
PLATELET AGGREGATION INHIBITORS	MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL		ASPIRIN ASPIRIN-DIPYRIDAMOLE ER CPMP 12HR BRILINTA 90mg DIPYRIDAMOLE TABS CLOPIDOGREL 75MG PRASUGREL HCL TAB	MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC/DEL	7 8 8 8 8 8 8	TICLOPIDINE HCL TABS BRILINTA 60mg DURLAZA EFFIENT PERSANTINE TABS PLAVIX TABS ZONTIVITY	Use PA Form# 20715 for Plavix, Effient & Brilinta Use PA form# 20420 for other requests 1. Dosing limits apply, see Dose Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. A special PA may be obtained at the pharmacy for members scheduled for "stent" placement or have had placement if in the last 12months. Please indicate on prescription date of stent placement. Brilinta - Concomitant use with strong CYP3A4 inhibitors should be avoided (including ketoconazole, itraconazole, atazanavir, and telithromycin). Doses of simvastatin and lovastatin >40mg should be avoided. DDI : exists for using maintenance ASA dose >100mg, as it reduces the effectiveness of Brilinta. DDI: Plavix will require prior authorization if being used in combination with omeprazole, esomeprazole, cimetidine, fluconazole, ketoconazole, intelence, fluoxetine, ticlopidine, and fluvoxamine.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
PLATELET AGGR. INHIBITORS / COMBO'S - MISC.	MC/DEL MC/DEL		CILOSTAZOL PENTOXIFYLLINE ER TBCR	MC/DEL MC/DEL MC MC		AGRYLIN CAPS ANAGRELIDE CAPS PLETAL TABS TRENTAL TBCR YOSPRALA	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
HEMATOLOGICALS								
MONOCLONAL ANTIBODY				MC MC/DEL MC MC/DEL MC MC MC/DEL MC MC		BKEMV EMPAVELI ENSPRYNG EPYSQLI FABHALTA GAMIFANT PIASKY SOLIRIS ULTOMIRIS UPLIZNA VOYDEYA	Use PA Form# 20420	A diagnosis of Paroxysmal nocturnal hemoglobinuria (PNH) using the HAM test or flow cytometry is required. In addition, the patient must show evidence of having received a meningitis vaccine at least 2 weeks prior to the start of therapy. Gamifant is recommended for the treatment of adult and pediatric (newborn and older) patients with primary hemophagocytic lymphohistiocytosis (HLH) with refractory, recurrent, or progressive disease or intolerance with conventional HLH therapy. Fabhalta and Ultomiris are recommended for the treatment of adults with paroxysmal nocturnal hemoglobinuria (PNH). Bkemv and Epysqli have updated criteria for a diagnosis of generalized myasthenia gravis (gMG): must have confirmation that patients are anti-acetylcholine receptor (AChR) antibody positive.
IMMUNE GLOBULIN	MC MC/DEL MC MC MC/DEL MC/DEL MC		BIVIGAM ¹ CUTAQUIG ¹ GAMUNEX-C GAMMAPLEX S-D ¹ HIZENTRA ¹ PANZYGA ¹ PRIVIGEN ¹	MC MC MC/DEL MC MC/DEL MC MC/DEL		ALYGLO ASCENIV ² CUVITRU GAMMAPLEX INJ HYQVIA OCTAGAM INJ ¹ XEMBIFY	Use PA Form# 20420 1. Clinical PA required 2. For the treatment of patients between 12 to 17 years of age.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Alyglo is indicated for treatment of primary humoral immunodeficiency in adults ages 17 or older. Cutaquig is indicated as replacement therapy for primary humoral immunodeficiency (PI) in adults. Xembify is indicated for treatment of primary humoral immunodeficiency (PI) in patients 2 years of age and older. Asceniv indicated for the treatment of primary humoral immunodeficiency (PI) in adults and adolescents (12 to 17 years of age). PI includes but is not limited to the humoral immune defect in congenital agammaglobulinemia, common variable immunodeficiency (CVID), X-linked agammaglobulinemia, Wiskott-Aldrich syndrome, and severe combined immunodeficiencies (SCID).
HEREDITARY ANGIOEDEMA	MC MC MC MC/DEL MC/DEL MC MC/DEL		PROPHYLAXIS			PROPHYHLAXIS	Use PA Form# 20420 1. Clinical PA is required to establish diagnosis and medical necessity. 2. For the treatment of patients ≥ 12 years of age. Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Haegarda is indicated for routine prophylaxis to prevent Hereditary Angioedema (HAE) attacks in adolescent and adult patients
			CINRYZE ¹ HAEGARDA ¹ ORLADEYO ^{1,2} TAKHZYRO ¹	MC	8	ANDEMBRY		
			TREATMENT			TREATMENT		
			BERINERT KIT ¹ FIRAZYR ¹ RUCONEST VIAL ¹	MC/DEL		KALBITOR VIAL		
HEMATOLOGICAL AGENTS-THROMBOPOIETIN RECEPTOR AGONISTS	MC MC		PROMACTA ¹ NPLATE ¹	MC MC/DEL MC/DEL		ALVAIZ DOPTELET MULPLETA	Use PA Form# 20420 1. Clinical PA required. Must see prior trial with insufficient response to corticosteroids and immunoglobulins.	Doptelet and Mulpelta : For the treatment of thrombocytopenia in adults with chronic liver disease who are scheduled to undergo a procedure.
HEMATOLOGICAL AGENTS-IgAN				MC/DEL MC		FILSPARI ¹ TARPEYO VANRAFIA	Use PA Form# 20420 1. PA required to confirm FDA-approved indication.	All preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists PA required to confirm FDA-approved indication. Vanrafia is for adults with biopsy proven primary IgAN AND eGFR>=30 cc/min/1.73m3 AND urine protein >=1 g/day AND on stable dose of maximally tolerated renin-angiotensin system inhibitor.
ANEMIA- BETA THALASSEMIA				MC MC		REBLOZYL ZYNTEGLO	Use PA Form# 20420	Reblozyl is indicated for three (3) treatments of anemia in adults: 1. in adult patients with beta thalassemia who require regular red blood cell (RBC) transfusions; 2. without previous erythropoiesis stimulating agent use (ESA-naïve) in adult patients with very low- to intermediate-risk myelodysplastic syndromes (MDS) who may require regular RBC transfusions; and 3. failing an ESA and requiring 2 or more RBC units over 8 weeks in adult patients with very low- to intermediate-risk MDS with ring sideroblasts (MDS-RS) or with myelodysplastic/myeloproliferative neoplasm with ring sideroblasts and thrombocytosis (MDS/MPN-RS-T). It is not indicated for use as a substitute for RBC transfusions in patients who require immediate correction of anemia. Zynteglo is indicated for the treatment of adult and pediatric patients with β-thalassemia who require regular red blood cell (RBC) transfusions.
HEMATOLOGIC DISORDER TREATMENT AGENTS				MC/DEL MC		CABLIVI TAVALISSE	Use PA Form# 20420	Tavalisse is recommended for patients at risk of bleeding when one line of therapy (steroids, IVIG, splenectomy) has failed. Cablivi is recommended for the treatment of adult patients with acquired thrombotic thrombocytopenic purpura (aTTP), in combination with plasma exchange and immunosuppressive therapy.
COMPLEMENT RECEPTOR ANTAGONIST				MC		TAVNEOS	Use PA Form# 20420	
WHIM SYNDROME AGENTS				MC		XOLREMDI	Use PA Form# 20420	Xolremdi : In patients 12 years of age and older with WHIM syndrome (warts, hypogammaglobulinemia, infections, and myelokathexis) to increase the number of circulating mature neutrophils and lymphocytes.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
HEMOSTATIC								
HEMOSTATIC	MC/DEL MC		AMICAR AMINOCAPROIC ACID	MC MC		FIBRYGA RIASTAP	Use PA Form# 20420	Fibryga and Riastap are indicated for the treatment of acute bleeding episodes in adults and adolescents with congenital fibrinogen deficiency, including afibrinogenemia and hypofibrinogenemia. Fibryga is not indicated for dysfibrinogenemia.
ACUTE HEPATIC PORPHYRIA (AHP)								
ACUTE HEPATIC PORPHYRIA (AHP)				MC		GIVLAARI	Use PA Form# 20420	Givlaari is indicated for the treatment of adults with acute hepatic porphyria (AHP).
PYRUVATE KINASE DEFICIENCY AGENTS								
PYRUVATE KINASE DEFICIENCY AGENTS				MC		PYRUKYND ¹	Use PA Form# 20420 1. PA required to confirm FDA approved indication.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s).
OP. - ANTIBIOTICS	MC MC MC/DEL MC MC/DEL MC MC MC/DEL MC/DEL		AK-SPORE OINT BACITRACIN/NEOMYCIN/POLYM BACITRACIN/POLYMYXIN B OINT CHLOROPTIC SOLN ERYTHROMYCIN OINT NEOSPORIN SOLN POLYSPORIN TRIMETHOPRIM SULFATE/POLY TOBRAMYCIN SULFATE SOLN	MC MC MC MC MC MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC MC MC MC/DEL MC/DEL MC		AK-POLY-BAC OINT AK-SULF OINT AK-TOB SOLN AZASITE BACITRACIN OINT BLEPH-10 SOLN GATIFLOXACIN DROPS GENTAMICIN SULFATE GENTAK ILOTYCIN OINT LEVOFLOXACIN DROPS NEOMYCIN/BACI/POLYM OINT NEOMYCIN/POLYMYXIN/GRAMIC NEOSPORIN OINT OCUSULF-10 SOLN OCUTRICIN SOLN POLYTRIM DROPS SULFACETAMIDE SODIUM DROPS SULFACETAMIDE SODIUM OINT TERAK OINT	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
OP. - ANTI-PARASITIC				MC		XDEMYV ¹	Use PA Form# 20420 1. For the treatment of Demodex blepharitis.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
OP. - RHO KINASE INHIBITORS	MC		RHOPRESSA				Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
OP. - QUINOLONES	MC/DEL MC/DEL MC/DEL MC/DEL		CILOXAN OINT CIPROFLOXACIN SOL 0.3% OFLOXACIN QUIXIN SOLN	MC/DEL MC/DEL MC		BESIVANCE CILOXAN SOLN OCUFLOX SOLN	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
OP. - QUINOLONES-4TH GENERATION	MC/DEL		MOXIFLOXACIN 0.5% SOLN (Generic Vigamox)	MC		ZYMAXID	Use PA Form# 20420	
OP. - ARTIFICIAL TEARS AND LUBRICANTS	MC/DEL MC/DEL MC MC MC/DEL MC MC MC MC MC MC MC MC		ARTIFICIAL TEARS OINT ARTIFICIAL TEARS SOLN CELLUVISC SOLN EYE LUBRICANT OINT GENTEAL LIQUITEARS SOLN MAJOR TEARS SOLN PURALUBE OINT PURALUBE TEARS SOLN REFRESH SOLN OP REFRESH PLUS SOLN ¹ REFRESH PM OINT	MC/DEL MC MC MC MC/DEL MC/DEL MC MC MC MC MC/DEL MC/DEL MC MC MC MC MC/DEL MC/DEL		ARTIFICIAL TEARS SOLN OP BION TEARS SOLN DRY EYES OINT DURATEARS OINT HYPO TEARS ISOPTO TEARS SOLN LACRI-LUBE LUBRIFRESH P.M. OINT MURINE SOLN MUROCEL SOLN NATURE'S TEARS SOLN REFRESH SOLN REFRESH TEARS SOLN ¹ TEARGEN SOLN TEARISOL SOLN TEARS NATURALE TEARS PURE SOLN	Use PA Form# 20420 1. Dosing limits apply, see Dose Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
				MC MC/DEL MC		TEARS RENEWED OINT THERATEARS SOLN V-R ARTIFICIAL TEARS SOLN		
OP. - BETA - BLOCKERS	MC/DEL MC/DEL MC/DEL MC/DEL		BETOPTIC-S SUSP CARTEOLOL HCL SOLN LEVOBUNOLOL HCL SOLN METIPRANOLOL SOLN	MC MC/DEL MC MC/DEL MC MC/DEL MC MC/DEL MC/DEL		BETAGAN SOLN BETAXOLOL HCL SOLN ISTALOL OCUPRESS SOLN OPTIPRANOLOL SOLN TIMOPTIC SOLN TIMOLOL DROP TIMOLOL SOL-GEL TIMOPTIC-XE SOLG	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
OP. - ANTI-INFLAMMATORY / STEROIDS OPHTH.	MC MC/DEL MC/DEL MC/DEL MC MC MC MC/DEL MC/DEL MC/DEL MC MC MC MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC		AK-SPORE HC OINT ALREX SUSP DEXAMETH SOD PHOS SOLN FLUORMETHOLONE SUSP FML DROPS SUSP 1% FML FORTE SUSP FML S.O.P. OINT LOTEMAX OINT LOTEMAX GEL LOTEMAX SUSP NEO/POLY/DEXAMETH OINT NEO/POLY/DEXAMETH SUSP PRED-G SUSP PRED FORTE SUSP 1% PRED MILD SUSP PREDNISOLONE TOBRADEX OINT TOBREX OINT SULFACETAMIDE/PREDNISOLONE ZYLET SUSP	MC MC MC MC MC MC MC/DEL MC MC/DEL MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC		AK-TROL SUSP BAC/POLY/NEOMY/HC OINT BLEPHAMIDE S.O.P. OINT BLEPHAMIDE SUSP BROMDAY EFLONE SUSP FLAREX SUSP FLUOR-OP SUSP ILUVIEN IMPLANT INVELTYS LOTEMAX SM DROPS GEL 0.38% MAXITROL OPTH OINT 0.1% NEO/POLY/BAC/HC OINT NEOM/POLY/DEX OPTH OINT 0.1% OMNIPRED DROPS SUSP OZURDEX PRED-G S.O.P. OINT PREDNISOLONE SODIUM PHOSHATE SOL RETISERT IMPLANT SULFACET SOD/PRED SOLN TRIESENCE VIAL TOBRADEX ST TOBRAMYCIN SUSP DEXAMETHASONE VASOCIDIN SOLN VEXOL SUSP XIPERE	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
OP. - PROSTAGLANDINS	MC/DEL MC MC/DEL MC/DEL		LATANOPROST SOL 0.005% LUMIGAN SOLN ROCKLATAN TRAVATAN-Z	MC/DEL MC/DEL MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL	7 8 8 8 8 8 8 8 8 8 8 8 8	ZIOPTAN BIMATOPROST 0.03% DROPS DURYSTA IYUZEH RESCULA ^{1,2,3} TRAVATAN SOLN TRAVOPROST VYZULTA XALATAN SOLN ¹ XELPROS	Use PA Form# 20420 1. All preferred must be tried. 2. Dosing limits apply, see Dosing Consolidation List. 3. Clinical PA is required to establish diagnosis and medical necessity.	Preferred drugs must be tried and failed, in step-order, due to lack of efficacy (failure to reach target IOP reduction) or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
OP. - CYCLOPLEGICS	MC MC/DEL MC/DEL MC/DEL		AK-PENTOLATE SOLN ATROPINE SULFATE CYCLOPENTOLATE HCL SOLN ISOPTO HYOSCINE SOLN	MC/DEL MC MC/DEL MC		CYCLOGYL SOLN ISOPTO ATROPINE SOLN ISOPTO HOMATROPINE SOLN MUROCOLL-2 SOLN	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
OP. - MIOTICS - DIRECT ACTING	MC/DEL MC MC MC/DEL MC/DEL		ISOPTO CARBACHOL SOLN ISOPTO CARPINE SOLN PILOCAR SOLN PILOCARPINE HCL SOLN PILOPINE HS GEL				Use PA Form# 20420	

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
OP. - SELECTIVE ALPHA ADRENERGIC AGONISTS	MC MC MC MC/DEL MC/DEL		ALPHAGAN SOLN ALPHAGAN P 0.1% SOLN ALPHAGAN P 0.15% SOLN BRIMONIDINE DROPS 0.2 % SIMBRINZA	MC/DEL MC/DEL		BRIMONIDINE TARTRATE DROPS 0.15 % IOPIDINE SOLN	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
OP. - ANTI-ALLERGICS	MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL		AZELASTINE HCL DROPS BEPREVE CROMOLYN SODIUM DROPS KETOTIFEN FUMARATE DROPS LASTACRAFT OLOPATADINE HCL 0.1% OLOPATADINE HCL 0.2% ZADITOR SOLN	MC MC/DEL MC/DEL MC MC/DEL MC MC/DEL	8 8 8 8 8 8 9	ALOCRI SOLN ALOMIDE SOLN EMADINE SOLN OPTICROM SOLN PATANOL SOLN ZERVIAE EPINASTINE	Use PA Form# 20420	All preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
OP. ANTI-ALLERGICS- MASTCELL STABILIZER CLASS				MC/DEL		ALAMAST SOLN	Use PA Form# 20420	
OP. - CARBONIC ANHYDRASE INHIBITORS/COMBO	MC/DEL MC MC/DEL MC/DEL		AZOPT SUSP COMBIGAN DORZOLAMIDE DORZOLAMIDE/TIMOLOL	MC/DEL		COSOPT SOLN PF	Use PA Form# 20420	
OP. - NSAID'S	MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		ACULAR SOLN ¹ DUREZOL KETOROLAC OPTH 0.4% KETOROLAC OPTH 0.5% MAXIDEX SUSP NEVANAC PREDNISOLONE DROPS	MC MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC MC MC MC/DEL	8 8 8 8 8 8 8 8 8 8 8 9	ACULAR LS ¹ BROMSITE ¹ DEXAMETHASONE DROPS DICLOFENAC OPTH 0.1% FLURBIPROFEN SODIUM SOLN ILEVRO LOTEMAX SM DROPS GEL 0.38% PROLENSA OCUFEN SOLN ¹ XIBROM ¹ VOLTAREN SOLN ¹ ACUVAIL ¹ BROMFENAC	Use PA Form# 20420 1. Must fail all preferred products before non-preferred.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
OP. - OF INTEREST	MC MC MC MC		EYSUVIS ² LUCENTIS RESTASIS DROPPERETTE XIIDRA	MC MC MC MC/DEL MC MC MC MC MC MC MC MC/DEL MC/DEL MC MC/DEL MC MC MC MC MC MC		BYOOVIZ BEOVU BOTOX SOLR CEQUA CIMERLI CYCLOSPORINE DROPERETTE CYSTADROPS ¹ CYSTARAN ¹ EYLEA EYLEA HD ¹ IZERVAY ¹ LUCENTIS LUXTURNAL MIEBO OXERVATE PAVBLU RESTASIS MULTIDOSE DROPS SUSVIMO SYFOVRE TRYPTYR ¹ TYRVAYA VABYSMO VERKAZIA VEVYE	Use PA Form# 20420 1. PA required to confirm appropriate diagnosis and clinical parameters for use. 2. For the short-term (up to two weeks) treatment of the signs and symptoms of dry eye disease.	Must fail adequate trials of multi agents from artificial tears and lubricant category. Beovu is non-preferred and indicated for the treatment of Neovascular (wet) Age-Related Macular Degeneration (AMD) Eylea is non-preferred and indicated for the treatment of: Neovascular (wet) Age-Related Macular Degeneration (AMD), Macular Edema following Retinal Vein Occlusion (RVO), Diabetic Macular Edema (DME), Diabetic Retinopathy (DR) Luxturna will be considered for the treatment of patients with confirmed biallelic RPE65 mutation-associated retinal dystrophy. Patients must have viable retinal cells as determined by the treating physician(s). Miebo is non-preferred and is indicated for the treatment of the signs and symptoms of dry eye disease (DED). Oxervate is non-preferred and is indicated for the treatment of neurotrophic keratitis. Pavblu : Clinical rationale for why eylea cannot be used Syfovre is non-preferred and is indicated for the treatment of geographic atrophy (GA) secondary to age-related macular degeneration (AMD). Vevye - Must fail adequate trials of multi agents from artificial tears and lubricant category and a preferred cyclosporine alternative.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
DERMATOLOGICAL								
ISOTRETINION, ACNE	MC MC MC MC		AMNESTEEM ¹ CLARAVIS ¹ MYORISAN ¹ ZENATANE ¹	MC MC		ABSORICA ABSORICA LD	Use PA Form# 20420 1. Users 24 or under, PA will not be required.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - ACNE PREPARATIONS	MC MC/DEL MC/DEL MC/DEL MC MC MC MC/DEL MC		ERYDERM SOLN ERYTHROMYCIN GEL ERYTHROMYCIN SOLN EVOCLIN ISOTRETINOIN METRONIDAZOLE CREA ² METRONIDAZOLE GEL ² METRONIDAZOLE LOTN ² TRETINOIN .025%, .05%, .01% GEL ¹ TRETINOIN CREA ^{1,2}	MC/DEL MC/DEL MC MC/DEL MC MC MC MC/DEL MC/DEL MC/DEL MC MC MC MC MC/DEL MC MC MC/DEL MC MC MC MC/DEL MC MC/DEL MC MC MC MC MC/DEL MC MC/DEL MC MC MC MC MC/DEL MC MC/DEL MC MC MC MC MC/DEL MC MC/DEL MC MC MC MC MC/DEL MC MC/DEL MC MC MC MC/DEL MC		ADAPALENE 0.3% GEL AKLIEF ⁶ ALTINAC CREA ALTRENO AMZEEQ ⁶ ARAZLO LOTION ⁶ AVITA CREA BENZAC BENZACLIN GEL ³ BENZAGEL-10 GEL BENZAMYCIN GEL BENZAMYCINPAK PACK BENZEOFAM BENZOYL PEROXIDE BREVOXYL CABTREO GEL ⁵ CLEOCIN-T ² CLINAC BPO GEL CLINDAGEL GEL CLINDAMYCIN PHOSPHATE CREAM ² CLINDETS SWAB DESQUAM-E GEL DESQUAM-X DIFFERIN 0.3% GEL DIFFERIN EMGEL GEL EPIDUO EPSOLAY ERYCETTE PADS FINEVIN CREA KLARON LOTN METROCREAM CREA ² METROGEL GEL ² METROLOTION LOTN ² NEOBENZ MICRO NORITATE CREA ONEXTON ⁵ PLIXDA RETIN-A GEL ² RETIN-A CREA ² RETIN-A MICRO GEL RHOFADÉ SODIUM SULFACET/SULF LOTN SOOLANTRA ⁴ TRIAZ TWYNEO VELTIN WINLEVI ⁵ ZENCIA WASH ZETACET ZIANA ZILXI	Use PA Form# 10220 for Brand Name requests Use PA Form# 20420 for all other requests 1. Users 24 or under, PA will not be required. 2. Dosing limits allowing one package per month. Refer to Dose Consolidation List. 3. Only available if component ingredients are unavailable. 4. Dosing limits apply, see Dosing Consolidation List. 5. Not approved for use in children <12 years of age. 6. For the treatment of patients ≥ 9 years of age.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
TOPICAL- ATOPIC DERMATITIS	MC/DEL	1	ELIDEL CREA	MC/DEL		CIBINQO	Use PA Form# 20420 1. Avoid live vaccines if treated with Dupixent. 2. Clinical PA required. 3. For the treatment of patients ≥ 12 years of age. 4. Preferred after a trial and failure of TCI.	Preferred drugs also indicated for this condition, including topical steroids, cyclosporin AND calcineurin inhibitors must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Note: If unable to use TCIs then a trial of Eucrisa could be recommended before Dupixent.
	MC/DEL	1	PIMECROLIMUS CRE (AUTH GENERIC LABELER 68682 Oceanside Pharmaceuticals)	MC MC		EBGLYSS ^{2,3} NEMLUVIO		
	MC/DEL	1	PROTOPIC OINT					
	MC/DEL	1	TACROLIMUS OINT					
	MC	2	ADBRY ^{2,4}					
	MC/DEL	2	DUPIXENT ^{1,2,4}					
	MC	2	EUCRISA ^{2,4}					
	MC	2	OPZELURA ^{2,3,4}					
TOPICAL - ANTIBIOTIC	MC		BACIT/NEOMYCIN/POLYM OINT	MC/DEL		CENTANY OINT 2% ¹	Use PA Form# 20420 1. Dosing limits apply, see Dosing Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL		BACITRACIN OINT	MC/DEL		MUPIROCIN CREA ¹		
	MC/DEL		GENTAMICIN SULFATE	MC/DEL		TRIPLE ANTIBIOTIC OINT		
	MC/DEL		MUPIROCIN OINT ¹	MC		XEPI		
TOPICAL - ANTIFUNGALS	MC/DEL		BETAMETHASONE CLOTRIMAZOLE CREA	MC/DEL	8	CICLOPIROX SOLN	Use PA Form# 10120 1. Diagnosis required.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Kerydin- Verify prior trials and failures or intolerance of preferred treatments, including both topical and oral agents DDI: Ketoconazole will now be non-preferred and require prior authorization if they are currently being used in combination with any of the following medications: prevacid, pantoprazole, onglyza or omeprazole.
	MC/DEL		BETAMETHASONE CLOTRIMAZOLE LOT	MC MC/DEL	8	EXELDERM		
					8	FUNGIZONE CREA		
	MC		CICLOPIROX 0.77 CREA	MC	8	HYDROCORT/ODOQ CREA		
	MC		CICLOPIROX 0.77 SUSP	MC	8	JUBLIA		
	MC/DEL		CLOTRIMAZOLE	MC/DEL	8	KERYDIN ¹		
	MC		ECONAZOLE NITRATE CREA	MC/DEL	8	LOPROX 0.77 LOTN		
	MC/DEL		KETOCONAZOLE CREA	MC/DEL	8	LOPROX 0.77 CREA		
	MC/DEL		KETOCONAZOLE SHAM	MC/DEL	8	LOPROX 0.77 SUSP		
	MC/DEL		LOPROX 1.0 CREA	MC	8	LOPROX SHAMPOO SHAM		
	MC/DEL		LOPROX 1.0 LOTN	MC/DEL	8	LOTRIMIN		
	MC/DEL		LOPROX GEL	MC/DEL	8	LOTRISONE LOT		
	MC/DEL		LOPROX TS LOTN	MC	8	LOTRISONE CREA		
	MC/DEL		MICONAZOLE NITRATE CREA	MC/DEL	8	LUZU		
	MC		MYCO-TRIACET II CREA	MC	8	MENTAX CREA		
	MC/DEL		NYSTATIN	MC	8	MYCOGEN II CREA		
	MC/DEL		NYSTATIN/TRIAMCINOLONE CREA	MC	8	NAFTIN		
	MC/DEL		NYSTOP POWD	MC/DEL	8	NIZORAL SHAM		
	MC		TRI-STATIN II CREA	MC	8	NYSTATIN/TRIAMCINOLONE OINT		
				MC/DEL	8	NYSTAT-RX POWD		
				MC/DEL	9	OXISTAT		
				MC/DEL		PENLAC NAIL LACQUER SOLN		
TOPICAL - ANTIPRURITICS	MC		ZONALON CREA	MC MC		KORSUVA	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
						PRUDOXIN CREA		
TOPICAL - ANTIPSORIATICS	MC/DEL		CALCIP/BETAMETHASONE SUS	MC/DEL	7	TACLONEX ¹	Use PA Form# 20420 1. Must fail all preferred products before non-preferred.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
				MC/DEL	8	DUOBRII		
				MC	8	ENSTILAR		
				MC	8	OXSORALEN ULTRA CAPS ¹		
				MC	8	PSORiatec CREA ¹		
				MC/DEL	8	SORIATANE CK KIT ¹		
				MC	8	VECTICAL ¹		
				MC	8	VTAMA		
				MC	8	ZORYVE		
TOPICAL - ANTISEBORRHEICS	MC/DEL		SELENIUM SULFIDE SHAM	MC MC MC		CARMOL SCALP TREATMENT KIT	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Zoryve Foam: For the treatment of seborrheic dermatitis in adult and pediatric patients 9 years of age and older.
						ZNP BAR		
						ZORYVE FOAM		

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
TOPICAL - ANTIVIRALS				MC/DEL MC/DEL MC MC MC		ACYCLOVIR OINT DENAVIR CREA ^{1,3} YCANTH ZELSUVMI ⁴ ZOVIRAX OINT ^{1,2}	Use PA Form# 20420 1. Must fail oral treatment with Acyclovir or Valacyclovir . 2. Approvals limited to 1 tube per 180 days. 3. Dosing limits apply, see Dosing Consolidation List. 4. For the topical treatment of molluscum contagiosum in adult and pediatric patients 1 year of age and older.	
TOPICAL - ANTINEOPLASTICS	MC		EFUDEX	MC/DEL MC/DEL MC MC/DEL		CARAC CREA FLUOROURACIL SOLARAZE GEL ZYCLARA	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - BURN PRODUCTS	MC MC/DEL MC MC MC/DEL		FURACIN CREA SILVER SULFADIAZINE CREA SSD AF CREA SSD CREA THERMAZENE CREA	MC/DEL		SILVADENE CREA	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - CORTICOSTEROIDS	MC MC/DEL MC MC MC MC/DEL MC MC MC MC MC MC/DEL MC		<div>LOW POTENCY</div> DERMA-SMOOTHIE- FS BODY HYDROCORTISONE CREA HYDROCORTISONE LOTN HYDROCORTISONE LOTN TEXACORT SOLN <div>MEDIUM POTENCY</div> DESOXIMETASONE 0.05% CREA/GEL FLUTICASONE PROPIONATE CREA/OINT HYDROCORTISONE BUTYRATE HYDROCORTISONE OINT HYDROCORTISONE VALERATE MOMETASONE FUROATE OINT TRIAMCINOLONE ACETONIDE .025-.1% <div>HIGH POTENCY</div> DESONIDE ¹ TRIAMCINOLONE ACETONIDE .5%	MC/DEL MC MC MC/DEL MC/DEL MC MC MC MC/DEL MC/DEL MC MC MC MC/DEL MC MC MC MC/DEL		<div>LOW POTENCY</div> ACLOVATE ANUSOL HC-1 OINT DESONATE GEL FLUOCINOLONE ACETONIDE FLUOCINOLONE HALOG HYDROCORTISONE POWD LIDA MANTLE HC CREA PROCTOCORT CREA VERDESO <div>MEDIUM POTENCY</div> BESER LOTION ³ CLODERM CREA CORDRAN CUTIVATE CREA / OINT CUTIVATE LOTN DERMATOP ELOCON OINT KENALOG AERS LOCOID LUXIQ FOAM PANDEL CREA TOPICORT TOPICORT LP CREA TOVET FOAM ³ WESTCORT <div>HIGH POTENCY</div> AMCINONIDE CREA BETAMETHASONE DIPROPIONATE DESOXIMETASONE 0.25% CREA/OINT	Use PA Form# 20420 1. Dosing limits apply, see Dosing Consolidation List. 2. Treatment beyond 4 weeks is not recommended. 3. For the treatment of patients ≥ 12 years of age. 4. For the treatment of patients ≥ 18 years of age.	At least 1 drug from each potency of preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
	MC/DEL MC/DEL MC MC		VERY HIGH POTENCY	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC		VERY HIGH POTENCY		
			AUGMENTED BETA DIP			BRYHALI LOTN		
			BETAMETHASONE VALERATE			CLOBETASOL PROPINATE LOTN		
			DIFLORASONE DIACETATE			CLOBETASOL PROPINATE SHAMPOO 0.05%		
			HALOBETASOL			CORMAX		
						DIPROLENE		
	MISCELLANEOUS	IMPEKLO ⁴						
MC		PROCTO-KIT CREA 1%				LEXETTE		
					OLUX FOAM			
						PSORCON		
						PSORCON E		
						SERNIVO SPRAY ²		
						TEMOVATE		
						ULTRAVATE		
TOPICAL - STEROID LOCAL ANESTHETICS				MC		EPIFOAM FOAM	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - STEROID COMBINATIONS	MC		DERMA-SMOOTHIE-FS SCALP	MC		CARMOL-HC CREA	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - EMOLLIENTS	MC/DEL MC MC		AMMONIUM LACTATE CREA ¹ AMMONIUM LACTATE LOTN 12% ¹ VITAMIN A & D MEDICATED OINT	MC MC MC MC		LAC-HYDRIN CREA ¹ LAC-HYDRIN LOTN 12% MEDERMA GEL MIMYX RENOVA CREA	Use PA Form# 20420 1. Dosing limits still apply, see Dose Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - ENZYMES / KERATOLYTICS / UREA				MC MC MC		CARMOL 40 CREA SALEX CREA SALEX LOTN	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Ziox, Panafil and Papain products have been removed from the PDL due to FDA safety concerns regarding drugs containing Papain.
TOPICAL - GENITAL WARTS	MC/DEL		IMIQUIMOD 5% ²	MC/DEL MC/DEL MC/DEL MC MC MC	5 8 8 8 8 8	PODOFILOX SOLN CONDYLOX ¹ ALDARA ¹ PICATO VEREGEN ¹ ZYCLARA ¹	Use PA Form# 20420 1. Non-preferred products must be used in specified order. 2. Dosing limits still apply, see Dose Consolidation List.	
TOPICAL - LOCAL ANESTHETICS	MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL		AF CAPSICUM OLEORESIN CREA CAPSAICIN CREA CAPSAICIN PATCH DIBUCAINE OINT ELA-MAX ¹ LIDOCAINE/PRILOCAINE CREA ¹ LIDOCAINE CREAM LIDOCAINE GEL LIDOCAINE PTCH 5%	MC/DEL MC/DEL MC MC MC MC MC/DEL		EMLA PADS EMLA CREA LIDA MANTLE CREA PONTOCAINE SOLN SYNERA ZOSTRIX ZTLIDO ²	Use PA Form# 20420 1. Lidocaine/Prilocaine cream and Ela-Max products require PA for users over 18 years of age. 2. Dosing limits still apply, see Dose Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - DEPIGMENTING AGENTS				MC MC MC MC/DEL MC/DEL MC MC MC	8 8 8 8 8 8 8 9	ALUSTRA CREA EPIQUIN MICRO GLYQUIN CREA HYDROQUINONE CREA HYDROQUINONE/SUNSCREENS SOLAQUIN FORTE CREA TRI-LUMA CREA ELDOQUIN	Use PA Form# 20420	As per Medicaid Policy, cosmetic drugs are not covered. Non-cosmetic clinical applications will be considered by prior authorization on a case by case basis.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
TOPICAL - SCABICIDES AND PEDICULICIDES	MC/DEL MC MC/DEL MC/DEL MC		ACTICIN CREA LICE KILLING SHAM LICE TREATMENT CREME RINS LIQD PERMETHRIN LOTN NATROBA ¹	MC MC MC/DEL MC MC MC/DEL		ELIMITE CREA EURAX LINDANE MALATHION OVIDE LOTN SPINOSAD SUSP	Use PA Form# 20420 1. Dosing limits apply, see Dosage Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - WOUND / DECUBITUS CARE				MC MC MC		FILSUVEZ REGRANEX GEL VYJUVEK	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Regranex will be approved for diabetic patients in good control (hgba1c <8), who are not smoking, with a stage III or IV WOCN AND NPUAP lower extremity diabetic ulcer and with an adequate blood supply (Tc p 02 >30, ABI>0.7 or ASP> 70), and where the underlying cause has been corrected. The wound must be free of infection and have been previously treated with preferred standard therapies for at least 2 months. Maximum approval for 20 weeks. Accuzyme and Ethezyme products have been removed from the PDL due to FDA concerns regarding drugs containing papain. Filsuvez: The patient has a diagnosis of dystrophic or junctional epidermolysis bullosa. The patient is at least 6 months old and does not have current evidence or history of squamous cell carcinoma or active infection in the area requiring filsuvez application. The patient has used standard wound care treatments, including silicone or foam dressings without wound resolution Vyjuvek: For the treatment of wounds in patients 6 months of age and older with dystrophic epidermolysis bullosa (DEB) with mutation(s) in the collagen type VII alpha 1 chain (COL7A1) gene.
TOPICAL - ASTRINGENTS / PROTECTANTS				MC MC MC		MOISTURIN DRY SKIN CREA PROSHIELD PLUS SKIN PROTE CREA SURGILUBE GEL	Use PA Form# 20420 1. Dosing limits apply, see Dosage Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
HYPERHIDROSIS THERAPY - AXILLARY	MC		XERAC AC SOLN	MC	8	SOFDRA ^{1,2}	1. Clinical PA is required to establish diagnosis and medical necessity. 2. For adults and pediatric patients 9 years of age and older.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. SOFDRA: prescribed by a dermatologist.
TOPICAL - ANTISEPTICS / DISINFECTANTS	MC/DEL		POVIDONE-IODINE SOLN	MC MC MC MC		BETADINE OINT FORMALYDE-10 AERS IODOSORB LAZERFORMALYDE SOLUTION SOLN	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
MISCELLANEOUS EYE								
OP. - EYE	MC MC MC MC MC MC/DEL		AK-DILATE SOLN EYE WASH SOLN NAPHAZOLINE HCL SOLN PHENYLEPHRINE HCL SOLN PONTOCAINE SOLN SODIUM CHLORIDE	MC MC/DEL MC		LENS PLUS REWETTING DROPS MURO 128 NEO-SYNEPHRINE SOLN	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
MISCELLANEOUS EAR								
EAR	MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC MC/DEL MC MC/DEL MC/DEL		A/B OTIC SOLN ACETASOL SOLN ACETASOL HC SOLN ACETIC ACID ACETIC ACID/HYDROCORTISON ALLERGEN SOLN CARBAMIDE PEROXIDE 6.5% OTIC SOLN. CIPRO HC SUSP CORTISPORIN-TC SUSP CORTOMYCIN COLY-MYCIN-S SUSP EAR DROPS SOLN EAR DROPS RX SOLN EAR WAX REMOVAL DROPS FLUOCINOLONE ACETONIDE OIL DROPS 0.01% NEOMYCIN/POLYMYXIN/HC OFLOXACIN 0.3% OTIC	MC MC MC/DEL MC/DEL MC/DEL MC MC MC MC MC MC MC MC MC/DEL MC/DEL		ANTIBIOTIC EAR SOLN ANTIBIOTIC EAR SUSP CIPRODEX CIPROFLOXACIN HCL DEBROX SOLN DERMOTIC FLOXIN OTIPRIO OTOVEL	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
MOUTH ANTISEPTICS								
MOUTH ANTI-INFECTIVES	MC MC/DEL		NILSTAT SUSP NYSTATIN SUSP	MC MC		MYCELEX TROC ORAVIG	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
MOUTH ANTISEPTICS	MC/DEL MC/DEL MC MC		CHLORHEXIDINE GLUCONATE LIDOCAINE VISCOUS SOLN TRIAMCINOLONE IN ORABASE PSTE TRIAMCINOLONE ORADENT PSTE	MC MC MC		APHTHASOL PSTE ¹ PERIOGARD SOLN ¹ TRIAMCINOLONE ACETONIDE PSTE ¹	Use PA Form# 20420 1. Must fail all preferred products before non-preferred.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
DENTAL PRODUCTS								
DENTAL PRODUCTS	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC		ETHEDENT CREA GEL-KAM CONC GEL-KAM GEL 0.4% PHOS FLUR SOLN SF 5000 PLUS CREA SF GEL STANNOUS FLUORIDE ORAL RI CONC	MC0MC MC/DEL MC/DEL MC		APF GEL DENTAGEL GEL PHOS-FLUR GEL THERA-FLUR-N GEL	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ARTIFICIAL SALIVA/STIMULANTS								
ARTIFICIAL SALIVA/ STIMULANTS	MC		SALIVA SUBSTITUTE SOLN	MC MC MC		EVOXAC CAPS RADIACARE SOLR SALAGEN TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
MISCELLANEOUS ANORECTAL								
ANORECTAL - MISC.	MC MC MC/DEL MC/DEL MC/DEL		CORTENEMA ENEM ELA-MAX 5 CREA HYDROCORTISONE ENEM PROCTOSOL HC CREA PROCTOZONE-HC CREA	MC/DEL MC/DEL MC/DEL MC/DEL MC		ANUSOL-HC CREA CORTIFOAM FOAM PROCTOFOAM HC FOAM PROCTO-KIT CREA 2.5% RECTIV OINT	Use PA Form# 20420	
T-CELL ACTIVATION INHIBITOR								
PSORIASIS BIOLOGICALS	MC MC MC MC MC/DEL MC		ADALIMUMAB-FKJP ENBREL ^{1,5} ENBREL SURECLICK ¹ HUMIRA ^{1,5} OTEZLA SIMLANDI SKYRIZI ⁶ TALTZ ²	MC MC/DEL MC MC/DEL MC MC/DEL MC MC/DEL MC MC MC MC/DEL MC MC MC MC MC MC		AMJEVITA BIMZELX ³ COSENTYX ⁴ CYLTEZO HADLIMA HULIO HYRIMOZ IDACIO ILUMYA ³ IMULDOSA OTULFI PYZCHIVA SELARSDI SILIQ SOTYKTU SPEVIGO STELARA STEQEYMA TREMIFYA YESINTEK YUFLYMA YUSIMRY	Use PA Form# 20910 1. Dosing limits apply, see Dosage Consolidation List. 2. Clinical PA required and will be preferred for the indication of plaque psoriasis, psoriatic arthritis and ankylosing spondylitis. 3. For the treatment of adults with moderate-to-severe plaque psoriasis who are candidates for systemic therapy or phototherapy. 4. Please see Criteria section. 5. Will not require a PA if at least one systemic drug such as methotrexate, cyclosporine, methoxsalen or acitretin is in members drug profile. 6. Clinical PA required and will be preferred for the indication of plaque psoriasis, psoriatic arthritis, Crohn's disease and ulcerative colitis.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Cosentyx approvals for 300mg dose(s) must use "300DOSE" package (containing 2 x 150mg pens or syringes). It is recommended to assess for TB infection prior to starting treatment with Taltz. Stelara will require using preferred trial of Skyrizi if unable please provide clinical rational as why inappropriate.
ALTERNATIVE MEDICINES								
ALTERNATIVE MEDICINES	MC MC		DIMETHYL SULFOXIDE SOLN MELATONIN	MC/DEL		CO-ENZYME Q-10	Use PA Form# 20420	Will only be approved for specific conditions supported by at least two double-blinded, placebo-controlled randomized trials that are not contradicted by other studies of similar quality.
CHELATING AGENTS								
CHELATING AGENTS	MC/DEL		CUPRIMINE CAPS	MC MC MC/DEL MC MC/DEL		CLOVIQUE DEPEN TITRATABS TABS EXJADE ¹ SYPRINE TRIENTINE CAPS	Use PA Form# 20420 1. FDA indication of treatment of chronic iron overload due to blood transfusions in members 2 years of age and older is required for approval of Exjade.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Clovique should be used when continued treatment with penicillamine is no longer possible because of intolerable or life endangering side effects.
ANTILEPTIC								
ANTILEPTIC				MC		THALOMID CAPS ¹	Use PA Form# 20420 1. All PA requests for 150mg dosing will require use of Thalomid 100mg and 50mg capsules.	Approved for indications of leprosy, treatment-resistant multiple myeloma and AIDS.

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
ANTINEOPLASTIC AGENTS								
ANTINEOPLASTIC AGENTS - ANTIADNDROGENS	MC/DEL		BICALUTAMIDE	MC/DEL		CASODEX	Use PA Form# 20420	
ANTINEOPLASTIC AGENTS- LHRH ANALOGS	MC/DEL MC/DEL MC/DEL MC/DEL		LUPRON DEPOTSYPHNGEKIT ¹ LUPRON DEPOT- PED KIT ¹ (1-month) LUPRON DEPOT-PED SYPHNGEKIT (3-month) TRIPYODUR VIAL	MC/DEL MC/DEL MC/DEL MC		FIRMAPON ² SUPPRELIN LA (IMPLANT) KIT TRELSTAR VANTAS ²	Use PA Form# 20420 1. Dosing limits apply, please refer to Dosage Consolidation List. 2. PA required to confirm FDA approved indication.	
ANTINEOPLASTIC AGENTS - TYROSINE KINASE INHIBITORS				MC MC/DEL MC		SPRYCEL ¹ TYKERB ² GLEEVEC ¹	Use PA Form# 20420 1. Verification of diagnosis is required. 2. PA required to confirm FDA approved indication and to monitor for potential drug-drug interactions.	
ANTINEOPLASTICS- MISCELLANEOUS	MC MC/DEL MC/DEL		AMIFOSTINE MERCAPTOPYRINE OXALIPLATIN	MC MC/DEL MC/DEL MC MC/DEL MC/DEL		DOCFREZ ELOXATIN ETHYOL LEUPROLIDE PURINETHOL ZOLINZA	Use PA Form# 20420	
ANTINEOPLASTICS- MONOCLONAL ANTIBODIES	MC/DEL		TRAZIMERA	MC/DEL MC/DEL MC MC/DEL MC MC MC/DEL		ENHERTU HERCEPTIN HERCESSI HERZUMA KANJINTI OGIVRI ONTRUZANT	Use PA Form# 20420	
CANCER								
CANCER	MC MC/DEL MC MC MC/DEL MC MC/DEL MC		ALIMTA ANASTROZOLE TABS ERBITUX IMATINIB MESYLATE LETOZOLE RUXIENGE VIDAZA ZIRABEV	MC MC MC MC/DEL MC MC MC/DEL MC MC MC MC/DEL MC/DEL MC MC MC MC/DEL MC/DEL MC MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC/DEL		ABECMA AKEEGA ALECENSA ALIQOPA ³ ALUNBRIG ¹ ALYMSYS ARIMIDEX AUCATZYL AUGTYRO AVMAPKI-FAKZYNJA AYVAKIT AVASTIN BALVERSA BAVENCIO ^{1,8} BENDEKA ³ BESPONSAS ³ BESREMI ¹ BIZENGRI BLENREP BOSULIF BRAFTOVI ¹ BREYANZI BRUKINSA CABOMETYX ³ CAMCEVI CALQUENCE ³ COMETRIQ ^{3,4,5} COTELLIC COPIKTRA DANZITEN DARZALEX ³ DATROWAY DAURISMO	Use PA Form# 20420 1. PA required to confirm appropriate diagnosis and testing. 2. Avoid CYP3A drug interaction. 3. Clinical PA required for appropriate diagnosis. 4. Re-approval will require documentation of response without disease progression and tolerance to treatment. 5. Dosing limits apply, see Dosage Consolidation List. 6. Max daily dose of 300mg. 7. Monitor liver enzymes periodically and stop treatment upon Grade 3 or higher elevation of liver enzymes approved indication. 8. For patients ≥ 12 years of age. 9. For the treatment of patients up to 25 years of age with B-cell acute lymphoblastic leukemia (ALL) that is refractory or in second or later relapse.	All non-preferred: A clinical PA is required to confirm appropriate clinical indication for the individual drug request. Specific to each drug all age, clinical testing requirements, previous step therapies, adjunctive drug therapy requirements, and response without disease progression will be also be evaluated for clinical appropriateness. The standard for the appropriate indication will include the FDA label as well as current NCCN guidelines. Scemblis is for the treatment of adult patients with: Philadelphia chromosome-positive chronic myeloid leukemia (Ph+ CML) in chronic phase (CP), previously treated with two or more tyrosine kinase inhibitors (TKIs).

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
				MC/DEL		ELREXFIO		
				MC/DEL		EMPLICITI(IV) ⁸		
				MC		EMRELIS		
				MC		EPKINLY		
				MC/DEL		ERLEADA		
				MC/DEL		ERIVEDGE		
				MC		EXKIVITY		
				MC		FARYDAK		
				MC/DEL		FEMARA		
				MC		FOLOTYN		
				MC		FOTIVDA		
				MC		FRUZAQLA		
				MC		GAVRETO		
				MC/DEL		GILOTRIF ^{4,5}		
				MC		GOMEKLI		
				MC		GRAFAPEX		
				MC/DEL		IBRANCE		
				MC		ICLUSIG ³		
				MC/DEL		IDHIFA ³		
				MC		IMBRUVICA		
				MC		IMDELLTRA		
				MC/DEL		IMFINZI		
				MC/DEL		IMJUDO		
				MC		IMKELDI		
				MC		IMLYGIC		
				MC/DEL		INLYTA		
				MC/DEL		INREBIC		
				MC		INQOVI		
				MC		ITOVEBI		
				MC		IWILFIN		
				MC		JAKAFI		
				MC		JAYPIRCA ^{1,2}		
				MC		JEMPERLI		
				MC/DEL		KEYTRUDA ¹		
				MC		KIMMTRAK		
				MC		KISQALI ¹		
				MC/DEL		KOSELUGO		
				MC		KRAZATI ³		
				MC		KYMRIAH ^{3,9}		
				MC		KYPROLIS ¹		
				MC		LARTRUVO ¹		
				MC		LAZCLUZE		
				MC		LENVIMA		
				MC/DEL		LIBTAYO ¹		
				MC		LONSURF		
				MC/DEL		LORBRENA		
				MC		LOQTORZI		
				MC		LUMAKRAS		
				MC/DEL		LUMOXITI ¹		
				MC		LUNSUMIO ¹		
				MC		LYNOZYFIC		
				MC		LYNPARZA ¹		
				MC		LYTGOBI		
				MC		NEXAVAR ¹		
				MC		NERLYNX ³		
				MC		NINLARO(PO)		
				MC/DEL		NUBEQA		
				MC		MARGENZA		
				MC/DEL		MEKINIST ^{3,4}		
				MC/DEL		MEKTOVI ¹		

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
				MC		MONJUVI		
				MC/DEL		MYLOTARG ³		
				MC/DEL		MVASI		
				MC		ODOMZO ^{1,2,5}		
				MC		OGSIVEO		
				MC		OJEMDA		
				MC		OJJAARA		
				MC		OMISIRGE		
				MC		ONUREG		
				MC/DEL		OPDIVO ³		
				MC		OPDIVO QVANTIG		
				MC		OPDUALAG		
				MC		ORGOVYX		
				MC		ORSERDU ^{2,3}		
				MC		PADCEV		
				MC		PEMAZYRE		
				MC		PEPAXTO		
				MC		PHESGO		
				MC/DEL		PIQRAY		
				MC		POLIVY		
				MC		POMALYST		
				MC		PORTRAZZA ³		
				MC		QINLOCK		
				MC		RETEVMO		
						REVUFORJ		
						ROMVIMZA		
				MC		REZLIDHIA		
				MC/DEL		ROZLYTREK		
				MC		RUBRACA		
				MC		RITUXAN		
				MC		RYBREVANT		
				MC		RYDAPT		
				MC		RYLAZE		
				MC		RYTELO		
				MC/DEL		SARCLISA		
				MC		SCEMBLIX ¹		
				MC/DEL		STIVARGA		
				MC/DEL		SUTENT ^{1,2}		
				MC/DEL		SYLATRON		
				MC		TABRECTA		
				MC		TALVEY		
				MC/DEL		TAFINLAR ^{3,4,5,6}		
				MC		TAZVERIK		
				MC/DEL		TALZENNA ¹		
				MC/DEL		TAGRISSO		
				MC		TECARTUS		
				MC		TECELRA		
				MC		TECENTRIQ ¹		
				MC		TECENTRIQ HYBREZA		
				MC		TEPMETKO		
				MC		TEVIMBRA		
				MC/DEL		TIBSOVO ¹		
				MC		TIVDAK		
				MC		TRODELVY		
				MC		TRUSELTIQ		
				MC/DEL		TRUXIMA		
				MC/DEL		TRUQAP		
				MC		TUKYSA		
				MC		UKONIQ		
				MC/DEL		VANFLYTA		

CATEGORY	Coverage Indicator	Step Order	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
				MC MC MC MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC MC MC MC/DEL MC MC MC		VEGZELMA VENCLEXTA ³ VERZENIO ³ VITRAKVI VIZIMPRO ¹ VONJO VORANIGO VYLOY WELIREG XALKORI XPOVIO XOSPATA XTANDI YERVOY YESCARTA ³ ZALTRAP ZEJULA ¹ ZELBORAF ZEPZELCA ZIIHERA ZYDELIG ZYKADIA ZYNLONTA ZYNYZ ¹ ZYTIGA		
IMMUNOSUPPRESSANTS								
IMMUNOSUPPRESSANTS	MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		CYCLOSPORINE MODIFIED GENGRAF CAPS MYCOPHENOLATE MYFORTIC NEORAL SOL SANDIMMUNE TACROLIMUS CAPS	MC/DEL MC/DEL MC/DEL MC MC MC/DEL MC MC MC MC/DEL		CELLCEPT CYCLOSPORINE CAPS CYCLOSPORINE SOL. MODIFIED ENVARUSUS XR MYHIBBIN ² NEORAL CAP PROGRAF CAPS REZUROCK ¹ ZORTRESS	Use PA Form# 20420 1. For the treatment of adult and pediatric patients 12 years and older with chronic graft-versus-host disease (chronic GVHD) after failure of at least 2 prior lines of systemic therapy. 2.Clinical PA is required.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Myhibbin: For the prophylaxis of organ rejection, in adult and pediatric recipients 3 months of age and older of allogeneic kidney, heart, or liver transplants, in combination with other immunosuppressants. DDI: All preferred immunosuppressants will require clinical PA for patients over 60 that are currently on fluoroquinolone therapy. DDI: Cyclosporine will now be non-preferred and require prior authorization if it is currently being used in combination with either Lipitor (doses greater than 20mg/day), crestor, or lovastatin (doses greater than 20mg). DDI: Cyclosporine will require prior authorization when used with livalo.
IMMUNOSUPPRESSANTS- Misc.				MC		HYFTOR ^{1,2}	Use PA Form# 20420 1. For the treatment of patients ≥ 6 years of age. 2. Clinical PA required for appropriate diagnosis and clinical parameters.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
PURINE ANALOG								
PURINE ANALOG	MC MC/DEL		AZASAN TABS AZATHIOPRINE TABS	MC/DEL		IMURAN TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
K REMOVING RESINS								
K REMOVING RESINS	MC/DEL MC/DEL		LOKELMA SODIUM POLYSTYRENE SULFON	MC/DEL MC/DEL MC		SPS SUSP SPS 30GM/120ML ENEMA SUSP VELTASSA	Use PA Form# 20420	
New drugs are initially non-preferred until reviewed by the DUR Committee and the State. According to State policy, any drug requiring specific diagnosis still requires the specific diagnosis unless otherwise noted within this document.								

Last update **09/2025**

PDL DOSAGE CONSOLIDATION LIST

Tabs/Caps/Patches: Quantities in units

Shaded areas are non-preferred agents - Quantities of these

Sprays/Inhalers/Nebulizers: Quantities in GM, ML, OR MCG

non-preferred agents are available up the limit only with

Injectibles: Quantities in ML

prior authorization

Drug Name	Strength	Limit/Day	Limit/Days
ABILIFY SOLUTION	1MG/ML	30ML	1020/34
ACCUPRIL	5MG	1	35/35
ACCUPRIL	10MG	1	35/35
ACCUPRIL	20MG	1	35/35
ACEON	2MG	1	35/35
ACEON	4MG	1	35/35
ACTONEL	5MG	1	35/35
ACTONEL	35MG	1/WK	5/35
ACTOS	All Strengths	1	35/35
ADDERALL XR	5MG	3	90/30
ADDERALL XR	10MG	3	90/30
ADDERALL XR	15MG	3	90/30
ADDERALL XR	20MG	2	60/30
ADDERALL XR	30MG	1	35/35
ADEMPAS	All Strengths	1	35/35
ADVAIR DISKUS	All Strengths	2	60/30
ADVAIR HFA	All Strengths	4	120/30
ADZENYS XR	All Strengths	1	30/30
AEROBID	250MCG	8 INHALATIONS	21/35
AEROBID-M	250MCG	8 INHALATIONS	21/35
ALAVERT-NON DROW	TAB	1	96/96
ALENDRONATE	All Strengths	1/WK	35/35
ALTABAX	5GM		1 TUBE/30
ALTABAX	15GM		1 TUBE/30
ALTABAX	30GM		1 TUBE/30
ALTACE	1.25MG	1	35/35
ALTACE	2.5MG	1	35/35
ALTACE	5MG	1	35/35
AMARYL	1MG	1	35/35
AMARYL	2MG	1	35/35
AMBIEN	5MG		12/34
AMBIEN	10MG		12/34
AMBIEN CR	6.25MG		12/34
AMBIEN CR	12.5MG		12/34
AMERGE (Step 8)	1MG		12/30
AMERGE (Step 8)	2.5MG	2.5MG	12/30
AMLODIPINE	2.5MG	1.5	53/35 DAYS
AMLODIPINE	5MG	1.5	53/35 DAYS
AMMONIUM LACTATE CREA	12%		1 TUBE/10
AMMONIUM LACTATE LOTN	12%		1TUBE/8
AMPHETAMINE/DEXTROAMPHET ER	5MG	3	90/30
AMPHETAMINE/DEXTROAMPHET ER	10MG	3	90/30
AMPHETAMINE/DEXTROAMPHET ER	15MG	3	90/30
AMPHETAMINE/DEXTROAMPHET ER	20MG	2	60/30
AMPHETAMINE/DEXTROAMPHET ER	30MG	1	90/90
AMPHETAMINE SALT	5,10,15MG	3	105/35
AMPHETAMINE SALT	20MG	2	70/35
AMPHETAMINE SALT	30MG	1	35/35
ANDRODERM	2.5MG	2	60/30
ANDRODERM	5MG	1	30/30
ARAVA	10MG	1	35/35
ARCAPTA	75MCG	1 INHALATION	35/35
ARICEPT	5MG	1	35/35
ARICEPT	10MG	1	35/35
ARIPIRAZOLE	2MG	2	180/90
ARIPIRAZOLE	5MG	2	180/90
ARIPIRAZOLE	10MG	2	180/90
ARIPIRAZOLE	15MG	2	180/90
ARIPIRAZOLE	20MG	1.5	135/90
ARIPIRAZOLE	30MG	1	90/90
ARIXTRA INJECTION	2.5MG/0.5ML		7/30
ARIXTRA INJECTION	5MG/0.4ML		7/30
ARIXTRA INJECTION	7.5MG/0.6ML		7/30
ARIXTRA INJECTION	10MG/0.8ML		7/30
ARMONAIR	All Strengths	1 INHALATION	60U/30
ASMANEX 30 UNITS	220MCG	1 INHALATION	30U/30
ASMANEX 60 UNITS	220MCG	2 INHALATIONS	60U/30
ASMANEX 120 UNITS	220MCG	4 INHALATIONS	120U/30
ATACAND	4MG	1.5	53/35

Drug Name	Strength	Limit/Day	Limit/Days
ATROVENT HFA	17MCG	12 INHALATIONS	25.8/34
ATROVENT 30ML	0.03%	12 SPRAYS	30/30
ATROVENT 15ML	0.06%	16 SPRAYS	45/30
AVANDIA	2MG	1.5	53/35
AVANDIA	4MG	1	35/35
AVAPRO	75MG	1.5	53/35
AVAPRO	150MG	1	35/35
AXERT (Step 8)	6.25MG		12/30
AXERT (Step 8)	12.5MG		12/30
AZELEX	20%		1 TUBE/18
AZILECT	All Strengths	1	35/35
BACTROBAN CREAM			1 TUBE/30
BECONASE AQ	42MCG	8 INHALATIONS	50/30
BENICAR-HCT	All Strengths	1	30/30
BENAZEPRIL	5MG	1	35/35
BENAZEPRIL	10MG	1.5	53/35
BENAZEPRIL	20MG	1	35/35
BENAZEP/HCTZ	5-6.25	1	35/35
BENAZEP/HCTZ	10/12.5	1	35/35
BEVESPI AERO		4 INHALATIONS	120/30
BONIVA	2.5MG	1	35/35
BOTOX (ADULTS)	100U/ML	1 session/90 days	600U/90
BOTOX (CHILDREN>12)	100U/ML	1 session/90 days	400U/90
BREO ELLIPTA	100/25MCG	1 INHALATIONS	60/60
BRILINTA	All Strengths	2	70/35
BRINTELLIX	All Strengths	1	35/35
BUTRANS		1 patch/WK	4/28
BYETTA	5mcg inj	0.04ML	1.2ML/30
BYETTA	10mcg inj	0.08ML	2.4ML/30
CALAN SR	120MG	1	35/35
CALAN SR	180MG	2	70/35
CALAN SR	240MG	2	70/35
CARDIZEM CD	120MG/24	1	35/35
CARDIZEM CD	180MG/24	1	35/35
CARDIZEM CD	240MG/24	1	35/35
CARDIZEM CD	300MG/24	1	35/35
CARDIZEM CD	360MG/24	1	35/35
CARDIZEM LA	120MG/24	1	35/35
CARDIZEM LA	180MG/24	1	35/35
CARDIZEM LA	240MG/24	1	35/35
CARDIZEM LA	300MG/24	1	35/35
CARDIZEM LA	360MG/24	1	35/35
CARDURA	1MG	1	35/35
CARDURA	2MG	1.5	53/35
CARDURA	4MG	1.5	53/35
CARTIA XT	120MG	1	90/90
CARTIA XT	180MG	1	90/90
CARTIA XT	240MG	1	90/90
CARTIA XT	300MG	1	90/90
CATAPRES-TTS1	0.1 MG/24HR		5/35
CATAPRES- TTS2	0.2 MG/24HR		5/35
CATAPRES- TTS3	0.3 MG/24HR		5/35
CEFIXIME	400MG	2	2/7
CELEBREX	100MG	1	35/35
CELEBREX	200MG	2	70/35
CELEBREX	400MG	1	35/35
CELEXA	20mg	0.5	17/34
CELEXA	40mg	1	51/34
CITALOPRAM	10MG	2	180/90
CITALOPRAM	20MG	2	180/90
CITALOPRAM	40MG	1	90/90
CLARINEX	REDI TAB	1	35/35
CLEOCIN-T		1 PACKAGE	1/30
CLINDAMYCIN PHOSPHATE		1 PACKAGE	1/30
COMBIVENT	103-18MCG	12 INHALATIONS	30/35
Drug Name	Strength	Limit/Day	Limit/Days
EFFEXOR XR	37.5MG	1	35/35
EFFEXOR XR	75MG	1	35/35
EMSAM	All Strengths	1	34/34

ATACAND	8MG	1.5	53/35
ATACAND	16MG	1	35/35
ATRIPLA	600MG	1	35/35
ATOMOXETINE	All Strengths	1	90/90
COMETRIQ	80MG	1	35/35
COMETRIQ	20MG	3	105/35
CONCERTA	18MG	1	30/30
CONCERTA	27MG	1	30/30
CONCERTA	36MG	2	60/30
COPAXONE INJ	20MG		1/32
COPAXONE KIT	20MG/ML		1/30
COREG CR	All Strengths	1	34/34
COSENTYX	150MG	1	1/30
CRESTOR	5MG	1	35/35
CRESTOR	10MG	1	35/35
CRESTOR	20MG	1	35/35
CRESTOR	40MG	1	35/35
CYMBALTA	All Strengths	1	35/35
DALMANE	15MG		10/30
DALMANE	30MG		10/30
DAYPRO	600MG	2	70/35
DAYTRANA	10mg/9hr (27.5mg)	1	34/34
DAYTRANA	15mg/9hr (41.3mg)	1	34/34
DAYTRANA	20mg/9hr (55.0mg)	1	34/34
DAYTRANA	30mg/9hr (82.5mg)	1	34/34
DDAVP	5ML		15/34
DENAVIR CREAM			2gm/30
DEPO-PROVERA	150MG/ML		1/90
DEPO-PROVERA	400MG/ML		2.5/90
DEPO-TESTOSTERONE	200MG/ML		20/90
DESMOPRESSIN	0.1MG	12	420/35
DESMOPRESSIN	0.2MG	6	210/35
DESONIDE	0.05%		2 TUBES/30
DESOWEN	0.05%		2 TUBES/30
DETROL LA	2MG	1	35/35
DEXEDRINE	All Strengths	3	90/30
DEXILANT	All Strengths	1	35/35
DEXTROAMPHETAMINE	All Strengths	3	90/30
DICLOFENAC 1% GEL	1% GEL		2 TUBES/30
DIFLUCAN	150MG		1/7
DILACOR XR	240MG/24	1	35/35
DILACOR XR	120MG/24	1	35/35
DILACOR XR	180MG/24	1	35/35
DILTIA - XT	120MG/24	1	90/90
DILTIA - XT	180MG	1	90/90
DILTIA - XT	240MG/24	1	90/90
DILTIAZEM CAP ER	120MG	1	90/90
DILTIAZEM CAP XR	120MG	1	90/90
DILTIAZEM CAP	120MG/24	1	90/90
DILTIAZEM CAP	180MG/24	1	90/90
DILTIAZEM CAP ER	240MG	1	90/90
DILTIAZEM CAP XR	240MG	1	90/90
DILTIAZEM XR CAP	240MG/24	1	90/90
DILTIAZEM CAP	240MG/24	1	90/90
DILTIAZEM CAP	300MG/24	1	90/90
DILTIAZEM CAP	360MG/24	1	90/90
DIOVAN	80MG	1	35/35
DIOVAN - HCT	80 - 12.5	1	35/35
DITROPAN XL	5MG	1	35/35
DITROPAN XL	10MG	2	70/35
DORAL	7.5MG		10/30
DOXAZOSIN	1MG	1	90/90
DOXAZOSIN	2MG	1.5	135/90
DOXAZOSIN	4MG	1.5	135/90
DRYSOL SOL	20%		1 BOTTLE/30DAYS
DURAGESIC PATCHES	12.5MCG/HR		11/33
DURAGESIC PATCHES	25MCG/HR		11/33
DURAGESIC PATCHES	50MCG/HR		11/33
DURAGESIC PATCHES	75MCG/HR		11/33
DURAGESIC PATCHES	100MCG/HR		22/33
DULOXETINE	20MG	3	270/90
DULOXETINE	30MG	3	270/90
DULOXETINE	60MG	2	180/90
EDEX	All Strengths		1/30

ENALAPRIL	2.5	1	90/90
ENALAPRIL	5MG	1.5	135/90
ENALAPRIL	10MG	1.5	135/90
ENALAPR/HCTZ	5-12.5	1	90/90
ENBREL	25MG/ML		8/28
ENBREL SURECLICK			8/28
ESTAZOLAM	1MG		10/30
ESTAZOLAM	2MG		10/30
ESTRING MIS	2MG		1/90
EVENITY		12 DOSES/LIFETIME	12 DOSES/LIFETIME
EVOTAZ	All Strengths	1	30/30
FELODIPINE	2.5MG	1	90/90
FELODIPINE	5MG	1.5	135/90
FENTANYL	25MCG/HR		11/33
FENTANYL	50MCG/HR		11/33
FENTANYL	75MCG/HR		11/33
FENTANYL	100MCG/HR		22/33
FETZIMA	All Strengths	1	35/35
FINASTERIDE	5MG	1	90/90
FLONASE	50MCG	4 SPRAYS	32/34
FLOVENT HFA 44MCG	44MCG	4 INHALATIONS	10.6/30
FLOVENT HFA 110MCG	110MCG	4 INHALATIONS	12/30
FLOVENT HFA 220MCG	220MCG	8 INHALATIONS	24/30
FLOVENT DISKUS	50MCG, 100MCG	4 INHALATIONS	60/30
FLOVENT DISKUS	250MCG	3 INHALATIONS	120/30
FLUCONAZOLE	150MG		1/7
FLUNISOLIDE SOLN	0.025%	16 SPRAYS	75/30
FLUOXETINE CAP	40MG	2	180/90
FLUOXETINE CAP	20MG	4	360/90
FLUOXETINE CAP	10MG	3	270/90
FLURAZEPAM	15MG		10/30
FLURAZEPAM	30MG		10/30
FLUTICASONE SPR		4 SPRAYS	48/90
FLUVOXAMINE	25MG	3	270/90
FLUVOXAMINE	50MG	3	270/90
FOCALIN	All Strengths	3	105/35
FOCALIN XR	All Strengths	1	35/35
FORFIVO XL	All Strengths	1	35/35
FOSAMAX	5MG	1	35/35
FOSAMAX	10MG	1	35/35
FOSAMAX	70MG	1/WK	5/35
FOSAMAX	40MG	2/WK	10/35
FOSINOPRIL	10MG	1.5	135/90
FOSINOPRIL	20MG	2	180/90
FRAGMIN INJ	10000U/ML	2ML	14/7
FRAGMIN INJ	2500U/.2ML	0.4ML	2.80/7
FRAGMIN INJ	25000U/ML	0.8ML	5.6/7
FRAGMIN INJ	5000U/.2ML	0.4ML	2.80/7
FRAGMIN INJ	7500U/.3ML	0.6ML	4.2/7
FROVA TAB (Step 8)	2.5MG		12/30
FULYZAQ	125MG	2	70/35
FUZEON	KIT	1	1/30
FYCOMPA	All Strengths	1	35/35
GABAPENTIN	300MG	9	810/90
GABAPENTIN	400MG	9	810/90
GABAPENTIN	600MG	6	540/90
GABAPENTIN	800MG	4	360/90
GEODON	20MG	2	70/35
GEODON	40MG	2	70/35
GEODON	60MG	2	70/35
GEODON	80MG	2	70/35
GEODON	INJ	2	70/35
GILOTRIF	All Strengths	1	35/35
GLIMEPIRIDE	1MG	1	90/90
GLIMEPIRIDE	2MG	1	90/90
GLUCOSE TES STRP		12	420/35
GLUCAGEN INJ. HYPOKIT			2/30
GLYCOLAX*	255GM		255GM/90
* Available for once daily dosing to members under the age of 18 years			
Drug Name	Strength	Limit/Day	Limit/Days
LUNESTA	2MG		12/34
LUNESTA	3MG		12/34
LUPRON DEPOT INJ	11.25MG	KIT	1/90

Drug Name	Strength	Limit/Day	Limit/Days
ILARIS			2/28
HALCION	0.125MG		10/35
HALCION	0.25		10/35
HUMIRA	40mg/0.8ml		4/28
HYDROXYZINE TAB	All Strengths	3	270/90
HYTRIN	1MG	1	35/35
HYTRIN	5MG	1	35/35
HYZAAR	50-12.5	1	35/35
IMDUR	30MG	1.5	53/35
IMDUR	60MG	1.5	53/35
IMITREX (step 8)	25MG		12/30
IMITREX (step 8)	50MG		12/30
IMITREX (step 8)	100MG		12/30
IMITREX VIAL	All Strengths		6 boxes/30
IMITREX CARTRIDGE	All Strengths		12/30
IMITREX NASAL SPRAY	All Strengths		12/30
IMITREX PEN INJCTR	All Strengths		12/30
IMIQUIMOD	5%		12/28
IMIQUIMOD	5%		12/28
INTAL	800MCG	8 INHALATIONS	28.4/34
INVOKANA	All Strengths	1	35/35
IPRATROPIUM 30ML	0.03%	12 SPRAYS	90/90
IPRATROPIUM 15ML	0.06%	16 SPRAYS	135/90
ISOPTIN SR	180MG	2	70/35
IRBESARTAN	All Strengths	1	90/90
ISOPTIN SR	240MG	2	70/35
ISOSORBIDE MONO	30MG	2	180/90
ISOSORBIDE MONO	60 MG	1.5	135/90
JANUMET	All Strengths	2	70/35
JANUVIA	All Strengths	1	35/35
JUVISYNC	All Strengths	1	35/35
KETOPROFEN	100MG	2	180/90
KETOPROFEN	200MG	1	90/90
KETOROLAC	10MG	4.8	24/30
KHEDEZLA	All Strengths	1	35/35
LAC-HYDRIN CREAM	12%		1TUBE/30
LAMICTAL	25MG	6	210/35
LAMICTAL	25MG CHW	6	210/35
LAMICTAL	100MG	2	70/35
LAMISIL	250MG	1	35/35
LAMOTRIGINE	25MG	6	540/90
LAMOTRIGINE	100MG	2	180/90
LANSOPRAZOLE CAPS	All Strengths	2	180/90
LATUDA	All Strengths	1	17/34
LESCOL	20MG	1	35/35
LEVAQUIN	250MG	1	35/35
LEXAPRO	5MG	0.5	15/30
LIPITOR	10MG	1	35/35
LIPITOR	20MG	1	35/35
LIPITOR	40MG	1.5	53/35
LISINOP/HCTZ	10/12.5MG	1	90/90
LINEZOLID	600mg		28/60
LINZESS	All Strengths	1	35/35
LOSARTAN	All Strengths	1	90/90
LOSARTAN- HCT	All Strengths	1	90/90
LOTENSIN	5MG	1	35/35
LOTENSIN	10MG	1.5	35/35
LOTENSIN	20MG	1	53/35
LOTENSIN - HCT	5 - 6.25	1	35/35
LOTENSIN - HCT	10 - 12.5	1	35/35
LOVASTATIN	10MG	1.5	135/90
LOVASTATIN	20MG	1.5	135/90
LOVENOX INJ	30MG/.3ML	0.6	14 injections/7
LOVENOX INJ	40MG/.4ML	0.8	14 injections/7
LOVENOX INJ	60MG/.6ML	1.2	14 injections/7
LOVENOX INJ	80MG/.8ML	1.6	14 injections/7
LOVENOX INJ	100MG/ML	2	14 injections/7
LOVENOX INJ	120MG/.8ML	1.6	14 injections/7
LOVENOX INJ	150MG/ML	2	14 injections/7
LUNESTA	1MG		12/34
Drug Name	Strength	Limit/Day	Limit/Days
NIFEDIPINE ER	90MG	1	90/90
NIFEDIPINE ER,CR	30MG	1	90/90

LUPRON DEPOT INJ	22.5	KIT	1/90
LUPRON DEPOT INJ	30MG		1/90
LUPRON DEPOT INJ	30MG	KIT	1/90
LYRICA	25,50,75MG	3	102/35
LYRICA	100,150,200MG	3	102/35
LYRICA	225,300MG	2	70/35
MAVIK	1MG	1	35/35
MAVIK	2MG	1	35/35
MAXAIR AUTO	200MCG	12 INHALATIONS	14/30
MAXALT (step 8)	5MG		12/30
MAXALT (step 8)	10MG		12/30
MAXALT MLT (step 1)	5MG		12/30
MAXALT MLT (step 1)	10MG		12/30
MEDROXYPR AC	150MG/ML		1/90
MELOXICAM TABS	All Strengths	1	90/90
METADATE ER	10,20MG	3	90/30
METFORMIN ER	500MG	4	360/90
METHYLIN	All Strengths	3	90/30
METHYLPHENIDATE ER	36mg	2	180/90
METHYLPHENIDATE	All Strengths	3	90/30
METROCREAM		1 PACKAGE	1/30
METROGEL		1 PACKAGE	1/30
METROLOTION		1 PACKAGE	1/30
METRONIDAZOLE CREAM		1 PACKAGE	1/30
METRONIDAZOLE GEL		1 PACKAGE	1/30
METRONIDAZOLE LOTION		1 PACKAGE	1/30
MEVACOR	10MG	1.5	53/35
MEVACOR	20MG	1.5	53/35
MIACALCIN		3.75ml	1 bottle/34
MICARDIS	All Strengths	1	30/30
MICARDIS-HCT	All Strengths	1	30/30
MIGRANAL NASAL SPRAY	All Strengths		12/30
MIRALAX	255G	8.5G	1 bottle/30
MIRALAX	17G/PACKET	0.5 packet	15 packets/30
MIRTAZAPINE	15mg	3	270/90
MOBIC	7.5 MG	1	35/35
MOBIC	15MG	1	35/35
MOEXIPRIL	7.5	1.5	135/90
MONOPRIL	10MG	1.5	53/35
MONOPRIL	20MG	2	70/35
MUPIROCIN			1 TUBE/30
NABUMETONE	500MG	2	180/90
NABUMETONE	750MG	2	180/90
NARATRIPTAN			12/30
NASACORT AERS	55 MCG	4 SPRAYS	9.3/25
NASONEX	50MCG	4 SPRAYS	17/30
NATROBA		120ML	1 bottle/30
NAYZILAM	All Strengths		5/30
NEUPOGEN INJ	300MCG/ML		10/30
NEUPOGEN INJ	480MCG/1.6		16/30
NEUPOGEN INJ	300MCG/.5ML		5/30
NEUPOGEN INJ	480MCG/.8ML		8/30
NEURONTIN	300MG	9	315/35
NEURONTIN	600MG	9	315/35
NEXIUM	20MG	1	35/35
NEXIUM	40MG	2	70/35
NEXIUM SUS	All Strengths	1	30/30
NIFEDIPINE CR	90MG	1	90/90
NIFEDIPINE ER	60MG	1	90/90
NIFEDIPINE ER	30MG	1	90/90
NIFEDIPINE ER	60MG	1	90/90
Drug Name	Strength	Limit/Day	Limit/Days
RELPAX	All Strengths		12/30
REMODULIN	All Strengths		1 MDV/30
RESTORIL	7.5MG		10/30
RESTORIL	15MG		10/30
RESTORIL	30MG		10/30
RETIN-A		1 TUBE	1 TUBE/30
REVLIMID	All Strengths	1	35/35
REYVOW	All Strengths		4/30
RHINOCORT AQ	32MCG	8 SPRAYS	18/30
REFRESH PLUS		15 ML	1 bottle/30

NORVASC	2.5MG	1.5	53/35 DAYS
NORVASC	5MG	1.5	53/35 DAYS
NURTEC ODT	All Strengths		8/30
NUVARING		1/MO	1/28
ODOMZO	200mg	1	30/30
OLMESARTAN	All Strengths	1	90/90
OLANZAPINE	2.5MG	3	270/90
OLANZAPINE	5MG	3	270/90
OLANZAPINE	7.5MG	3	270/90
OLANZAPINE	10MG	3	270/90
OLANZAPINE	15MH	2	180/90
OLANZAPINE	20MG	1.5	135/90
OLANZAPINE ODT	All Strengths	1	90/90
OMEPRAZOLE	10MG	2	180/90
OMEPRAZOLE	20MG	2	180/90
OMEPRAZOLE	40MG	2	180/90
OMNARIS	50MCG	4 sprays	12.5/30
ONGLYZA	All Strengths	1	35/35
OPSUMIT	All Strengths	1	35/35
ORUVAIL	100MG	2	70/35
ORUVAIL	200MG	1	35/35
OXAPROZIN	600MG	2	180/90
OXYCODONE ER	10,20,40MG	2	70/35
OXYCODONE ER	80MG	4	140/35
OXYCONTIN**	10,20,30,40MG	2	70/35
OXYCONTIN**	80MG	4	140/35
PANTOPRAZOLE	All Strengths	2	180/90
PAROXETINE	10MG	2	180/90
PAROXETINE	20MG	2	180/90
PAXIL	10MG	1.5	53/35
PAXIL	20MG	1	35/35
PEGASYS KIT		KIT	1/28
PLAN B			2/15 or 4/30
PLENDIL	2.5MG	1	35/35
PLENDIL	5MG	1.5	53/35
PRAVACHOL	10MG	1	35/35
PRAVACHOL	20MG	1	35/35
PRAVACHOL	40MG	1	35/35
PRAVACHOL	80MG	1	35/35
PRAVASTATIN	10MG	1	35/35
PRAVASTATIN	20MG	1	35/35
PRAVASTATIN	40MG	2	180/90
PRAVASTATIN	80MG	1	35/35
PREVPAC MIS	500MG-30MG		14/30
PRILOSEC OTC	20MG	2	168/84
PRINIVIL	2.5MG	1	35/35
PRINIVIL	5MG	1	35/35
PRINIVIL	10MG	1.5	53/35
PRINIVIL	20MG	1.5	53/35
PRINZIDE	10-12.5	1	35/35
PROAIR HFA	90mcg	12 INHALATIONS	17/34
PROTONIX	20MG	2	70/35
PROTONIX	40MG	2	70/35
PROZAC	10MG	1.5	53/35
PULMICORT	200MCG	8 INHALATIONS	1/25
PULMICORT FLEX	All Strengths	8 Inhalations	2/30
QUETIAPINE	25MG	3	270/90
QUETIAPINE	50MG	3	270/90
QUETIAPINE	100MG	3	270/90
QUETIAPINE	200MG	3	270/90
QUINAPRIL	5MG	1	90/90
QUINAPRIL	10MG	1	90/90
QUINAPRIL	20MG	1	90/90
QVAR AERS	All Strengths	8 Inhalations	14.6/25
RANITIDINE SYRUP***	15MG/ML	20ML	700ML/35
RELAFEN	500MG	2	70/35
RELAFEN	750MG	2	70/35
REMERON	15MG	1.5	53/35
Drug Name	Strength	Limit/Day	Limit/Days
SULAR	10MG	1.5	53/35
SULAR	20MG	1	35/35
SUMATRIPTAN PEN INJ	All Strengths		12/30
SUMATRIPTAN NASAL SPRAY	All Strengths		12/30
SUMATRIPTAN SYRINGE	All Strengths		12/30

REFRESH PLUS		30 ML	2 bottles/30
REFRESH TEARS		15 ML	1 bottle/30
REFRESH TEARS		30 ML	2 bottles/30
RESCULA			2 bottles/35
REYATAZ	All Strengths	1	35/35
RISPERDAL	0.5MG	1.5	53/35
RISPERDAL	0.25MG	1.5	53/35
RISPERDAL	1MG	1.5	53/35
RISPERDAL	2MG	1.5	53/35
RISPERDAL	3MG	2	70/35
RISPERDAL	4MG	2	70/35
RISPERDAL INJ	25MG		2/28
RISPERDAL INJ	37.5		2/28
RISPERDAL INJ	50MG		2/28
RISPERDAL M-TAB	0.5MG	1.5	53/35
RISPERDAL M-TAB	1MG	1.5	53/35
RISPERDAL M-TAB	2MG	4	140/35
RISPERDAL SOL.	1MG/ML	8ML	280/35
RISPERIDONE	0.5MG	3	270/90
RISPERIDONE	0.25MG	3	270/90
RISPERIDONE	1MG	3	270/90
RISPERIDONE	2MG	3	270/90
RISPERIDONE	3MG	2	180/90
RISPERIDONE	4MG	2	180/90
RISPERIDONE SOL.	1MG/ML	8ML	280/35
RITALIN LA	All Strengths	1	35/35
RITALIN LA	30mg	2	70/35
SAVELLA	All Strengths	2	70/35
SEREVENT DISKUS	50MCG	2 INHALATIONS	60/30
SEROQUEL	100MG		45/30
SEROQUEL XR	150MG	1	35/35
SEROQUEL XR	200MG	1	35/35
SEROQUEL XR	300MG	2	70/35
SEROQUEL XR	400MG	2	70/35
SERTRALINE	25MG	3	270/90
SERTRALINE	50MG	3	270/90
SERTRALINE	100MG	3	270/90
SIMVASTATIN	5MG	1	35/35
SIMVASTATIN	10MG	1.5	53/35
SIMVASTATIN	20MG	1.5	53/35
SIMVASTATIN	40MG	1.5	53/35
SIMVASTATIN	80MG	1	35/35
SINGULAIR	4MG	1	35/35
SINGULAIR	5MG	1	35/35
SINGULAIR	10MG	1	35/35
SONATA	5MG		12/34
SONATA	10MG		12/34
SPIRIVA	HANDIHLR	1 INHALTION	30/30
SPORANOX SOL	10MG/ML	10ML/ML	300cc/30
SPORANOX PULSEPAK	F		30/30
SPORANOX	100MG		30/30
STADOL INJ	1MG/ML		9/35
STADOL INJ	2MG/ML		9/35
STRATTERA	All Strengths	1	35/35
SUPRAX	400MG	1	1/7
Drug Name	Strength	Limit/Day	Limit/Days
XOPENEX HFA		12 INHALATIONS	2 INHALERS/34
XOPENEX NEB		12CC	408/34
ZALEPLON	All Strengths		30/30
ZECUITY	6.5		4/28
ZEMBRACE	All Strengths		3boxes/30
ZESTORETIC	10-12.5	1	35/35
ZESTRIL	2.5MG	1	35/35
ZESTRIL	5MG	1	35/35
ZESTRIL	10MG	1.5	53/35
ZESTRIL	20MG	1.5	53/35
ZETONNA	37MCG	2	60/30
ZIPRASIDONE	20MG	3	270/90
ZIPRASIDONE	40MG	3	270/90
ZOCOR	5MG	1	35/35
ZOCOR	10MG	1.5	53/35
ZOCOR	20MG	1.5	53/35
ZOCOR	40MG	1.5	53/35

SUMATRIPTAN TAB	All Strengths		12/30
SYNVISC INJ	8MG/ML		2/30
SYRINGES		10	1000/100
TAFINLAR	50MG	6	210/35
TAFINLAR	75MG	4	140/35
TAMIFLU CAPS	75MG		10/30
TAZTIA XT CAP	120MG/24	1	90/90
TAZTIA XT CAP	180MG/24	1	90/90
TAZTIA XT CAP	240MG/24	1	90/90
TAZTIA XT CAP	300MG/24	1	90/90
TAZTIA XT CAP	360MG/24	1	90/90
TELMISARTAN	All Strengths	1	90/90
TEMAZEPAM	7.5MG		10/30
TEMAZEPAM	15MG		10/30
TEMAZEPAM	30MG		10/30
TEQUIN	200MG	1	35/35
TERAZOSIN	1MG	1	90/90
TERAZOSIN	5MG	1	90/90
TERBINAFINE	250MG	1	35/35
TEST STRIPS	Blood Glucose	12	420/35
TIAZAC	120MG/24	1	35/35
TIAZAC	180MG/24	1	35/35
TIAZAC	240MG/24	1	35/35
TIAZAC	300MG/24	1	35/35
TIAZAC	360MG/24	1	35/35
TIAZAC	420MG/24	1	35/35
TILADE	1.75MG	8 INHALATIONS	48.6/35
TOPAMAX SPRINKLES	All Strengths	1	35/35
TOPROL XL	25MG	1.5	53/35
TOPROL XL	50MG	1.5	53/35
TRADJENTA	All Strengths	1	35/35
TRAMADOL	50MG	8	720/90
TRAMADOL/ APAP	37.5/325MG	8	720/90
TRETINOIN		1 TUBE	1 TUBE/30
TRELEGY ELLIPTA	All Strengths	1INHALATION	30U/30
TREXIMET	85/500	2.5	12/30
TRIAZOLAM	0.125MG		10/30
TRIAZOLAM	0.25MG		10/30
TROKENDI XR	25MG	1	35/35
TROKENDI XR	50MG	1	35/35
TROKENDI XR	100MG	1	35/35
TROKENDI XR	200MG	2	70/35
UBRELVY	All Strengths		10/30
ULTRAM	50MG	8	280/35
UNIVASC	7.5MG	1.5	53/35 DAYS
UTIBRON	7.5mcg/15.6mcg	2 INHALATIONS	60/30
VALTOCO	All Strengths		10/30
VALSARTAN-HCT	All Strengths	1	90/90
VASERETIC	5-12.5MG	1	35/35
VASOTEC	2.5MG	1	35/35
VASOTEC	5MG	1.5	53/35
VASOTEC	10MG	1.5	53/35
VENLAFAXINE TABS	25	3	270/90
VENLAFAXINE TABS	37.5	3	270/90
VENLAFAXINE TABS	100	3	270/90
VENLAFAXINE ER CAPS	37.5	3	270/90
VENLAFAXINE ER CAPS	75	3	270/90
VENLAFAXINE ER	150	2	180/90
VENTOLIN HFA	90MCG	12 INHALATIONS	36/34
VERAPAMIL ER, SR	120MG	1	90/90
VERAPAMIL ER, CR, SR	180MG	2	90/90
VERAPAMIL ER, CR, SR	240MG	2	90/90
VERELAN	180MG	1	35/35
VERELAN SR	120MG	1	35/35
VERELAN SR	180MG	1	35/35
VERELAN SR	240MG	2	70/35
VERAMYST	27.5MCG	4 sprays	10/30
VYEPTI	All Strengths		4/30
VYVANSE	All Strengths	1	35/35
VYVANSE CHEW	All Strengths	1	35/35

ZOFRAN*	4MG	3	90/30
ZOFRAN*	8MG	1.5	45/30
ZOFRAN*	24MG	0.5	15/30
ZOFRAN*	4MG/5ML	15ML	450/30
ZOLMITRIPTAN TAB	All Strengths		12/30
ZOLOFT	25MG	0.5	18/35
ZOLOFT	50MG	0.5	18/35
ZOLOFT	100MG	3	105/35
ZOLPIDEM (step 1)	5MG		30/30
ZOLPIDEM (step 1)	10MG		30/30
ZOMIG (Step 8)	5MG		12/30
ZTLIDO	All Strengths	3	90/30
ZYPREXA	2.5MG	1.5	53/35
ZYPREXA	5MG	1	35/35
ZYPREXA	7.5MG	1	35/35
ZYPREXA	10MG	1	35/35
ZYPREXA	15MG	1	35/35
ZYPREXA	20MG	1	35/35
ZYPREXA ZYDIS	5MG	1	35/35
ZYPREXA ZYDIS	10MG	1	35/35
ZYPREXA ZYDIS	15MG	1	35/35
ZYPREXA ZYDIS	20MG	1	35/35

*Cancer diagnosis with non-daily chemotherapy required
**Available without pa with CA and HO diag.
*** Ranitidine syrup available without PA to users less than 6 years old.
MDV=Multidose Vial