

MaineCare PDL

PDL Effective January 1, 2026

* PLEASE NOTE: For a **search** box hit Ctrl F

* PLEASE NOTE: All **cost effective** generics applicable to DEL are considered **PREFERRED Drugs**. "BASIC" Covered Drugs are **bolded** with the Coverage Indicator of "MC / DEL".

General Criteria for all PDL categories- For more information or help using the PDL, providers may call 1-888-445-0497; members should call 1-866-796-2463. To access PDL and PA materials via the internet: www.maineicarepdl.org

A: Preferred Drugs- Unless otherwise specified, preferred drugs are available without prior authorization. Step order may apply for preferred drugs in some drug categories as indicated on the PDL. (See item "D" below for explanation of step order.)

B: Requests for Non-preferred Drugs- Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

C: Adequate Drug Trials- 1. The minimum trial period for each preferred and step order drug is two weeks, unless otherwise stated within specific PDL drug categories; trials with less than a two week duration will be reviewed on a case-by-case basis; 2. A trial will not be considered valid if preferred or non-preferred products were readily available (by override, individual purchase, samples, etc.); 3. Certain drug trials, such as with controlled substances, may require evidence that the preferred drugs were actually tried (example: with random pill counts and with random urine drug tests, using the methods of GC/MS with no lower threshold); 4. Adequate trials require documentation of attempts to titrate dose of preferred agents toward desired clinical response. 5. Adequate trials include prevention/treatment of common adverse effects associated with preferred agents (example: antinausea, antipruritic, etc.)

D: Step Order- When numbers appear in the "step order" column, it means drugs in this category must be used in the order specified, with the lower numbers having preference over the higher numbers. Chart notes should be provided to confirm drug trials that do not appear in the member's MaineCare drug profile.

E: The Department will institute strategies to ensure cost effectiveness through the use of an enhanced Drug Benefit. Preferred brand drugs will no longer be preferred in any PDL drug category where preferred generic drugs are also available. It is expected that preferred generics will be used prior to any preferred brands. This will be operated as a form of step care. Preferred brands in these categories will require prior authorization for these high utilization / high cost members.

F: Brand Name Medication Requests- (Must be submitted on the Brand Name PA request form)- According to MaineCare Benefits Manual Chapter II (80.07-5), when medically necessary covered brand-name drugs have an A-rated generic equivalent available, the most cost effective medically necessary version will be approved and reimbursed, since the brand-name and A-rated generic drugs have been determined by the FDA to be chemically and therapeutically equivalent. The Bureau does not make determinations as to whether or not a generic drug is clinically inferior or inequivalent to its brand version. This is the proper role of the FDA. Physicians should submit their reports of generic inequivalence directly to the FDA via the MEDWATCH.

G: PA requests for non- FDA Approved Indications- Decisions will be made on a case-by-case basis until the DUR committee is able to review the evidence and make a recommendation. Interim approvals and DUR recommendations for approval of a drug for a non- FDA approved indication will require a minimum of two published, peer reviewed, non contradicted, double blind, placebo-controlled randomized clinical studies establishing both safety and efficacy.

H: Dose Consolidation Requirements- Some drugs may also be affected by dose consolidation requirements. Please see Dose Consolidation List and/or Splitting Tables provided in the PDL.

I: Trials from Multiple Drug Classes - Trial/failure/intolerance to preferred agents from multiple classes within the same category or other categories of drugs may be required prior to the approval of non-preferred agents (e.g., Cymbalta, Zofran and others).

J: Drug-specific PA Forms- Drug-specific PA forms contain medical necessity documentation requirements and/or criteria that may not be repeated in the PDL. Drug-specific PA forms may be obtained on the web at www.maineicarepdl.org.

K: PA Exemptions for Prescribers- According to MaineCare Benefits Manual Chapter II (80.07-4), providers may receive a three (3) month exemption from prior authorization requirement for certain categories of drugs when they demonstrate high compliance with the Department's PDL. The Department will notify providers in writing which drug categories are included and what dates apply to the exemption. If a provider loses his/ her exemption, members who previously were not required to obtain a PA while the prescriber was exempt will be required to do so, and criteria for approval of that medication will need to be met.

L: Drug-Drug Interactions (DDI)- The DUR Committee has implemented new drug-drug interaction edits requiring prior authorization. Several drug-drug combinations and PDL drug categories are affected by new PA requirements. These will be indicated in the PDL with DDI notation. Please see the DDI document provided in the PDL.

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
ANTI INFECTIVE COMBO'S - MISC.	MC/DEL	ERYTHROMYCIN/SULF SUSR	MC		BACTRIM DS TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL	SEPTRA/DS TABS	MC		VABOMERE ¹		
	MC/DEL	SULFAMETHOXAZOLE/TRIMETH					
	MC/DEL	TRIMETHOPRIM/SULFAMETHOXA					
ANTI - FUNGALS							
ANTIFUNGALS - ASSORTED	MC/DEL	FLUCONAZOLE ¹	MC/DEL	6	LAMISIL TABS ⁴	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. The other criteria are listed on the Antifungal PA form including the required proof of a non-cosmetic fungal infection.
	MC/DEL	KETOCONAZOLE TABS ⁷	MC/DEL	6	ITRACONAZOLE		
	MC/DEL	NYSTATIN	MC	8	BREXAFEMME		
	MC/DEL	TERBINAFINE TABS ⁴	MC/DEL	8	CRESEMDA ⁹		
	MC/DEL	VORICONAZOLE TABS	MC/DEL	8	GRIFULVIN V TABS		
			MC	8	GRISEOFULVIN SUSP		
			MC	8	GRISEOFULVIN ULTRAMICROS TABS ⁸		
			MC	8	GRIS-PEG TABS		
			MC	8	REZZAYO ⁹		
			MC/DEL	8	SPORANOX SOLN ²		
			MC/DEL	8	SPORANOX PULSEPAK CAPS ³		
			MC/DEL	8	SPORANOX CAPS ³		
			MC/DEL	8	DIFLUCAN		
			MC/DEL	8	ERAXIS INJ ⁶		
			MC	8	GRIFULVIN SUSP		
			MC/DEL	8	ONMEL		
			MC/DEL	8	NOXAFL ⁵		
			MC/DEL	8	TOLSURA		
			MC/DEL	8	VFEND TABS		
			MC	8	VIVJOA		
ANTI - VIRALS							
ANTIRETROVIRALS - PREP	MC	APRETUDE	MC	8	TRUVADA ¹	Use PA Form# 20420	DDI: The concomitant use of the following drugs with Descovy is not recommended: tipranavir/ritonavir, St. John's wort, and the antimycobacterials rifabutin, rifampin, or rifapentine.
	MC	DESCOY ¹					
	MC	EMTRICITABINE-TENOFOVIR					
	MC	DISOP (ORAL) TAB					
	MC	YEZTUGO					
ANTIRETROVIRALS	MC/DEL	ABACAVIR TABS	MC/DEL	8	ABACAVIR SOL	Use PA Form# 20420	Fuzeon: Prescriber is either an HIV specialist provider or has consulted with one. Documentation of genotype testing is supplied and shows that there is no other potent, appropriate two or three drug oral regimen available, AND patient has a positive HIV viral load within past 6 months while on his/her current antiretroviral regimen AND the drug will be prescribed with at least two other drugs that are likely to be active based on the genotype testing.
	MC/DEL	ATAZANAVIR	MC/DEL	8	APTVUS		DDI: Reyataz requires prior authorization if it is currently being used in combination with either prevacid, pantoprazole, prilosec, or any currently non preferred PPI .
	MC	BIKTARVY	MC	8	ATRIPLA ¹		DDI: Norvir requires prior authorization if it is currently being used in combination with either enablex 15mg or vesicare 10mg.
	MC	CABENUVA	MC/DEL	8	CIMDUO		DDI: Preferred Crixivan caps requires prior authorization if it is currently being used in combination with either enablex 15mg or vesicare 10mg.
	MC	COMPLERA ¹	MC/DEL	8	COMBIVIR TABS		DDI: Administration with the following drugs: the anticonvulsants carbamazepine, oxcarbazepine, phenobarbital, and phenytoin; the antimycobacterials rifampin and rifapentine; proton pump inhibitors such as dexlansoprazole, esomeprazole, lansoprazole, omeprazole, pantoprazole, rabeprazole; systemic dexamethasone (more than a single dose); and St. John's wort with Odefsey is contraindicated.
	MC/DEL	DELSTRIGO	MC/DEL	8	EDURANT		Stribild: PA required; must provider rationale as to why the member's medical need cannot be met with preferred agents, particularly genvoya or combinations of preferred and agents AND must be antiretroviral treatment-naïve or virologically controlled on current therapy (HIV-1RNA < copies/ml) AND be HBV negative AND not be combined with other anti-retroviral agents.
	MC	DIDANOSINE	MC/DEL	8	EPZICOM ¹		DDI: Tivicay will require prior authorization is used with nevirapine, oxcarbazepine, phenytoin, phenobarbital, carbamazepine, and St. John's wort.
	MC/DEL	DOVATO	MC/DEL	8	FUZEON		DDI: Aatazanavir or Darunavir and the following drugs are contraindicated (due to potential for serious and/or life-threatening events or loss of therapeutic effect): aifuzosin, dronedarone, rifampin, irinotecan, ergotamine, methylergonovine, cisapride, St. John's wort, lovastatin, simvastatin, pimozide, nevirapine, sildenafil (when given as revatio for treatment of PAH), indinavir, triazolam, or PO midazolam will be non-preferred and require prior authorization if it is currently being used in combination with
	MC	EFAVIRENZ TAB	MC/DEL	8	INTELENCE		
	MC/DEL	EFAVIRENZ CAP	MC/DEL	8	ISENTRESS ³		
	MC	EFAVIRENZ-EMTRICITABINE-TENOFOVIR DF TAB	MC/DEL	8	ISENTRESS HD		
	MC	EMTRIVA ¹	MC	8	JULUCA		
			MC	8	KALETRA		

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
	MC	EPIVIR SOL	MC/DEL	8	LAMIVUDINE SOLN		tyost. DDI: Combined P-gp, UGT1A1 and strong CYP3A inhibitors may significantly increase plasma concentrations of Sunlenca. Concomitant administration of Sunlenca with these inhibitors is not recommended. Sunlenca: In combination with other antiretroviral(s) for the treatment of HIV-1 infection in heavily treatment-experienced adults with multidrug resistant HIV-1 infection failing their current antiretroviral regimen due to resistance, intolerance, or safety considerations.
	MC/DEL	EVOTAZ ¹	MC/DEL	8	LEXIVA		
	MC	GENVOYA ^{1,4}	MC/DEL	8	NEVIRAPINE		
	MC/DEL	ISENTRESS 400MG ⁵	MC	8	NORVIR		
	MC/DEL	ISENTRESS CHEW ³	MC/DEL	8	PIFELTRO		
	MC/DEL	ISENTRESS POWDER	MC	8	RETROVIR		
	MC/DEL	LAMIVUDINE TABS	MC	8	REYATAZ		
	MC/DEL	LAMIVUDINE/ZIDOVUDINE	MC/DEL	8	SELZENTRY		
	MC/DEL	LOPINAVIR-RITONAVIR SOL	MC	8	STAVUDINE		
	MC	LOPINAVIR-RITONAVIR TAB	MC	8	STRIBILD ¹		
	MC	ODEFSEY ¹	MC/DEL	8	SYMF ⁴		
	MC/DEL	PREZCOBIX	MC/DEL	8	SYMF ⁴ LO ⁴		
	MC	PREZISTA ²	MC/DEL	8	SYMTUZA		
	MC/DEL	RITONAVIR TAB 100MG	MC/DEL	8	TRIZIVIR TABS		
	MC	RUKOBIA ⁴	MC/DEL	8	VIRACEPT TABS		
	MC	SUNLENCA ⁴	MC	8	VITEKTA		
	MC	SUSTIVA ¹	MC	8	ZERIT		
	MC	TIVICAY	MC	8	VIDEX EC		
	MC	TIVICAY PD	MC	8	VIREAD TABS ¹		
	MC	TRIUMEQ ¹	MC/DEL	8	ZIAGEN TABS		
	MC	TROGARZO ⁴	MC/DEL	8	ZIAGEN SOL		
	MC	TYBOST	MC/DEL	9	VIRAMUNE XR		
	MC	VIREAD POW					
	MC/DEL	ZIDOVUDINE					
CYTO-MEGALOVIRUS AGENTS	MC	CIDOFOVIR	MC	4	VALCYTE TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC	FOSCARNET SODIUM	MC	8	FOSCAVIR		
	MC/DEL	GANCICLOVIR	MC/DEL	8	LIVTENCY ¹		
	MC/DEL	VALGANCICLOVIR	MC	8	PREVYMIS		
HERPES AGENTS	MC/DEL	VALACYCLOVIR HCL	MC/DEL	8	FAMCICLOVIR ¹	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
			MC	8	SITAVIG		
			MC	8	VALTREX TABS ¹		
			MC/DEL	9	FAMVIR TABS ¹		
INFLUENZA AGENTS	MC	AMANTADINE CAPS	MC		AMANTADINE TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC	RELENZA DISKHALER AEPB	MC		FLUMADINE TABS		
	MC/DEL	OSELTAMIVIR ¹	MC		FLUMIST		
			MC/DEL		RIMANTADINE HCL TABS		
			MC/DEL		TAMIFLU ¹		
			MC/DEL		TAMIFLU SUS		
			MC/DEL		XOFLUZA		
IMMUNE SERUMS							
IMMUNE SERUMS	MC	HYPERRHO INJ					
HEPATITIS AGENTS							
HEPATITIS C AGENTS	MC	SOFSBUVIR/VELPATASVIR ³ (Authorized generic labeler 72626 Aseuga Therapeutics)	MC/DEL		COPEGUS TABS	Use PA Form #10700	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC	MAVYRET ³	MC/DEL		DAKLINZA		
	MC/DEL	PEGASYS KIT ¹	MC		EPCLUSA ²		
	MC/DEL	PEGASYS SOLN	MC		HARVONI ²		
	MC/DEL	PEG-INTRON KIT ¹	MC/DEL		REBETOL CAPS		
	MC	RIBAVIRIN	MC		RIBAPAK		
	MC/DEL	RIBASPHERE	MC		SOVALDI ²		
			MC		VIEKIRA PAK ²		
			MC		VIEKIRA XR ²		
			MC		VOSEVI		
			MC/DEL		ZEPATIER ²		
HEPATITIS AGENTS - MISC.			MC		ACTIMMUNE	Use PA Form# 20420	Approved for chronic granulomatous disease, osteopetrosis and idiopathic pulmonary fibrosis.

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
					receptor (AChR) antibody positive. 6. For the treatment of patients between ages 4-17 years of age.	Migraine: Consideration for Botox approvals will only be made after failures of required trials of the following preferred medications: tricyclic or venlafaxine, beta blocker, valproic acid, topiramate. Firdapse is recommended for the treatment of Lambert-Eaton myasthenic syndrome (LEMS) in adults.	
NEUROLOGICS- hATTR AGENTS			MC MC MC MC MC/DEL MC/DEL MC/DEL	8 8 8 8 8 8 8	AMVUTTRA ¹ ATTRUBY ONPATRO ¹ TEGSEDI ¹ VYNDAMAX ¹ VYNDAQEL ¹ WAINUA ¹	Use PA Form# 20420 1. PA required for appropriate diagnosis.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval. Tegsedi should be non-preferred and approved for patients for whom other treatments, including Onpatro, have been ineffective. Vyndamax will be considered for the treatment of the cardiomyopathy of wild type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) in adults to reduce cardiovascular mortality and cardiovascular-related hospitalization.
NEUROLOGICS- SMA		GENE			GENE	Use PA Form# 20420 1. Clinical PA is required to establish diagnosis and medical necessity. 2. For patients 2 months of age and older.	Zolgensma: The patient is less than 2 years of age AND The diagnosis is spinal muscular atrophy (SMA) AND the patient has bi-allelic mutations of the SMN1 gene AND the patient does not have advanced SMA (e.g. complete paralysis of limbs or permanent ventilator dependence) AND medication is prescribed per the dosing. Spinraza: The diagnosis is spinal muscular atrophy (SMA) type 1, 2, or 3 (results of genetic testing must be submitted) AND The patient has at least 2 copies of the SMN2 gene AND The prescriber is a neurologist, pulmonologist, or other physician with expertise in treating SMA AND Baseline motor ability has been established using one of the following exams: Hammersmith Infant Neurological Exam (HINE) Hammersmith Functional Motor Scale Expanded (HFMSE) Upper Limb Module Test (non-ambulatory) Children's Hospital of Philadelphia Infant Test of Neuromuscular Disorders (CHOP INTEND) AND Prior to starting therapy, and prior to each dose, the following laboratory tests will be conducted: Treating provider attests the member has a platelet count > 50,000/ml or greater Treating provider agrees to do platelet count and coagulation test before each dose Treating provider agrees to do a quantitative spot urine protein test before each dose Concomitant use of Spinraza and Zolgensma is investigational and will not be approved AND Use of Spinraza after gene replacement therapy, including Zolgensma is investigational and will not be approved Note: Initial approval will be granted for 4 loading doses (the first 3 loading doses should be administered at 14-day intervals; the 4th loading dose should be administered 30 days after the 3rd dose). Renewal may be granted for up to 12 months with a maximum of 3 doses approved per year (12mg (5ml) every 4 months). For therapy continuation, clinical documentation must be submitted documenting improvement or maintenance of motor ability OR slower progression of disease than would otherwise be expected.
NEUROLOGICS- RETT SYNDROME			MC		DAYBUE ^{1,2}	Use PA Form# 20420 1. Clinical PA required for appropriate diagnosis 2. For the treatment of patients 2 years of age and older.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ALS DRUGS	MC/DEL	RILUZOLE	MC MC MC MC MC MC		EXSERVAN QALSODY RILUTEK TABS RADICAVA ¹ RELYVRI ¹ TIGLUTIK	Use PA Form# 20420 1. Clinical PA for indication required.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Qalsody: For the treatment of amyotrophic lateral sclerosis (ALS) in adults who have a mutation in the superoxide dismutase 1 (SOD1) gene. Continued approval for this indication may be contingent upon verification of clinical benefit in confirmatory trial(s).
MOVEMENT DISORDERS	MC MC MC MC	AUSTEDO ¹ AUSTEDO XR ¹ INGREZZA ¹ TETRABENAZINE ¹	MC/DEL		XENAZINE	Use PA Form# 20420 1. Clinical PA required for appropriate diagnosis	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Avoid concomitant use of VMAT2 inhibitors with MAO inhibitors (e.g. isocarboxazid, phenelzine, or selegiline). Concomitant use with strong CYP3A4 inducers (e.g. rifampin, carbamazepine, phenytoin, St. John's wort) is not recommended

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
			MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC MC/DEL MC/DEL		METHITEST TAB METHYLTESTOSTERONE CAP OXANDROLONE STRIANT MUC ER TESTIM TESTOSTERONE GEL PACKETS TESTOSTERONE SOL TESTRED CAPS TLANDO VOGELXO XYOSTED		
ESTROGENS - PATCHES / TOPICAL	MC MC/DEL MC/DEL	EVAMIST MINIVELLE PATCH VIVELLE-DOT PTTW	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	5 8 8 8 8	ESTRADIOL PTWK DIVIGEL ¹ CLIMARA PTWK ELESTRIN ¹ MENOSTAR PATCH	Use PA Form# 20420 1. Step order drugs must be used in specified step order.	Approved for failures on multiple oral estrogen agents after 90 day trials or if unable to swallow any oral medication.
ESTROGENS - TABS	MC/DEL MC/DEL	ESTRADIOL PREMARIN TABS	MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC		ENJUVIA ESTRADIOL-NORETHINDRONE ESTRACE TABS ESTRATAB TABS MENEST TABS NORETHINDRON-ETHINYL ORTHO-EST TABS	Use PA Form# 20420 Must fail preferred products before non-preferred products.	Preferred drugs must be tried for at least 90 days and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ESTROGEN COMBO'S	MC/DEL MC/DEL MC/DEL MC/DEL	ANGELIQ COMBIPATCH PTTW PREMPHASE TABS PREMPRO TABS	MC/DEL MC/DEL MC MC/DEL MC/DEL		FEMHRT 1/5 TABS ¹ FYAVOLV LOPREEZA TAB ORTHO-PREFEST TABS ¹ SYNTEST H.S. TABS ¹	Use PA Form# 20420 1. Must fail Premphase and Prempro products before non preferred products.	Preferred drugs must be tried for at least 90 days and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
PROGESTINS	MC/DEL MC/DEL MC MC	MEDROXYPROGESTERONE ACETA ¹ NORETHINDRONE ACETATE TABS ¹ 17-ALPH HYDROXYPROGESTERONE PWDR PROGESTERONE CAPS	MC/DEL MC MC MC/DEL MC/DEL		AYGESTIN TABS CYCRIN TABS PROGESTERONE POWD PROMETRIUM CAPS PROVERA TABS	Use PA Form# 20420 1. Must fail Medroxyprogesterone and Norethindrone products before non-preferred products.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ENDOMETROSIS							
CENTRAL PRECOCIOUS PUBERTY AGENTS	MC	FENSOLVI ¹				Use PA Form# 20420 1. For pediatric patients 2 years of age and older with central precocious puberty (CPP).	
ENDOMETROSIS- NASAL	MC/DEL	SYNAREL (NASAL) SPRAY				Use PA Form# 20420	Synarel is also indicated for central precocious puberty.
ENDOMETROSIS/ UTERINE FIBROIDS- ORAL	MC MC MC/DEL	MYFEMBREE ^{1,2} ORIAHNN ¹ ORILISSA ¹				Use PA Form# 20420 1. Prior treatment of NSAID and hormonal contraceptives required. 2. Limited to 24 months due to the risk of continued bone loss, which may not be reversible.	
ENDOMETROSIS- INJECTABLE	MC/DEL	DEPO-SUBQ PROVERA 104				Use PA Form# 20420	
CONTRACEPTIVES							
CONTRACEPTIVES - PROGESTIN ONLY	MC/DEL MC/DEL MC MC MC/DEL MC/DEL	CAMILA TABS ERRIN INCASSIA TAB HEATHER TAB NORETHINDRONE ACETATE 0.35MG TABS SLYND	MC/DEL MC/DEL MC		JOLIVETTE NORA-BE TABS ORTHO MICRONOR TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. If member experienced adverse reactions, consider using Oral Contraceptives from other groups. DDI: Preferred Oral Contraceptives will now be non-preferred and require prior authorization if it is currently being used in combination with Tracleer.
CONTRACEPTIVES - INJECTABLE	MC/DEL	MEDROXYPROGESTERONE ACETATE 150mg IM	MC/DEL		DEPO-PROVERA 150 mg SUSP	Use PA Form# 20420	The preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
DIABETIC - DPP- 4 ENZYME INHIBITOR	MC/DEL MC/DEL	JANUVIA ^{1,2} TRADJENTA ²	MC MC/DEL MC/DEL MC		BRYNOVIN NESINA QTERN ZITUVIO	Use PA Form# 20420 1. Preferred if therapeutic doses of metformin are seen in members drug profile for at least 60 days within the past 18 months or if phosphate binder is currently seen in the members drug profile. 2. Dosing limits apply. Please refer to Dose Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. BRYNOVIN: In addition to tried and failed Preferred Agents, Brynovin requires tried and failed Non-Preferred Agent Zituvio.
DIABETIC - DPP- 4 ENZYME INHIBITOR-COMBO	MC/DEL MC/DEL MC/DEL	JANUMET ^{1,2} JANUMET XR ^{1,2} JENTADUETO ¹	MC/DEL MC/DEL MC MC/DEL MC MC		JENTADUETO XR KAZANO KOMBIGLYZE XR OSENI ZITUVIMET ZITUVIMET XR	Use PA Form# 20420 1. Preferred if therapeutic doses of metformin are seen in members drug profile for at least 60 days within the past 18 months or if phosphate binder is currently seen in the members drug profile. 2. Dosing limits apply. Please refer to Dose consolidation list.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Zituvimet/ Zituvimet XR: Approvals will require trial of preferred sitagliptin/metformin products or other preferred diabetic agents.
DIABETIC - LANCET-LANCET DEVICE						Use PA Form# 20420	Please refer to the MaineCare Preferred Diabetic Supply List available at www.maineicarepdi.org
DIABETIC - SYRINGES-NEEDLES						Use PA Form# 20420	Please refer to the MaineCare Preferred Diabetic Supply List available at www.maineicarepdi.org
DIABETIC - OTHER			MC		SYMLIN	Use PA Form #20420	
SGLT 2 INHIBITORS	MC/DEL MC/DEL	FARXIGA JARDIANCE	MC/DEL MC/DEL		INVOKANA ¹ STEGLATRO	Use PA Form# 20420 1. Dosing limits apply please refer to Dose Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
SGLT 2 INHIBITOR COMBINATIONS	MC/DEL MC/DEL MC/DEL	SYNJARDY SYNJARDY XR XIGDUO XR	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL		GLYXAMBI INVOKAMET INVOKAMET XR SEGLUROMET STEGLUJAN TRIJARDY XR	Use PA Form# 20420	Preferred drugs must be tried for at least 3 months at full therapeutic doses and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Glyxambi /Xigduo XR- Verify prior trials and failures or intolerance of preferred treatments from other diabetic categories. Synjardy XR is not recommended for patients with type 1 DM or for the treatment of diabetic ketoacidosis.
DIABETIC MONITOR	MC	RELION TRUEMETRIX AIR BLOOD GLUCOSE MONITORING SYSTEM TRUEMETRIX AIR BLOOD GLUCOSE MONITORING SYSTEM TRUEMETRIX BLOOD GLUCOSE MONITORING SYSTEM	MC MC MC MC MC MC MC MC MC		ACCUCHECK ASCENSA ASSURE CONTOUR BREEZE Z EXACTECH FREESTYLE INSULIN FREESTYLE LITE SYSTEM KIT PRECISION XTRA METER PRODIGY	Use PA Form# 20420	Effective October 25th 2007, approvals for all non preferred meters/ test strips will require medical necessity documenting clinically significant features that are not available on any of the preferred meters.
DIABETIC TEST STRIPS	MC/DEL MC	RELION TRUEMETRIX TRUEMETRIX	MC MC MC MC MC MC MC MC MC MC		ACCUCHECK ASCENSA ASSURE CONTOUR BREEZE Z EXACTECH FREESTYLE FREESTYLE LITE FREESTYLE INSULIN PRECISION XTRA PRODIGY	Use PA Form# 20420	Effective October 25th 2007, approvals for all non preferred meters/ test strips will require medical necessity documenting clinically significant features that are not available on any of the preferred meters. Effective October 1, 2023, a maximum of 100 blood glucose test strips every 90 days will be available without Prior Authorization for members currently utilizing continuous glucose monitors (CGM).
INCRETIN MIMETIC	MC/DEL MC MC/DEL	RYBELSUS TRULICITY VICTOZA	MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL	5 8 8 8 8 8	OZEMPIC ADLYXIN BYDUREON BCISE MOUNJARO SOLIQUA XULTOPHY	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Soliqua must try both insulin and a preferred incretin mimetic and have a medical necessity for use that is not based on convenience or simply due to the fact that one injection is needed instead of two.

CATEGORY	Coverage Indicator	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
DIABETIC - ORAL SULONYLUREAS	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	CHLORPROPAMIDE TABS GLIMEPIRIDE GLIPIZIDE TABS GLIPIZIDE ER TABS GLYBURIDE MICRONIZED TABS GLYBURIDE TABS ¹ TOLAZAMIDE TABS TOLBUTAMIDE TABS	MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL		AMARYL TABS DIABETA TABS GLUCOTROL TABS GLUCOTROL XL TBCR GLYNASE TABS MICRONASE TABS	Use PA Form# 20420 1. PA required for members ≥ 65 . Glyburide has a greater risk of severe prolonged hypoglycemia in older adults.	Preferred drugs must be tried for at least 3 months at full therapeutic doses and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: All sulfonylureas (except glyburide) will now be non-preferred and require prior authorization if it is currently being used with either ranitidine or cimetidine. DDI: Glimepiride will now be non-preferred and require prior authorization if it is currently being used with either fluconazole (except 150mg strength) or fluvoxamine. Amaryl is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for concurrent use with either fluconazole or fluvoxamine.
DIABETIC -ORAL BIGUANIDES	MC/DEL MC/DEL	METFORMIN HCL TABS METFORMIN ER	MC MC MC MC/DEL		GLUCOPHAGE TABS GLUCOPHAGE XR TB24 FORTAMET METFORMIN ER OSMOTIC	Use PA Form# 20420	Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
DIABETIC - THIAZOL / BIGUANIDE COMBO			MC/DEL MC/DEL MC MC		ACTOPLUS MET ¹ ACTOPLUS MET XR AVANDARYL ¹ AVANDAMET TABS ¹	Use PA Form# 20420 1. Requires use of Actos, Metformin, or other preferred anti-diabetics.	DDI: Actos, Avandia, or any combination product with Actos or Avandia will now be non-preferred and require prior authorization if it is currently being used with gemfibrozil.
DIABETIC - / THIAZOL	MC/DEL	PIOGLITAZONE HCL ¹	MC/DEL MC		ACTOS TABS ³ AVANDIA TABS ²	Use PA Form# 20420 1. Pioglitazone HCL is non-preferred as monotherapy. Pioglitazone HCL is preferred if therapeutic doses of metformin, sulfonylurea or insulin are seen in members drug profile for at least 60 days within the past 18 months. 2. Current users of Avandia who have tried Actos will be able to continue use of Avandia. 3. Dosing limits apply. See Dose Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Actos, Avandia, or any combination product with Actos or Avandia will now be non-preferred and require prior authorization if it is currently being used with gemfibrozil.
DIABETIC - ALPHAGLUCOSIDASE			MC		PRECOSE TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
DIABETIC - SULONYLUREA / BIGUANIDE	MC/DEL	GLYBURIDE/METFORMIN	MC MC MC/DEL		GLUCOVANCE TABS ¹ METAGLIP TABS ¹ DUETACT ²	Use PA Form# 20420 1. Use individual ingredients. 2. Use Actos with generic glimepiride.	Approved for patients failing to achieve good diabetic control with maximal doses of individual components.
DIABETIC - MEGLITINIDES	MC	NATEGLINIDE	MC/DEL MC/DEL		PRANDIN TABS STARLIX TABS	Use PA Form# 20420	Preferred drugs from other diabetic sub-categories must be tried and failed due to lack of inadequate diabetic control or intolerable side effects before non-preferred drug will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Prandin is non-preferred but with any prior authorization requests, the member's drug profile will also be monitored for current use with both sporanox and gemfibrozil, due to a significant drug-drug interaction.
GLUCOSE ELEVATING AGENTS							
GLUCOSE ELEVATING AGENTS	MC/DEL MC/DEL MC/DEL	BAQSIMI ¹ GVOKE ² ZEGALOGUE ³	MC		GLUCAGON DIAGNOSTIC KIT	Use PA Form# 20420 1. For the treatment of patients ≥ 4 years of age. 2. For the treatment of patients ≥ 2 years of age. 3. For the treatment of patients ≥ 6 years of age.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved (in step order), unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
THYROID							
THYROID EYE DISEASE			MC		TEPEZZA	Use PA Form# 20420	
THYROID HORMONES	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	ARMOUR THYROID TABS CYTOMEL TABS ERMEZA ¹ LEVOTHROID TABS LEVOTHYROXINE SODIUM TABS LEVOXYL TABS UNITHYROID TABS	MC MC/DEL MC MC/DEL		LEVOTHYROXINE SODIUM SOLR LIOTHYRONINE SYNTHROID TABS THYQUIDITY	Use PA Form# 20420 1. Clinical PA is required to confirm diagnosis of dysphagia.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTITHYROID THERAPIES	MC/DEL MC/DEL	METHIMAZOLE TABS PROPYLTHIOURACIL TABS	MC/DEL		TAPAZOLE TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required			Criteria
SOMATOSTATIC AGENTS			MC/DEL MC MC MC/DEL MC	7 8 8 8 8	OCTREOTIDE INJ ¹ BYNFEZIA ¹ MYCAPSSA ¹ SANDOSTATIN ¹ SOMATULINE ¹	Use PA Form# 10710 1. Non-preferred products must be used in specified step order.		
GROWTH HORMONE ANTAGONISTS								
GH ANTAGONISTS			MC		SOMAVERT	Use PA Form# 10710	Approved for acromegaly patients failing surgery/radiation/drug therapy including bromocriptine and sandostatin.	
VASOPRESSIN RECEPTOR ANTAGONIST								
VASOPRESSIN RECEPTOR ANTAGONIST			MC MC/DEL		JYNARQUE ¹ SAMSCA	Use PA Form# 20420 1. Clinical PA required for appropriate diagnosis	Samsca Drug Warning- Avoid use in patients with underlying liver disease, including cirrhosis, because the ability to recover from liver injury may be impaired. Limit duration of therapy to 30 days to minimize the risk of liver injury. DDI: Jynarque- Concomitant use with strong CYP3A inhibitors is contraindicated. Avoid concomitant use of Jynarque with OATP1B1/B3 and OAT3 substrates (e.g. statins, bosantan, glyburide, nateglinide, repaglinide, methotrexate, furosemide).	
URINARY INCONTINENCE								
VASOPRESSINS	MC/DEL MC/DEL	DESMOPRESSIN TABS DDAVP SOLN	MC/DEL MC/DEL MC MC/DEL MC	5 6 8 8 8	DDAVP TABS DESMOPRESSIN SPRAY ¹ DESMOPRESSIN ACETATE SOLN ¹ NOCDURNA ¹ NOCTIVA ¹	Use PA Form# 20420 1. Products must be used in specified step order. Nocturnal enuresis patients will be encouraged to periodically attempt stopping DDAVP.	Approved for central diabetes insipidus and for nocturnal enuresis. For nocturnal enuresis- must be over 6 years old, must fail an adequate trial of alarm training (higher success rate, lower relapse rate) and must periodically attempt weaning (at 6 month intervals).	
ANTISPASMODICS	MC/DEL MC/DEL	OXYBUTYNIN TOLTERODINE	MC/DEL MC/DEL MC/DEL	8 8 8	DARIFENACIN ER TAB DITROPAN FLAVOXATE HCL TAB	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.	
ANTISPASMODICS - LONG ACTING	MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	FESOTERODINE GELNIQUE GEL PACKET MYRBETRIQ OXYBUTYNIN ER TABS OXYTROL SOLIFENACIN SUCCINATE TAB TROSPiUM	MC MC/DEL MC MC/DEL MC/DEL MC MC	8 8 8 8 8 8 8	DITROPAN XL TBCR ENABLEX ^{1,2} GEMTESA ² TOLTERODINE TAB TOVIAZ VESICARE ¹ VESICARE LS ³	Use PA Form# 20420 1. See Criteria Section. 2. Use a preferred long acting antispasmodic. 3. For the treatment of patients \geq 2 years of age.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. 1. Vesicare 5mg and Enablex 7.5mg maximum doses if given with drugs known to be significant CYP3A4 inhibitors (ketoconazole, sporanox, erythromycin, fluconazole, nefazodone, neflunavir, and ritonavir). DDI: Enablex 15mg and Vesicare 10mg will now be non-preferred and require prior authorization if they are currently being used in combination with any of the following medications: clarithromycin, erythromycin, ketek, crixivan, norvir, ketoconazole, fluconazole (except 150mg strength), sporanox, or nefazodone.	
CHOLINERGIC	MC/DEL	BETHANECHOL	MC/DEL		URECHOLINE	Use PA Form# 20420		
HYPERAMMONIA TREATMENTS	MC	CARBAGLU TABS	MC		CARGLUMIC ACID TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.	
UREA CYCLE DISORDER	MC MC	BUPHENYL TABLET PHEBURANE GRANULES	MC MC MC MC/DEL MC/DEL		BUPHENYL POWDER RAVICTI LIQUID OLPRUVA SODIUM PHENYLBUTYRATE POWDER SODIUM PHENYLBUTYRATE TAB	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Olpruva: As adjunctive therapy to standard of care, which includes dietary management, for the chronic management of adult and pediatric patients weighing 20kg or greater and with a body surface area (BSA) of 1.2m ² or greater, with urea cycle disorders (UCDs) involving deficiencies of carbamylphosphate synthetase (CPS), ornithine transcarbamylase (OTC), or argininosuccinic acid synthetase (AS).	
METABOLIC MODIFIER								
HERED. TYROSINEMIA			MC MC MC	6 6 8	ORFADIN NITYR HARLIKU ¹	Use PA Form# 20420 1. Clinical PA is required to establish diagnosis and medical necessity.	Approved for Type 1 hereditary tyrosinemia patients. Must include laboratory evidence of dx at first PA. Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.	
FABRY DISEASE AGENTS			MC MC MC/DEL		ELFABRIO ¹ FABRAZYME ² GALAFOLD ¹	Use PA Form# 20420 1. Clinical PA to verify appropriate diagnosis. 2. For the treatment of patients 2 years of age and older.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Elfabrio and Galfold: For the treatment of adults with confirmed Fabry disease.	
ANTIHYPERTENSIVES / CARDIAC								
CARDIAC GLYCOSIDES	MC/DEL MC/DEL MC/DEL	DIGITEK TABS DIGOXIN LANOXIN				Use PA Form# 20420		
CARDIAC MYOSIN INHIBITORS			MC		CAMZYOS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Camzyos: For the treatment of adults with symptomatic New York Heart Association (NYHA) class II-III obstructive hypertrophic cardiomyopathy (HCM) to improve functional capacity and symptoms. DDI: Concomitant use of Camzyos with a moderate to strong CYP2C19 inhibitor or a strong CYP3A4 inhibitor is contraindicated.	
CARDIAC - SINUS NODE INHIBITORS			MC		CORLANOR	Use PA Form# 20420	In patients with stable, symptomatic chronic heart failure with left ventricular ejection fraction \leq 35%, who are in sinus rhythm with resting heart rate \geq 70 beats per minute (bpm) and either are on maximally tolerated doses of beta-blockers or have a contraindication to beta-blocker use.	

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
CARDIAC- ERAs			MC		TRYVIO	Use PA Form#20420	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p> <p>Tryvio: In combination with other antihypertensive drugs, is indicated for the treatment of resistant hypertension, to lower blood pressure (BP) in adult patients who are not adequately controlled on other drugs. Resistant HTN is defined as a patient who takes at least 3 different class antihypertensive medications with complementary mechanisms including thiazide, ACE inhibitor, ARB, long-acting calcium channel blocker, with a trial of spironolactone, unless contra-indicated.</p>
CARDIAC- SOLUBLE GUANYLATE CYCLASE STIMULATORS			MC/DEL		VERQUVO	Use PA Form# 20420	
CARDIAC RISK REDUCTION- SGLT2/GLP-1			MC MC MC/DEL		INPEFA ¹ LODOCO WEGOVY	Use PA Form #23976 1. To reduce the risk of cardiovascular death, hospitalization for heart failure, and urgent heart failure visit in adults with: Heart failure or Type 2 diabetes mellitus, chronic kidney disease, and other cardiovascular risk factors.	<p>Other Preferred SGLT inhibitors must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p> <p>Lodoco: Patient must have tried and failed generic colchicine due to lack of efficacy or intolerable side effects</p> <p>Wegovy: Patient does not have diagnosis of diabetes, end stage renal disease/dialysis, or HFrEF (EF < 45%)</p> <ul style="list-style-type: none"> • Patient has BMI > 27 kg/m², and is not being used for weight loss only • Patient has history of at least one of the following: <ul style="list-style-type: none"> o Stroke o Myocardial Infarction o Symptomatic peripheral arterial disease
ANTIANGINALS--Isosorbide Dinitrate/ Mono-Nitrates	MC/DEL MC/DEL	ISOSORBIDE MONONITRATE TABS ISOSORBIDE MONONITRATE ER	MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC		DILATRATE SR CPCR ISORDIL TITRADOSE TABS ISOSORBIDE DINITRATE SUBL ISOSORBIDE DINITRATE CR TBCR ISOSORBIDE DINITRATE ER TBCR ISOSORBIDE DINITRATE TD TBCR IMDUR TB24 ISMO TABS MONOKET TABS	Use PA Form# 20420	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p>
NITRO - OINTMENT/CAP/CR	MC/DEL MC/DEL MC MC	NITROBID OINT NITROGLYCERIN CPCR NITROL OINT NITRO-TIME CPCR				Use PA Form# 20420	
NITRO - PATCHES	MC/DEL MC/DEL	NITROGLYCERIN PT24 NITRO-DUR PT 24 0.8MG	MC MC/DEL		NITRODISC PT24 NITRO-DUR PT24	Use PA Form# 20420	<p>All preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p>
NITRO - SUBLINGUAL/ SPRAY	MC/DEL	NITROSTAT SUBL	MC/DEL MC MC		NITROQUICK SUBL NITROLINGUAL SOLN NITROLINGUAL TABS	Use PA Form# 20420	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p>
BETA BLOCKERS - NON SELECTIVE	MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	CARVEDILOL LEVATOL TABS NADOLOL TABS PINDOLOL TABS PROPRANOLOL HCL SOLN ¹ PROPRANOLOL HCL TABS ¹ PROPRANOLOL HCL 60MG TABS PROPRANOLOL LA CAPS RANOLAZINE ER TABS SOTALOL AF SOTALOL HCL TABS TIMOLOL MALEATE TABS	MC MC/DEL MC MC MC MC/DEL MC/DEL MC/DEL MC MC MC MC MC MC MC MC		ASPRUZO BETAPACE TABS BETAPACE AF TABS COREG CR ³ COREG TABS CORGARD TABS INDERAL TABS HEMANGEOL SOL INDERAL XL CAP INDERAL LA CPCR INNOPRAN XL RANEXA	Use PA Form# 20420 1. Recommend using BID since its effects do not last 24 hours. 2. Please use other strengths in combination to obtain this dose. 3. Dosing limits still apply. Please see dose consolidation list.	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p> <p>DDI: Concomitant use of Ranolazine products with strong CYP3A inhibitors, including ketoconazole, itraconazole, clarithromycin, nefazodone, nefazodone, ritonavir, indinavir, and saquinavir, is contraindicated.</p>
BETA BLOCKERS - CARDIO SELECTIVE	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	ACEBUTOLOL HCL CAPS ATENOLOL TABS ¹ BETAXOLOL HCL TABS BISOPROLOL FUMARATE TABS BYSTOLIC METOPROLOL TARTRATE TABS ¹ METOPROLOL ER NEBIVOLOL HCL TAB	MC MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC MC MC MC MC MC MC		KERLONE TABS LOPRESSOR TABS SECTRAL CAPS TENORMIN TABS TOPROL XL TB24 ZEBETA TABS	Use PA Form# 20420 1. Recommend using Atenolol (and Metoprolol) BID since its effects do not last 24 hours.	<p>Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.</p>

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
BETA BLOCKERS - ALPHA / BETA	MC/DEL	LABETALOL HCL TABS	MC		TRANDATE TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
BETA BLOCKERS & DURECTIC COMBOS	MC/DEL	METOPROLOL-HYDROCHLOROTHIAZIDE TAB	MC/DEL		DUTOPROL	Use PA Form# 20420	
CALCIUM CHANNEL BLOCKERS-- Amlodipine, Felodipines, Nifedipines, Nisoldipine, and Verapamil	MC/DEL	AMLODIPINE ¹	MC/DEL MC MC/DEL		KATERZIA NORLIQVA NORVASC TABS ¹	Use PA Form# 20420 1. Dosing limits apply, see Dose Consolidation List.	
	MC	DILTIA XT CP24	MC/DEL	5	DILACOR XR CP24 ¹	Use PA Form# 20420	Preferred drugs must be tried and failed (in step-order) due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL	DILTAZEM HCL ER CP24	MC/DEL	6	TAZTIA ¹	Use PA Form# 20420	1. Products must be used in specified order or PA will be required. Just write "Diltiazem 24-hour" and the pharmacy will use a preferred long acting Diltiazem that does not require PA.
	MC/DEL	DILTAZEM HCL XR CP24	MC/DEL	8	DILTAZEM HCL TABS ¹	Use PA Form# 20420	DDI: All preferred Diltiazem will now be non-preferred and require prior authorization if they are currently being used in combination with either enablex 15mg or vesicare 10mg. All non-preferred Diltiazem require prior authorization, but with any prior authorization request, the member's drug profile will also be monitored for current use with enablex 15mg or vesicare 10mg.
	MC/DEL	DILTAZEM CD 300MG CP24	MC/DEL	8	DILTAZEM HCL ER CP12 ¹	Use PA Form# 20420	
	MC/DEL	DILTAZEM CD 360MG CP24	MC/DEL	8	DILTAZEM HCL ER CP12 ¹	Use PA Form# 20420	
	MC	CARTIA XT CP24 ¹					
	MC/DEL	DILTAZEM CD CP24 ¹					
	MC/DEL	DILTAZEM HCL ER CP24 ¹					
	MC/DEL	DILTAZEM XR CP24 ¹					
	MC/DEL	TIAZAC CP24 ¹					
			MC/DEL MC/DEL		PLENDIL TB24 FELODIPINE	Use PA Form# 20420	Other Preferred calcium channel blockers must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
			MC MC		CARDENE SR CPCR NICARDIPINE HCL CAPS	Use PA Form# 20420	Other Preferred calcium channel blockers must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL	AFEDITAB CR	MC/DEL		ADALAT CC TBCR ¹	Use PA Form# 20420	Preferred drug must be tried and failed in step order due to lack of efficacy or intolerable side effects before non-preferred drugs in step order will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL	NIFEDIAC CC	MC/DEL		NIFEDIPINE CAPS	Use PA Form# 20420	
	MC/DEL	NIFEDICAL XL TBCR	MC/DEL		PROCARDIA CAPS	Use PA Form# 20420	
	MC/DEL	NIFEDIPINE TBCR	MC/DEL		PROCARDIA XL TBCR	Use PA Form# 20420	
	MC/DEL	NIFEDIPINE ER TBCR					
			MC MC		SULAR TB24 SULAR CR ¹	Use PA Form# 20420 1. Established users of 10MG and 20MG strengths are grandfathered.	
	MC/DEL	VERAPAMIL HCL CR TBCR	MC/DEL		CALAN TABS	Use PA Form# 20420	Preferred drugs must be tried and failed (in step-order) due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL	VERAPAMIL HCL ER TBCR	MC/DEL		CALAN SR TBCR	Use PA Form# 20420	
	MC/DEL	VERAPAMIL HCL SR TBCR	MC/DEL MC MC/DEL		COVERA-HS TBCR ISOPTIN-SR VERAPAMIL HCL ER CP24 VERAPAMIL HCL SR CP24 VERAPAMIL HCL TABS VERELAN CP24 VERELAN PM CP24	Use PA Form# 20420 Products must be used in specified order or PA will be required. Just write "Verapamil 24-hour" and the pharmacy will use a preferred long acting generic that does not require PA.	
ANTIARRHYTHMICS	MC/DEL	AMIODARONE HCL	MC/DEL		CORDARONE	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL	DISOPYRAMIDE	MC/DEL		DISOPYRAMIDE	Use PA Form# 20420	
	MC/DEL	FLECAINIDE	MC/DEL		MULTAQ	Use PA Form# 20420	
	MC/DEL	MEXILETINE HCL	MC/DEL		NORPACE	Use PA Form# 20420	DDI: Amiodarone will now be non-preferred and require prior authorization if it is currently being used in combination with either lovastatin (doses greater than 40mg/day) or lipitor (doses greater than 20mg/day) or levofloxacin or gemifloxacin, or moxifloxacin, or ofloxacin.
	MC/DEL	PROCAINAMIDE	MC/DEL		PACERONE	Use PA Form# 20420	
	MC/DEL	PROPAFENONE	MC		QUINIDEX	Use PA Form# 20420	DDI: Multaq will be preferred unless the following medications are seen in the member's drug profile within the last 35 days for brand name medications or 90 days for generic medications: erythromycin, amiodarone and other antiarrhythmics, TCA's, phenothiazine, ketoconazole, itraconazole, voriconazole, cyclosporine, telithromycin, clarithromycin, nefazodone, ritonavir.
	MC	QUINAGLUTE	MC/DEL		TAMBOCOR	Use PA Form# 20420	
	MC/DEL	QUINIDINE GLUCONATE	MC/DEL		TIKOSYN ¹	Use PA Form# 20420	
	MC/DEL	QUINIDINE SULFATE	MC/DEL		RYTHMOL SR	Use PA Form# 20420	
			MC/DEL		RYTHMOL	Use PA Form# 20420	
ACE INHIBITORS	MC/DEL	BENAZEPRIL HCL	MC	5	MAVIK TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Non-preferred products are subject to step-order requirements unless clinical circumstances warrant exception.
	MC/DEL	CAPTOPRIL TABS	MC/DEL	5	ACCUPRIL TABS	Use PA Form# 20420	
	MC/DEL	ENALAPRIL MALEATE TABS	MC/DEL	8	ACEON TABS ¹	Use PA Form# 20420	
	MC/DEL	FOSINOPRIL SODIUM	MC/DEL	8	ALTACE CAPS ¹	Use PA Form# 20420	
	MC/DEL	LISINOPRIL TABS	MC	8	EPANED	Use PA Form# 20420	
	MC/DEL	RAMIPRIL	MC/DEL	8	LOTENSIN TABS ¹	Use PA Form# 20420	
	MC/DEL	QUINAPRIL HCL	MC/DEL	8	MOEXIPRIL HCL ¹	Use PA Form# 20420	
			MC	8	MONOPRIL HCT TABS ¹	Use PA Form# 20420	

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
			MC/DEL MC MC/DEL MC/DEL	8 8 8 8	PRINIVIL TABS ¹ QBRELIS UNIVASC ¹ ZESTRIL TABS ¹		
ANGIOTENSIN RECEPTOR BLOCKER	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	AMLODIPINE-OLMESARTAN TAB ³ IRBESARTAN ¹ LOSARTAN ¹ MICARDIS TABS ³ OLMESARTAN ¹ TELMISARTAN ¹	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC	8 8 8 8 8 8 8	ATACAND TABS AVAPRO BENICAR TABS COZAAR DIOVAN EDARBI TEVETEN TABS	Use PA Form# 20420 1. Dosing limits apply, please see Dose Consolidation List. 2. Use preferred active ingredients which are available without PA. 3. Preferred without a PA only if patient on a diabetic therapy or prior ACE therapy.	Per best practices patient should have trialed prior therapy of ACE inhibitor or currently on a diabetic therapy.
DIRECT RENIN INHIBITOR			MC/DEL MC/DEL MC/DEL		AMTURNIDE TEKTURN ¹ TEKAMLO	Use PA Form# 20420 1. Must show failure of single and combination therapy from all preferred antihypertensive categories.	
ANTIHYPERTENSIVES - CENTRAL	MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	CLONIDINE HCL TABS GUANFACINE HCL TABS HYDRALAZINE HCL TABS HYLOREL TABS METHYLDOPA TABS MINOXIDIL TABS PRAZOSIN HCL CAPS RESERPINE TABS	MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC/DEL		CLONIDINE PATCH CLONIDINE TTS GUANABENZ ACETATE TABS ISMELIN TABS MINIPRESS CAPS NEXICLON TENEX TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ACE INHIBITORS AND CA CHANNEL BLOCKERS			MC/DEL MC MC MC/DEL	8 8 8 9	AMLODIPINE/BENAZEPRIL PRESTALIA ¹ TARKA TBCR LOTREL CAPS	Use PA Form# 20420 1. Prestalia will only be approved for patients \geq 18 years of age. Use individual preferred generic medications.	
ACE AND THIAZIDE COMBO'S	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	BENAZEPRIL HCL/HYDROCHLOR CAPTOPRIL/HYDROCHLOROTHIA ENALAPRIL MALEATE/HCTZ TABS LISINOPRIL-HCTZ TABS LOTENSIN HCT TABS	MC/DEL MC MC/DEL MC/DEL MC/DEL		ACCURETIC TABS MONOPRIL HCT TABS PRINZIDE TABS UNIRETIC TABS ZESTORETIC TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
BETA BLOCKERS AND DIURETIC COMBO'S	MC/DEL MC/DEL MC/DEL	ATENOLOL/CHLORTHALIDONE BISOPROLOL FUMARATE/HCTZ PROPRANOLOL/HCTZ	MC/DEL MC/DEL MC MC MC/DEL		CORZIDE TABS LOPRESSOR HCT TABS TENORETIC TIMOLIDE 10/25 TABS ZIAC TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ARB'S AND CA CHANNEL BLOCKERS	MC/DEL MC/DEL MC/DEL	AMLODIPINE/VALSARTAN AMLODIPINE/VALSARTAN HCT TRIBENZOR	MC/DEL MC MC/DEL MC/DEL		AZOR BYVALSON EXFORGE EXFORGE HCT	Use PA Form# 20420	DDI: Byvalson will be non-preferred and require a prior authorization if it is currently being used in combination with drugs known to be significant CYP2D6 inhibitors (e.g. quinidine, propafenone, fluoxetine, paroxetine). Per best practices, patient should have trialed prior therapy of ACE inhibitor or currently on a diabetic therapy.
ARB'S AND DIURETICS	MC/DEL MC/DEL MC/DEL MC/DEL	BENICAR HCT ¹ LOSARTAN HCT ¹ MICARDIS HCT TABS ¹ VALSARTAN-HCT ¹	MC/DEL MC/DEL MC MC/DEL MC/DEL MC	7 8 8 8 8 8	IRBESARTAN HYDROCHLOROTHIAZIDE ATACAND HCT TABS AVALIDE TABS ¹ DIOVAN HCT TABS ¹ HYZAAR TABS TEVETEN HCT TABS	Use PA Form# 20420 1. Dosing limits apply, see Dose Consolidation List.	Per best practices, patient should have trialed prior therapy of ACE inhibitor or currently on a diabetic therapy.
ANGIOTENSIN MODULATORS-ARB COMBINATION	MC	ENTRESTO	MC/DEL MC		EDARBYCLOR ENTRESTO SPRINKLES	Use PA Form# 20420	
ARB'S AND DIRECT RENIN INHIBITOR COMBINATION			MC/DEL		VALTURNA	Use PA Form# 20420	

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
CHOLESTEROL - HMG COA + ABSORB INHIBITORS LESS POTENT DRUGS/- COMBINATIONS	MC/DEL MC/DEL MC/DEL	EZETIMIBE TABS LOVASTATIN TABS ² PRAVASTATIN ²	MC MC/DEL MC/DEL MC MC/DEL MC MC MC/DEL MC/DEL MC	8 8 8 8 8 8 8 8 8 8 8	ALTOPREV TB24 FLUVASTATIN TAB ER LESCOL XL TB24 LIVALO MEVACOR TABS NEXLETOL NEXLIZET PRAVACHOL TABS PRAVIGARD ZETIA TABS	Use PA Form# 20420 2. Dosing limits apply, please see Dosage Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Zetia will be approved for patients unable to tolerate all other therapies or unable to achieve cholesterol goal with maximally tolerated dose of most potent statins. DDI: Lescol will now be non-preferred and require prior authorization if it is currently being used in combination with diclofenac. DDI: Lovastatin (doses greater than 40mg/day) will now be non-preferred and require prior authorization if it is currently being used in combination with amiodarone. DDI: Lovastatin (doses greater than 20mg per day) will now be non-preferred and require prior authorization if it is currently being used in combination cyclosporine. DDI: All preferred statins will now be non-preferred and require prior authorization if it is currently being used in combination with gemfibrozil.
CHOLESTEROL - HMG COA + ABSORB INHIBITORS STATIN/ NIACIN COMBO	MC	SIMCOR	MC		ADVICOR TBCR	Use PA Form# 20420	
FAMILIAL HYPERCHOLESTEROLEMIA	MC MC	PRALUENT (LABLER 72733) PEN ^{1,2,3,5} REPATHA ^{1,2,3}	MC MC MC MC		EVKEEZA ^{1,4} JUXTAPI ¹ KYNAMRO ¹ LEQVIO	Use PA Form# 20420 1. Clinical PA required for appropriate diagnosis. 2. Quantity limits apply. 3. Documented adherence to lipid lowering medications and abstinence from tobacco for previous 90 days. 4. For treatment of patients \geq 12 years of age. 5. Approval of Praluent NDC's with labeler code 00024 will be considered only if labeler code 72733 NDC's are on a long-term backorder and unavailable from the manufacturer.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Juxtapid is contraindicated with strong CYP3A4 inhibitors. Juxtapid dosage should not exceed 30mg daily when it is used concomitantly with weak CYP3A4 inhibitors. Kynamro requires an appropriate lab testing prior to starting (ALT <ast), alkaline="" and="" bilirubin,="" every="" first="" for="" liver-related="" monthly="" months.<br="" phosphatase="" tests="" the="" then="" three="" total="" year,=""></ast),> Repatha and Praluent Criteria for approval: The patient's age is FDA approved for the given indication AND • Concurrent use with statin therapy AND • Documented adherence to prescribed lipid lowering medications for the previous 90 days AND • Recommended or prescribed by a lipidologist or cardiologist AND • Inability to reach goal LDL-C despite a trial of 2 or more maximum tolerated dose of statins (one of which must be atorvastatin or rosuvastatin) and ezetimibe 10mg daily. Additional criteria for the diagnosis of heterozygous familial hypercholesterolemia (HeFH): (both are required): Total cholesterol > 290 mg/dL OR LDL-C > 190 mg/dL AND one of the following • Presence of tendon xanthomas OR • In 1st or 2nd degree relative-documented tendon xanthomas, MI at age \leq 60 years or TC > 290 mg/dL. Additional criteria for the diagnosis of clinical atherosclerotic cardiovascular disease: History of MI, angina, coronary or other arterial revascularization, stroke, TIA, or PVD of atherosclerotic origin. Additional criteria for the diagnosis of homozygous familial hypercholesterolemia (Repatha only): Total cholesterol levels > 290mg/dL or LDL-C > 190mg/dL (adults) OR Total cholesterol levels > 260mg/dL or LDL-C > 155mg/dL (children $<$ 16 years) and TG within reference range OR Confirmation of diagnosis by gene testing.
FAMILIAL HYPERCHOLESTEROLEMIA AND HYPERTRIGLYCERIDEMIA					TRYNGOLZA	Use PA Form# 20420	Tryngolza requires fasting triglycerides of \geq 880 mg/dL and confirmed genetically identified familial chylomicronemia syndrome (FCS).
HYPERPHAGIA - MISC							
HYPERPHAGIA - MISC			MC	8	VYKAT XR		FDA approved for the treatment of hyperphagia in adults and pediatric patients 4 years of age and older with Prader-Willi syndrome (PWS).
PULMONARY ANTI-HYPERTENSIVES							
PULMONARY ANTI-HYPERTENSIVES	MC MC/DEL MC/DEL	EPOPROSTENOL INJ ³ SILDENAFIL TADALAFIL	MC/DEL MC MC/DEL MC MC MC MC MC/DEL MC MC MC MC MC MC MC MC MC/DEL MC		ADEMPAS ^{1,3} ADCIRCA ⁴ ALYQ TAB FLOLAN ³ LIQREV OPSUMIT ^{1,2} OPSYNVI ⁴ ORENITRAM REMODULIN ³ REVATIO ⁴ TADLIQ ⁴ TYVASO UPTRAVI VELVETRI ³ WINREVAIR ⁴ YUTREPIA	Use PA Form# 20420 1. Requires previous trials/failure of multiple preferred medications. 2. Dosing limits apply, see the Dose Consolidation List. 3. Require WHO Group 1 diagnosis of primary PAH (Primary Pulmonary Hypertension) and NYHA functional class 3 or 4. 4. Require WHO Group 1 diagnosis of primary PAH (Primary Pulmonary Hypertension) and NYHA functional class 2 or 3.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Sildenafil will be preferred with clinical PA for treatment of pulmonary arterial hypertension (WHO Group 1) in adults to improve exercise ability and delay clinical worsening. Avoid concomitant use of sildenafil with moderate or strong Cyp3A inhibitors. DDI: Uptravi will require a prior authorization if it is currently being used in combination with strong inhibitors of CYP2C8 (gemfibrozil). DDI: Opsumit will require a prior authorization if it is currently being used in combination with drugs known to be significant CYP3A inhibitors (ketoconazole, itraconazole, clarithromycin, indinavir, nefazodone, neflunavir, ritonavir, atazanavir, saquinavir and telithromycin). DDI: Adempas will require a prior authorization if it is currently being used in combination with drugs known to be PDE inhibitors should be avoided (including dipyridamole, adcirca and tadalafil) with adempas. Liqrev: treatment of pulmonary arterial hypertension (WHO Group 1) in adults to improve exercise ability and delay clinical worsening. Avoid concomitant use of liqrev with moderate or strong CYP3A inhibitors.

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
ALLERGY / ASTHMA THERAPIES							
ANAPHYLACTIC DEVICES	MC MC/DEL MC/DEL MC/DEL	AUVI-Q EPINEPHRINE EPIPEN EPIPEN JR	MC MC		NEFFY TWINJECT	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ALLERGEN IMMUNOTHERAPY			MC MC MC MC MC		ODACTRA ORALAIR ¹ PALFORZIA RAGWITEK GRASTEK	Use PA Form# 20420 1. See criteria section	Prescriber must provide the testing to show that the patient is allergic to the components in the prescribed therapy and must provide a clinically valid rationale why single agent sublingual therapy is being chosen over subcutaneous therapy. Palforzia is approved for use in patients with a confirmed diagnosis of peanut allergy. Initial dose escalation may be administered to patients aged 4 through 17 years. Up-dosing and maintenance may be continued in patients 4 years of age and older. Odactra is approved for use in persons 12 through 65 years of age. Note that Odactra is not indicated for the immediate relief of allergic symptoms. Treatment must start 12 weeks before expected onset of pollen season and only after confirmed by positive skin test or in vitro testing for pollen-specific IgE antibodies for any of the 5 grass species contained in Oralair. Oralair: Patient age ≥10 years and ≤65 years. Have an auto-injectable epinephrine on-hand.
ANTIASTHMATIC - ANTICHOLINERGICS - INHALER	MC MC/DEL MC/DEL	INCRUSE ELLIPTA ³ SPIRIVA HANDIHALER ^{1,2} SPIRIVA RESPIMAT	MC MC/DEL		LONHALA MAGNAIR TUDORZA	Use PA Form# 20420 1. Quantity limit of 1 inhalation daily (1 capsule for inhalation daily). Spiriva will require PA if Combivent or Atrovent nebulizer solution is in member's current drug profile. 2. We ask physicians to write "asthma" on the prescription whenever Spiriva is primarily being used for that condition. 3. Quantity limit of 1 inhalation daily.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIASTHMATIC - Dipeptidyl Peptidase 1 Inhibitors	MC	BRINSUPRI				Use PA Form# 20420 1. Clinical PA is required to establish diagnosis and medical necessity.	BRINSUPRI required criteria include: <ul style="list-style-type: none">• Imaging confirming bronchiectasis and no overlapping asthma/COPD required.• Documented airway clearance.• Greater than 2 exacerbations requiring antibiotic therapy in the last 12 months.• Must be approved by pulmonologist.
ANTIASTHMATIC - PHOSPHODIESTERASE 4 INHIBITORS	MC/DEL	ROFLUMILAST	MC/DEL MC		DALIRESP OHTUVAYRE ¹	Use PA Form# 20420 1. For the maintenance treatment of chronic obstructive pulmonary disease (COPD) in adult patients.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIASTHMATIC - ANTICHOLINERGICS - NEBULIZER	MC/DEL	IPRATROPIUM BROMIDE SOLN	MC MC/DEL		ATROVENT SOLN YUPELRI	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIASTHMATIC - ANTIINFLAMMATORY AGENTS	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	CROMOLYN SODIUM NEBU DUPIXENT ^{2,4} FASENRA ⁶ FASENRA ⁶ AUTO INJCT XOLAIR ^{1,4}	MC MC MC MC MC	8 8 8 8	CINQAIR ³ NUCALA ² RHAPSIDO ⁴ TEZSPIRE ⁵	Use PA Form# 20420 1. Need max inhaled steroids and written by pulmonary or allergy specialist. Must have elevated IgE and ≥ age 6. 2. For patients with severe asthma aged 12 years or older and eosinophilia. 3. For patients ≥ 18 years of age with eosinophilia. 4. Clinical PA required to establish diagnosis and medical necessity. 5. For adult and pediatric patients aged 12 years and older with severe asthma. 6. For patients ≥ 6 years of age for eosinophilia.	All will require suboptimal response to maximal doses of inhaled steroid as evidenced by asthmatic ER/Hospital admissions and Allergy/Pulmonary specialist management. Dupixent limited to patient with asthma not controlled on high dose ICS-LABA who have eosinophil greater than or equal to 150 cells or the patient is depend on an oral corticosteroid. Fasenra , Nucala and Cinqair are not indicated for treatment of other eosinophilic conditions and are not indicated for the relief of acute bronchospasm or status asthmaticus. Rhapsido for Chronic Spontaneous Urticaria - must have had an inadequate clinical response of at least 14-days with at least two different second-generation antihistamines at 4 times standard dose. Must continue use of second-generation antihistamine. Must be prescribed by or in consultation with either allergist-immunologist, dermatologist, pulmonologist, or otolaryngologist.
ANTIASTHMATIC - NASAL STEROIDS	MC/DEL MC/DEL MC MC/DEL MC/DEL MC	BUDESONIDE SPRAY FLUTICASONE SPR ³ OLOPATADINE SPRAY OMNARIS SPR ³ TRIAMCINOLONE NS QNASL	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC	8 8 8 8 8 8	DYMISTA FLONASE SUSP ^{2,3} FLUNISOLIDE SOLN ^{1,3} NASONEX SUSP RHINOCORT AERO ^{2,3} RHINOCORT AQUA SUSP ^{2,3} RYALTRIS ⁴ TRI-NASAL SOLN ^{2,3}	Use PA Form# 20420 1. All preferred drugs must be tried before moving to non preferred steps. 2. All step 5 medications need to be tried before moving to step 8's. 3. Dosing limits apply to whole category, see Dosage Consolidation List.	Preferred drugs and step therapy must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Xhance will be considered for the treatment of nasal polyps in patients 18 years of age or older. The patient has had a documented side effect, allergy, or treatment failure of two preferred nasal glucocorticoids, one of which must be fluticasone.

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
			MC MC/DEL MC MC/DEL	8 8 8 8	VANCENASE POCKETHALER AERS ^{2,3} VERAMYST ^{2,3} XHANCE ² ZETONNA ³	4. Use of individual ingredients or other preferred agents.	
ANTIASTHMATIC - NASAL MISC.	MC/DEL MC/DEL MC	AZELASTINE CROMOLYN NASAL 4% IPRATROPIUM NASAL SOL ¹	MC/DEL MC/DEL	8 8	ASTEPRO ² PATANASE	Use PA Form# 20420 1. Ipratropium will be approved if submitted with documentation supporting use of CPAP machine. 2. Utilize Multiple preferred, as well as step therapy Azelastine.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Approved if patient fails on nonsedating antihistamines and steroid nasal sprays.
ANTIASTHMATIC - BETA - ADRENERGICS	MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC MC/DEL MC MC/DEL MC/DEL MC/DEL MC	ALBUTEROL 0.63mg/3ml ALBUTEROL HFA ALBUTEROL NEB LEVALBUTEROL TARTRATE METAPROTERENOL PROAIR DIGIHALER ⁴ PROAIR RESPICLICK PROVENTIL HFA SEREVENT STRIVERDI TERBUTALINE SULFATE TABS VENTOLIN HFA AERS	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC MC MC MC MC/DEL MC/DEL	8 8 8 8 4 8 8	ACCUNEB NEBU AIRSUPRA ALBUTEROL HFA (labeler 66993001968) BRETHINE VOLMAX TBCR VOSPIRE ER TB12 XOPENEX HFA ³ XOPENEX NEBU ^{1,2}	Use PA Form# 20420 1. Xopenex users w/ prior asthma hospitalization due to albuterol nebulizer failure will be grandfathered. 2. Quantity Limit: 12 cc/day.. 3. Dosing limits apply, see Dosage Consolidation List. 4. For the treatment of patients \geq 4 years of age.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. AIRSUPRA has new PA criteria that include the patient is aged \geq 18, AND the patient has had a documented side effect or allergy, AND treatment failure/intolerance or contraindication to Symbicort® and Dulera® SMART therapy, AND the patient is unable to use albuterol and budesonide separately.
ANTIASTHMATIC - ADRENERGIC COMBINATIONS	MC MC MC MC MC/DEL MC/DEL	ADVAIR DISKUS ¹ ADVAIR HFA ¹ AIRDUO RESPICLICK ² BREO ELLIPTA ¹ DULERA FLUTICASONE-SALMETEROL SYMBICORT	MC MC/DEL MC/DEL MC	8 8 8 8	AIRDUO DIGIHALER ² BREYNA BREZTRI AEROSPHERE TRELEGY ELLIPTA ¹	Use PA Form# 20420 1. Dosing limits apply, see Dosage Consolidation List. 2. For patients \geq 12 years and older.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. AirDuo® Respclick be non-preferred and require prior authorization and be available to those who are unable to tolerate or who have failed on preferred medications. DDI: Avoid concomitant use of strong CYP3A4 inhibitors (e.g. ritonavir, atazanavir, clarithromycin, indinavir, itraconazole, nefazodone, neflifavir, saquinavir, ketoconazole, telithromycin) with AirDuo® Respclick is not recommended due to increased systemic corticosteroid and increased cardiovascular adverse effects.
ANTIASTHMATIC - ADRENERGIC ANTICHOLINERGIC	MC/DEL MC MC/DEL MC/DEL	ALBUTEROL/IPRATROPIUM NEB. SOLN ANORO ELLIPTA COMBIVENT RESPIMAT STIOLTO	MC/DEL MC/DEL MC/DEL		BEVESPI AEROSPHERE ^{2,3} DUAKLIR PRESSAIR DUONEB SOLN ¹	Use PA Form# 20420 1. Please use preferred individual ingredients Albuterol and Ipratropium. 2. Dosing limits apply, see Dosing Consolidation List. 3. The safety and efficacy of use in children under the age of 18 years have not been established.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Bevespi should be used with extreme caution in patients being treated with MAO inhibitors, TCAs, or other drugs known to prolong the QTc interval. DDI: Avoid concomitant use of Bevespi with other anticholinergic-containing drugs, due to an increased risk of anticholinergic adverse events. Bevespi should be used with extreme caution in patients being treated with MAO inhibitors, TCAs, or other drugs known to prolong the QTc interval.
ANTIASTHMATIC - XANTHINES	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	AMINOPHYLLINE TABS THEOCHRON TB12 THEOLAIR-SR TB12 THEOPHYLLINE CR TB12 THEOPHYLLINE ELIX THEOPHYLLINE SOLN THEOPHYLLINE ER CP12 THEOPHYLLINE ER TB12	MC/DEL MC MC/DEL		THEO-24 CP24 THEOLAIR TABS UNIPHYL TBCR	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIASTHMATIC - STEROID INHALANTS	MC MC/DEL MC/DEL MC/DEL MC/DEL MC	ARNUITY ELLIPTA ASMANEX TWISTHALER ^{3,4} ASMANEX HFA BUDESONIDE NEB 0.25MG & 0.5MG ¹ PULMICORT FLEXHALER ³ QVAR AERS ³	MC MC/DEL MC MC/DEL MC/DEL	8 8 8 8 8	AEROSPA ALVESCO ³ ARMONAIR DIGIHALER BUDESONIDE NEB 1MG PULMICORT SUSP	Use PA Form# 20420 1. Budesonide Neb 0.25mg & 0.5mg will be preferred for members under the age of 8 years old. PA will be required for members 8 years of age and older, please consider other preferred options. 2. All preferred must be tried before moving to non preferred steps. 3. Dosing limits apply, see Dosage Consolidation List. 4. Asmanex 110mcg will be limited to member between the ages of 4-11 years old.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
ANTIASTHMATIC - 5-Lipoxygenase Inhibitors			MC		ZYFLO CR TABS	Use PA Form# 20420	Other Preferred asthma controller drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANTIASTHMATIC - LEUKOTRIENE RECEPTOR ANTAGONISTS	MC/DEL MC/DEL MC/DEL	MONTELUKAST GRANULE ¹ MONTELUKAST SODIUM TAB MONTELUKAST SODIUM CHEW TAB	MC/DEL MC/DEL MC/DEL	8 8 8	ACCOLATE TABS SINGULAIR ² SINGULAIR GRANULES	Use PA Form# 20420 1. Montelukast Granules will only be approved if between ages of 6 - 24 months. 2. Singulair Chewable 4mg from 2 years- 5 years and Singulair Chewable 5mgs from 6 years- 14 years old.	
ANTIASTHMATIC - ALPHA-PROTEINASE INHIBITOR			MC MC/DEL MC MC	8 8 8 8	ARALAST ZEMAIRA GLASSIA PROLASTIN SUSR	Use PA Form# 20420	Prolastin and Azemaira will be approved for members with A1AT deficiency and clinically demonstrable panacinar emphysema.
ANTIASTHMATIC - HYDRO-LYTIC ENZYMES			MC/DEL		PULMOZYME SOLN	Use PA Form# 20420	Will be approved for cystic fibrosis patients.
ANTIASTHMATIC - MUCOLYTICS	MC/DEL	ACETYLCYSTEINE ¹	MC		MUCOMYST	Use PA Form# 20420 1. Acetylcysteine is covered with diagnosis of CF.	
ANTIASTHMATIC-CFTR POTENTIATOR AND COMBINATIONS			MC MC MC MC MC MC/DEL		ALYFTREK BRONCHITOL ¹ KALYDECO ORKAMBI SYMDEKO TRIKAFTA	Use PA Form# 20420 1. For the treatment of patients \geq 18 years of age with CF.	Alfytrek will be considered for the treatment of patients 6 years and older with at least one responsive mutation, including 31 additional mutations not responsive to other CFTR modulator therapies. Bronchitol will be considered as add-on maintenance therapy to improve pulmonary function in adult patients 18 years and older with cystic fibrosis (CF). Use Bronchitol only for adults who have passed the Bronchitol Tolerance Test (BTT). (see Recommended Dosage section for further information). Kalydeco will be considered for patients with cystic fibrosis (CF) aged 1 month and older who have at least one mutation in the CFTR gene that is responsive to ivacaftor potentiation based on clinical and/or in vitro assay data. If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to detect the presence of a CFTR mutation followed by verification with bi-directional sequencing when recommended by the mutation test instructions for use. Orkambi will be considered for patients with cystic fibrosis (CF) aged 1 year and older who are homozygous for the F508del mutation in the CFTR gene. If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to detect the presence of the F508del mutation on both alleles of the CFTR gene. The efficacy and safety of Orkambi have not been established in patients with CF other than those homozygous for the F508del mutation. Symdeko will be considered for patients with cystic fibrosis (CF) aged 6 years and older who are homozygous for the F508del mutation or who have at least one mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to tezacaftor/ivacaftor based on in vitro data and/or clinical evidence. If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to detect the presence of a CFTR mutation followed by verification with bi-directional sequencing when recommended by the mutation test instructions for use. Trikafta will be considered for the treatment of cystic fibrosis (CF) in patients aged 2 years and older who have at least one F508del mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene or mutation in the CFTR gene that is responsive based on in vitro data. If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to confirm the presence of at least one F508del mutation or a mutation that is responsive based on in vitro data.
IDIOPATHIC PULMONARY FIBROSIS	MC/DEL	OFEV ¹ PIRFENIDONE	MC MC/DEL	8 8	ESBRIET ¹ JASCAYD ¹	Use PA Form# 20420 1. Clinical PA is required to establish diagnosis and medical necessity.	Ofev- Avoid concomitant use with P-gp and CYPA4 inducers (e.g. carbamazepine, phenytoin, and St. John's wort). Esbriet- The concomitant use with strong CYP1A2 inhibitors (e.g. fluvoxamine, enoxacin) is not recommended.
COUGH/COLD							
COUGH/COLD	MC/DEL MC/DEL MC/DEL MC/DEL MC MC	DEXTROMETHORPHAN CAPS ¹ DEXTRO-GUAIF SYRP ¹ GUAIFENESIN SYRP ¹ PSEUDOEPHEDRINE ¹ ROBITUSSIN DM SYRP ¹ ROBITUSSIN SUGAR FREE SYRP ¹				Use PA Form# 20420 1. All of cough cold preparations are not covered except these preferred products.	All non-preferred products are not covered as permitted by Federal Medicaid regulations and MaineCare Policy.
DIGESTIVE AIDS / ASSORTED GI							
GI - ANTIPERISTALTIC AGENTS	MC/DEL MC/DEL MC/DEL MC/DEL MC	DIPHENOXYLATE DIPHENOXYLATE/ATROPINE LOPERAMIDE HCL CAPS/LIQ OPIUM TINCTURE TINC PAREGORIC TINC	MC/DEL MC MC		LOFENE TABS LONOX TABS MOTOFEN TABS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval.

Category	Coverage Indicator	Preferred Drugs		Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
ANTI-CATAPLECTIC AGENTS								
PSYCHOTHERAPEUTIC AGENTS - MISC.				MC MC		NUDEXTA XENAZINE	Use PA Form# 20710 for Xenazine	
WEIGHT LOSS								
WEIGHT LOSS						No longer covered: Phentermine, Xenical, Didrex, and Meridia	Weight loss drugs are not covered as permitted by Federal Medicaid regulations and Maine Medicaid (MaineCare) Policy.	
ALZHEIMER DISEASE								
ALZHEIMER - Cholinomimetics/Others	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	DONEPEZIL HYDROCHLORIDE TABS ¹ DONEPEZIL HYDROCHLORIDE ODT ¹ EXELON DIS ¹ GALANTAMINE CAPS ¹ GALANTAMINE TAB ¹ MEMANTINE ¹ RIVASTIGMINE TARTRATE CAPS ¹		MC MC MC/DEL MC MC/DEL MC/DEL MC/DEL	6 6 7 8 8 8 8	ARICEPT TABS ² ARICEPT ODT ² DONEPEZIL HYDROCHLORIDE TABS 23MG ADLARITY ³ EXELON CAP GALANTAMINE HYDROBROMIDE SOL KISUNLA ¹	Use PA Form# 20420 1. PA is required to establish dementia diagnosis and baseline mental status score. 2. Must fail all preferred products before moving to non-preferred. 3. Approvals will require trials and failure or clinical rationale why preferred patches can't be used.	Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Kisunla and Leqembi: Testing to rule out reversible causes of dementia (CBC, CMP, TSH, B12, urine drug screen, RPR/VDRL, (folate if alcohol abuse is present), HIV (if risk present) and an assessment including a review of current medications as a cause of intellectual decline - Prescribed by or in consultation with a neurologist or geriatrician or geriatric psychiatrist. Diagnosis of Alzheimer's disease defined as: <ul style="list-style-type: none">Confirmed presence of amyloid pathology and mild cognitive impairment or mild dementia stage of disease, consistent with Stage 3 and Stage 4 Alzheimer's disease ORConfirmed presence of amyloid pathology and prodromal or mild dementia stage of disease, consistent with Stage 3 & Stage 4 Alzheimer's disease Testing: <ul style="list-style-type: none">Clinical Dementia Rating (CDR) global score of 0.5 or 1.0 ORRepeatable Battery for Assessment of Neuropsychological Status (RBANS) delayed memory index score ≤ 85 ORMini-Mental State Examination (MMSE) score of 20-30 ORMontreal Cognitive Assessment (MoCA) score ≤ 22 <ul style="list-style-type: none">Member is age 50 or olderObtain recent (within one year) brain magnetic resonance imaging (MRI) prior to initiating treatmentProvider attestation to obtain MRIs prior to the 7th infusion (first dose of 10 mg/kg) and 12th infusion (sixth dose of 10 mg/kg)Member does NOT have history or increased risk of amyloid related imaging abnormalities-edema (ARIA-E), which includes brain edema or sulcal effusions and amyloid related imaging abnormalities hemosiderin deposition (ARIA-H), which includes microhemorrhage and superficial siderosisMember does NOT have hypersensitivity to any components of these drugs
SMOKING CESSATION								
NICOTINE PATCHES / TABLETS	MC/DEL MC/DEL MC/DEL MC/DEL	CHANTIX TAB ¹ CHANTIX STARTER PACK NICOTINE DIS PT24 ¹ VARENICLINE TAB		MC/DEL		NICODERM CQ PT24 ¹	Use PA Form# 20420 1. See criteria section for exemptions	As of July 1, 2014 per MaineCare policy, smoking cessation products will be covered without a copay(including MEDEL). No annual or lifetime limits, must follow FDA approved indications and therapy guidelines. Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Note: MaineCare policy, smoking cessation product were "not covered" except for during pregnancy between 9/1/12 and 1/1/14, between 1/1/2014 and 7/1/14 smoking cessation products were covered with limitations. Patients may qualify for the medication through The Maine Tobacco Helpline if they do not have MaineCare or MEDEL. Patients are encouraged to call The Maine Tobacco helpline at 1-800-207-1230.
NICOTINE REPLACEMENT - OTHER	MC/DEL MC/DEL MC/DEL	NICOTINE POLACRILEX GUM ¹ NICOTINE LOZENGE MINI NICOTINE LOZENGE		MC/DEL MC/DEL MC/DEL MC	8 8 8 8	NICOTROL INHALER ^{1,2} NICOTROL NASAL SPRAY ^{1,2} NICORETTE GUM ^{1,2} NICORETTE LOZENGES	Use PA Form# 20420 1. See criteria section for exemptions. 2. Must use non-preferred products in specified step order.	As of July 1, 2014 per MaineCare policy, smoking cessation products will be covered without a copay(including MEDEL). No annual or lifetime limits, must follow FDA approved indications and therapy guidelines. Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Note: MaineCare policy, smoking cessation product were "not covered" except for during pregnancy between 9/1/12 and 1/1/14, between 1/1/2014 and 7/1/14 smoking cessation products were covered with limitations. Patients may qualify for the medication through The Maine Tobacco Helpline if they do not have MaineCare or MEDEL. Patients are encouraged to call The Maine Tobacco helpline at 1-800-207-1230.
ALCOHOL DETERRENTS								
ALCOHOL DETERRENTS	MC/DEL MC MC MC/DEL	ACAMPROSATE ANTABUSE TABS DISULFIRAM TABS NALTREXONE HCL TABS		MC/DEL		ACAMPRO ¹	Use PA Form# 20420 1. Should only be used in conjunction with formal structured outpatient detoxification program.	Preferred generic drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
MISCELLANEOUS ANALGESICS								
ANALGESICS - MISC.	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	ACETAMINOPHEN ASPIRIN ASPRIN/ APAP/ CAFF TAB BUTAL/ASA/CAFF BUTALBITAL COMPOUND BUTALBITAL/ACET TABS		MC MC/DEL MC/DEL MC MC MC/DEL		AXOCET CAPS ESGIC-PLUS FIORICET TABS FIORINAL CAPS FIORTAL CAPS FORTABS TABS	Use PA Form# 20420 1. QL: 1. QL: No greater than 14-day supply within 90 days.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Journavx requires patient must have documented clinical reason as to why they are unable to use acetaminophen and NSAIDS (which can include Cox-II inhibitors). Journavx is FDA approved for moderate to severe ACUTE pain in adults.

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
	MC/DEL	BUTALBITAL/APAP CAPS	MC		JOURNAVX ¹		
	MC/DEL	BUTALBITAL/APAP/CAFFEINE TABS	MC		PHRENILIN TABS		
	MC/DEL	CHOLINE MAGNESIUM TRISALI	MC		PHRENILIN FORTE CAPS		
	MC/DEL	DIFLUNISAL TABS	MC		TRILISATE LIQD		
	MC	EXCEDRIN	MC		TRILISATE TABS		
	MC/DEL	SALSALATE TABS	MC		ZEBUTAL CAPS		
			MC		ZORPRIN TBCR		
LONG ACTING NARCOTICS							
NARCOTICS - LONG ACTING	MC/DEL	FENTANYL PATCH ⁴	MC	8	ARYMO ER	Use PA Form# 20510	Preferred drugs (Fentanyl Patch, Morphine Sulfate ER tab, and Butrans) must be tried for at least 2 weeks each & failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug & the preferred drug(s) exists. Adequate trials include prevention/treatment of common adverse effects associated w/ narcotics (antinausea, antipruritic, etc.) as well as adequate equianalgesic dosing when converting from one narcotic to another. Also, adequate documentation of attempts to titrate dose of preferred agents to achieve adequate pain relief & desired clinical response must be provided. Member's drug regimen for additions &/or discontinuations of medications that may affect absorption &/or metabolism of preferred agents must be monitored. Approvals will not be granted if patient had access to either non-preferred products or high doses of short acting narcotics during the trial period. Non-preferred drugs will not be approved for patients showing evidence of usage patterns consistent w/ controlled substance abuse such as:
	MC/DEL	BUTTRANS ⁴	MC	8	AVINZA	Use PA Form #10300 for PAs over the opiate limit	1. Frequent or persistent early refills of controlled drugs;
	MC/DEL	MORPHINE SULFATE ER TB12	MC	8	BELBUCA		2. Multiple instances of early refill overrides due to reports of misplacement, stolen, dropped in toilet or sink, distant travel, etc.
			MC	8	EXALGO		3. Breaches of narcotic contracts with any provider;
			MC/DEL	8	HYSINGLA ER		4. Failure to comply with patient responsibilities in attached opioid documentation (see PA form) including but not limited to failing to submit to and pass
			MC	8	KADIAN		5. Failing to take or pass random drug testing;
			MC/DEL	8	METHADONE ⁶		6. Failing to provide old records regarding prior use of narcotics;
			MC/DEL	8	METHADOSE ⁶		7. Receiving controlled substances from other prescribers that the provider submitting the PA is unaware of;
			MC/DEL	8	MORPHABOND ER		8. Documented history of substance abuse. Substance abuse evaluations may be required for patients with medical records displaying documented substance abuse or potential signs of narcotic misuse and abuse such as chronic early refills, short dosing intervals, frequent dose increases, multiple lost/stolen etc scripts and intolerance or "allergy" to all products but Oxycontin.
			MC/DEL	8	MORPHINE SULFATE ER CAP		9. Circumventing MaineCare prior authorization requirements for narcotics by paying cash for affected narcotics (prescribers failed to submit prior authorization prior to cash narcotic scripts being filled by member).
			MC/DEL	8	MORPHINE SULFATE SUPP		10. Requests for any Brand name controlled substance, considered by authorities to be highly abused and diverted (Oxycontin, Percocet, Tylox, Vicodin, Dilaudid, Ultracet...) with an available AB rated generic equivalent will be denied unless it will be provided in a setting that virtually eliminates the risk of diversion.
			MC/DEL	8	MS CONTIN TB12		11. Allergic reactions to any product within a specific narcotic class will justify and preclude use of any other product in the same class due to the risk of cross-hypersensitivity.
			MC	8	OPANA ER		Hysingla ER - Concomitant use should be avoided with mixed agonist/antagonist analgesics, partial agonist analgesics, and MAOIs. Verify prior trials and failures or intolerance of preferred treatments.
			MC/DEL	8	ORAMORPH SR TB12		Methadone - Established users must have a trial and failure of at least 2 preferred drugs for least 2 weeks. Otherwise they will be allowed 180 days to transition to a preferred product.
			MC/DEL	8	OXYCONTIN TB12 ¹		
			MC	8	XARTEMIS ER		
			MC	8	ZOHYDRO ER		
			MC	8	OXYCODONECONC		
			MC/DEL	9	OXYCODONE ER ^{3,5}		
NARCOTICS - SELECTED	MC/DEL	TRAMADOL HCL TABS 50 mg ²	MC/DEL	7	RYZOLT	Use PA Form# 20420	Preferred drugs from this and other narcotic classes must be tried for at least 2 weeks each and failed due to lack of efficacy or intolerable side effects before non-preferred drugs from this class will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Approvals will not be granted if patient had access to either non-preferred products or high doses of short acting narcotics during the trial period. Substance abuse evaluations may be required for patients with medical records displaying potential signs of narcotic misuse and abuse such as chronic early refills, short dosing intervals, frequent dose increases, multiple lost/stolen etc scripts and intolerance or "allergy" to all products but desired product. Allergic reactions to any product within a specific narcotic class will justify and preclude use of any other product in the same class due to the risk of cross-hypersensitivity.
	MC/DEL	TRAMADOL/APAP TABS	MC	8	BUPRENEX SOLN	Use PA form #10300 for PAs over the opiate limit	Non-preferred drugs will not be approved for patients showing evidence of usage patterns consistent with controlled substance abuse such as:
			MC/DEL	8	BUTORPHANOL		1. frequent or persistent early refills of controlled drugs;
			MC	8	NALBUPHINE HCL SOLN		2. multiple instances of early refill overrides due to reports of misplacement, stolen, dropped in toilet or sink, distant travel;
			MC	8	QDOLO SOLN		3. breaches of narcotic contracts with any provider;
			MC	8	SEGLENTIS ¹		4. failure to comply with patient responsibilities in attached opioid documentation (see PA form) including but not limited to failing to submit to and pass pill counts;
			MC	8	STADOL NS SOLN		5. failing to take or pass random drug testing;
			MC	8	TRAMADOL ER		6. failing to provide old records regarding prior use of narcotics;
			MC	8	ULTRACET TABS ¹		7. receiving controlled substances from other prescribers that the provider submitting the PA is unaware of. Substance abuse evaluations may be required for patients with medical records displaying potential signs of narcotic misuse and abuse such as chronic early refills, short dosing intervals, frequent dose increases, multiple lost/stolen etc scripts and intolerance or "allergy" to all products but Oxycontin. Allergic reactions to any product within a specific narcotic class will justify and preclude use of any other product in the same class due to the risk of cross-hypersensitivity.
			MC	9	ULTRAM ER		Beginning January 2017, all current opiate users who are above the maximum combined daily dose of 100 MME must titrate their total daily dose of opioid medications below 30 MME. Also, the maximum daily supply of an opiate prescription for acute pain will be limited to 7-day supplies. The maximum day supply of an opiate prescription for chronic pain will be limited to 30-day supplies. As of July 1, 2017 all users of opioid medications must comply with the maximum combined daily dose of 100 MME.
							However, for MaineCare members, effective January 1, 2017, opioid prescription(s) for more than a 7-day supply and/or more than 30 MME/ day will require a prior authorization. Please note that MaineCare implemented a 30 MME limit January 1, 2013 that is still effective.
							Post-surgical members may receive prior authorizations for opiates up to 60 days in length if medical necessity is provided by the surgical provider.
							An MME conversion chart is available at www.maineicarepd.org . Click on "General Pharmacy Info."
							Please see the Pain Management Policy tab for the complete criteria.

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
MISCELLANEOUS NARCOTICS							
NARCOTICS - MISC.	MC/DEL	ACETAMINOPHEN/CODEINE	MC/DEL	8	ABSTRAL	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Please refer to General Criteria category E.
	MC/DEL	ASPIRIN/CODEINE TABS	MC/DEL	8	APADAZ	Use PA form #10300 for PAs over the opiate limit	
	MC/DEL	BUTAL/ASA/CAFF/COD CAPS	MC/DEL	8	ASCOMP/CODEINE CAPS		
	MC	BUTALBITAL/ASPIRIN/CAFFE CAPS	MC/DEL	8	BUTALBITAL/APAP/CAFFEINE/ CAPS	1. Fentanyl OT loz (Barr) and Capital and Codeine Suspension products require PA for users over 18 years of age. PA is not required if under 18 years of age.	Beginning January 2017, all current opiate users who are above the maximum combined daily dose of 100 MME must titrate their total daily dose of opioid medications below 30 MME. Also, the maximum daily supply of an opiate prescription for acute pain will be limited to 7-day supplies. The maximum day supply of an opiate prescription for chronic pain will be limited to 30-day supplies. As of July 1, 2017 all users of opioid medications must comply with the maximum combined daily dose of 100 MME.
	MC	CAPITAL AND CODEINE SUSP ¹	MC/DEL	8	BUTALBITAL COMPOUND- CODEINE CAP		
	MC/DEL	CODEINE PHOSPHATE SOLN	MC	8	DEMEROL		
	MC/DEL	CODEINE SULFATE TABS	MC/DEL	8	DILAUDID		
	MC/DEL	ENDOCET TABS ³	MC	8	DILAUDID-HP SOLN		
	MC/DEL	ENDODAN TABS	MC	8	FENTANYL CITRATE SOLN		
	MC/DEL	FENTANYL OT LOZ ¹	MC/DEL	8	FENTORA		
	MC/DEL	HYDROCODONE/ACETAMINOPHEN	MC/DEL	8	FIORICET/CODEINE CAPS		
	MC/DEL	HYDROMORPHONE HCL ³	MC	8	FIORINAL/CODEINE #3 CAPS		
	MC	LORTAB ELX	MC	8	FIORTAL/CODEINE CAPS		
	MC/DEL	MEPERIDINE SOL	MC/DEL	8	HYDROCODONE/IBUPROFEN		
	MC/DEL	OXYCODONE TAB	MC/DEL	8	HYDROMORPHONE ER		
	MC/DEL	OXYCODONE/ACETAMINOPHEN ^{2,3}	MC/DEL	8	HYDROMORPHONE RECTAL SUPP		
	MC/DEL	ROXICET	MC	8	IBUDONE		
	MC	ROXIPRIN TABS	MC/DEL	8	LEVORPHANOL TARTRATE TAB		
			MC/DEL	8	LORCET		
			MC	8	LORTAB		
			MC	8	MAXIDONE TABS		
			MC/DEL	8	MEPERIDINE TABS		
			MC/DEL	8	NORCO TABS		
			MC/DEL	8	ONSOLIS		
			MC/DEL	8	OXECTA		
			MC/DEL	8	OXYCODONE CAP		
			MC/DEL	8	OXYCODONE/APAP 10/650		
			MC/DEL	8	OXYCODONE/APAP 7.5/500		
			MC/DEL	8	PENTAZOCINE/ACET TABS		
			MC/DEL	8	PENTAZOCINE/NALOXONE TABS		
			MC	8	PERCOSET TABS		
			MC	8	PHRENILIN W/CAFFEINE/CODE CAPS		
			MC/DEL	8	ROXICET 5/500 TABS		
			MC	8	ROXICODONE TABS		
			MC/DEL	8	ROXYBOND		
			MC	8	SYNALGOS-DC CAPS		
			MC	8	TALACEN TABS		
			MC	8	TREZIX		
			MC	8	TYLENOL/CODEINE #3 TABS		
			MC	8	TYLOX CAPS		
			MC	8	XOLOX		
			MC	8	VICODIN		
			MC	8	VICOPROFEN TABS		
			MC	8	ZYDONE TABS		
			MC	9	ACTIQ LPOP		
			MC	9	CONZIP		
			MC	9	OPANA		
OPIOID DEPENDENCE TREATMENTS	MC	SUBOXONE FILM ²	MC/DEL		BUPRENORPHINE ¹	Use PA form #20200 for Extended Release Buprenorphine	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL	BUPRENORPHINE/NALOXONE TABS ²	MC		ZUBSOLV	Use PA Form #20100 for all others	Members will continue to be required to follow the criteria listed below:
						1-Induction period for 30 days	
						2-Max dose of 32 mg for induction	
						3-Max dose of 24 mg for maintenance	
						4-There is not more than one opioid fill in member's drug profile between current fill of Buprenorphine and a prior Buprenorphine fill within the past 90 days	
						5- Should provide evidence of monthly monitoring including random pill counts, urine drug tests and use of Maine Prescription Monitoring Program reports.	
						6- Buprenorphine monotherapy is preferred if member is pregnant and dose not > 24 mg day and pregnancy diagnosis is noted on the prescription.	

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
EXTENDED RELEASE BUPRENORPHINE	MC MC	BRIXADI ¹ SUBLOCADE ¹				Use PA form #20200 for Extended Release Buprenorphine 1. Clinical PA required.	Brixadi and Sublocade: The prescriber can attest (and medical record should document) that: -member has a documented history of opioid use disorder (OUD), -XRB is being used for the treatment of OUD (rather than pain or any other non-FDA approved indication) and -member's total daily dose of sublingual buprenorphine is less than or equal to 24 mg daily. AND at least one of the following is true: -The member's previous use of sublingual buprenorphine has included misuse, overuse, or diversion. -The member is at high risk of overdose (e.g., individuals leaving incarceration or abstinence-based treatment programs; individuals who are unsheltered; or those facing potential gaps in care due to delays in care or geographically limited treatment access). -The member has experienced significant medical complications of OUD and/or injection drug use. Occurrence should be in the last 5 years, or it should be clearly documented that the risk indicated by this infection or complication is ongoing (Examples of medical complications of OUD include: threatened the function of organs or life or limb threatening and required medical and/or surgical therapy. Examples of medical complications of injection drug use include osteomyelitis, endocarditis, renal failure, joint infection or other serious medical complications directly related to OUD.) -The member has treatment-resistant OUD, including those with ongoing illicit substance use in the context of sublingual buprenorphine treatment as documented by positive urine drug screens or other clear objective evidence, and/or further functional decline with explicit documentation of the functional decline. -The member has a significant intolerance of, or documented allergy to, sublingual buprenorphine (either buprenorphine monotherapy or buprenorphine/naloxone combination therapy) that has resulted in the patient's inability to comply with continued treatment using the sublingual product. (A true allergy is usually accompanied by rash, respiratory symptoms, or anaphylaxis. Other complaints such as bad taste, mouth tingling, etc. do not constitute evidence of allergy or significant intolerance. Formulation preference or convenience are not, in and of themselves, indications for using XRB.) -The member is in ongoing treatment with XRB and would like to continue the medication.
OPIOID WITHDRAWAL AGENTS			MC		LUCEMYRA ¹	Use PA Form#20420 1. Clinical PA for appropriate approved use and patient has documented contraindication to Clonidine.	
NARCOTIC ANTAGONISTS							
NARCOTIC - ANTAGONISTS	MC/DEL MC MC MC MC MC	NALTREXONE HCL TABS NALOXONE INJ NARCAN NS NALOXONE SPRAY OTC VIVITROL INJ ZIMHI	MC MC MC/DEL	8 8 8	OPVEE ² KLOXXADO ZURNAL ²	Use PA Form# 20420 2. For the treatment of adult and pediatric patients 12 years of age and older.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
COX 2 / NSAIDS							
COX 2 INHIBITORS - SELECTIVE / HIGHLY SELECTIVE	MC/DEL MC/DEL MC/DEL MC/DEL	CELECOXIB ^{4,5} KETOROLAC TROMETHAMINE ^{2,3,5} NABUMETONE TABS ⁵ MELOXICAM TABS ^{1,5}	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC		CELEBREX CAPS ^{4,5} MELOXICAM CAPS ⁵ MOBIC ⁵ MOBIC SUSP ⁵ RELAFEN TABS ⁵ QMIIZ ODT VIVLODEX XIFYRM ⁵	Use PA Form# 20420 1. Meloxicam and Xifir have dosing limits allowing one tablet daily of all strengths without PA. 2. Ketorolac Tromethamine is indicated for the short term (up to 5 days) management of moderately severe acute pain that requires analgesic at the opioid level in adults. Not indicated for minor or chronic pain conditions. 3. Ketorolac has dosing limits allowing 24 tablets for a 5-day supply every 30 days. 4. Dosing limits will be set at a maximum of 400mg daily. 5. The FDA has issued a Public Health Advisory warning of the potential for increased cardiovascular risk & GI bleeding with NSAID use.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
NSAIDS	MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	CHILDRENS IBUPROFEN DICLOFENAC POTASSIUM TABS DICLOFENAC SODIUM TABS DICLOFENAC SODIUM 1% GEL ¹ ETODOLAC FENOPROFEN CALCIUM TABS FLURBIPROFEN TABS IBUPROFEN INDOMETHACIN KETOPROFEN	MC MC MC MC MC/DEL MC MC MC MC MC/DEL MC/DEL MC/DEL MC/DEL		ADVIL TABS ANAPROX TABS ANAPROX DS TABS CAMBIA CATAFLAM TABS CHILDREN'S ADVIL SUSP CHILD'S IBUPROFEN SUSP CHILDREN'S MOTRIN SUSP CLINORIL TABS DAYPRO TABS	Use PA Form# 20420 The FDA has issued a Public Health Advisory warning of the potential for increased cardiovascular risk & GI bleeding with NSAID use. 1. Dosing limits apply, see Dosage Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Approvals will be granted for other requests based on failure of at least one generic NSAID from at least 3 different NSAID classes as described in the COX-II PA form. The FDA has issued a Public Health Advisory warning of the potential for increased cardiovascular risk & GI bleeding with NSAID use. DDI: Diclofenac will now be non-preferred and require prior authorization if it is currently being used in combination with lescol.

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required			Criteria
	MC/DEL	MECLOFENAMATE SODIUM CAPS	MC/DEL		DICLFENAC GEL			
	MC/DEL	NAPROSYN SUSP	MC/DEL		EC-NAPROSYN TBEC			
	MC/DEL	NAPROXEN SUSP	MC/DEL		ETODOLAC ER 600MG			
	MC/DEL	NAPROXEN TABS	MC		FELDENE CAPS			
	MC/DEL	NAPROXEN SODIUM TABS	MC/DEL		FLECTOR PATCH			
	MC/DEL	NAPROXEN SODIUM CAPS	MC/DEL		IBU-200			
	MC/DEL	NAPROXEN DR TBEC	MC		INDOCIN			
	MC/DEL	OXaprozin TABS	MC		LICART			
	MC/DEL	SULINDAC TABS	MC/DEL		LODINE			
	MC/DEL	TOLMETIN SODIUM	MC		LOFENA			
	MC/DEL	VOLTAREN GEL	MC/DEL		MOTRIN			
			MC		NALFON CAPS			
			MC/DEL		NAPRELAN TBCR			
			MC/DEL		NAPROSYN TABS			
			MC/DEL		NAPROXEN SODIUM TBCR			
			MC		PENNSAID			
			MC/DEL		PIROXICAM CAPS			
			MC		PONSTEL CAPS			
			MC		RELAFEN DS			
			MC		SB IBUPROFEN TABS			
			MC		SPRIX			
			MC		TIVORBEX			
			MC		TOLECTIN			
			MC		V-R IBUPROFEN TABS			
			MC		ZORVOLEX			
NSAID - PPI			MC		PREVACID NAPRA-PAC			
			MC/DEL		VIMOVO ¹			
						1. Use a preferred NSAID and PPI separately.		
RHEUMATOID ARTHRITIS								
RHEUMATOID ARTHRITIS	MC/DEL	ACTEMRA VIALS	MC	8	AMJEVITA			
	MC/DEL	ACTEMRA SYRINGES	MC/DEL	8	ARAVA			
	MC/DEL	ADALIMUMAB-FKJP ⁷	MC	8	AVTOZMA			
	MC	AVSOLA	MC/DEL	8	CIMZIA			
	MC/DEL	AZATHIOPRINE	MC/DEL		CYLTEZO			
	MC	ENBREL ²	MC/DEL		ENTYVIO			
	MC	ENBREL SURECLICK ²	MC		HADLIMA			
	MC	HUMIRA ^{1,2}	MC/DEL		HULIO			
	MC	KINERET SOLN	MC/DEL		HYDROXYCHLOROQUINE ²			
	MC/DEL	LEFLUNOMIDE	MC/DEL		HYRIMoz			
	MC/DEL	METHOTREXATE	MC/DEL		ILARIS ^{3,4}			
	MC	ORENICA	MC/DEL		INFLECTRA			
	MC/DEL	RINVOQ ³	MC		INFILXIMAB VIAL			
	MC	SIMLANDI ⁷	MC		JYLMV0			
	MC	SIMPONI PEN	MC/DEL		KEVZARA			
	MC	SIMPONI AUTOINJECTOR	MC		OLUMIANT			
	MC/DEL	SULFASALAZINE TABS	MC		OMVOH			
	MC	TYENNE ⁸	MC		OTREXUP			
	MC/DEL	XELJANZ ^{3,6}	MC		RASUVO ⁶			
	MC/DEL	XELJANZ XR	MC		REMICADE			
			MC/DEL		RENFLEXIS			
			MC		TOFIDENCE			
			MC		VELSIPITY			
			MC/DEL	8	XELJANZ XR SOL			
			MC		XATMEP ⁵			
			MC		YUFLYMA			
			MC		YUSIMRY			
			MC		ZYMFENTRA			
						8. See additional criteria on the RA PA form.		

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
MIGRAINE - PREVENTATIVE TREATMENT	MC MC/DEL MC/DEL MC/DEL MC/DEL	AIMOVIG ¹ AJOVY ¹ AJOVY AUTO INJECT ¹ EMGALITY SYRINGE ¹ 120mg/ml EMGALITY PEN ¹ 120mg/ml	MC MC		NURTEC ODT QULIPTA VYEPTI ²	Use PA Form# 10110 1. See criteria section. 2. Dosing limits apply, see Dosage Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Aimovig, Ajoyv and Emgality: The patient is 18 years of age or older AND patient has a diagnosis of episodic migraine (4-14 headache days per month with migraine lasting 4 hours or more) or chronic migraine (\geq 15 headache days per month, of which \geq 8 are migraine days, for at least 3 months) AND patient has failed or has a contraindication to an adequate trial (\geq 60 days) of at least 2 medications for migraine prophylaxis from at least 2 different classes.
MIGRAINE - ACUTE TREATMENT	MC MC MC	NURTEC ODT ¹ SPASTRIN TABS UBRELVY ¹	MC MC MC/DEL MC/DEL MC MC/DEL		BELCOMP-PB SUPP ELYXYB MIGRAZONE CAPS MIGERGOT SUP REYVOW ZAVZPRET	Use PA Form# 10110 1. Dosing limits apply, see Dosage Consolidation List.	Nurtec ODT will be preferred after 2 adequate trials of at least two preferred triptans. Reyvow is non-preferred and is indicated for the acute treatment of migraine with or without aura in adults. Reyvow is not indicated for the preventive treatment of migraine. Ubrelvy is preferred after 2 adequate trials of at least two preferred triptans for the acute treatment of migraine with or without aura in adults. It is not indicated for migraine prevention. Zavzpret: The patient must have a documented side effect, allergy, or treatment failure to preferred oral CGRP Inhibitor and two non-preferred oral CGRP Inhibitors.
GOUT							
GOUT	MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL	ALLOPURINOL TABS COLCHICINE TAB FEBUXOSTAT TAB MITIGARE PROBENECID TABS PROBENECID/COLCHICINE TABS	MC/DEL MC MC MC/DEL MC		COLCHICINE CAP COLCRYS GLOPERBA ULORIC ¹ ZYLOPRIM TABS	Use PA Form# 20420 1. Failure of therapeutic (300mg) dose of Allopurinol (failure define as not being able to get uric acid levels below 6mg/dl) or severe renal disease.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: The concomitant use of Gloperba and CYP3A4 inhibitors (e.g. clarithromycin, ketoconazole, grapefruit juice, erythromycin, verapamil, etc.) should be avoided due to the potential for serious and life-threatening toxicity.
MISC.							
ACID SPHINGOMYELINASE DEFICIENCY (ASMD)			MC		XENPOZYME ^{1,2}	Use PA Form# 20420 1. For treatment of non-central nervous system manifestations of acid sphingomyelinase deficiency (ASMD) in adult and pediatric patients. 2. Clinical PA required for appropriate diagnosis and clinical parameters.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
ANESTHETICS - MISC.	MC MC MC	BUPIVACAINE HCL SOLN LIDOCAINE HCL SOLN MARCaine SOLN	MC MC/DEL MC		SENSORCAINE-MPF SOLN SYNVISC INJ XYLOCAINE SOLN	Use PA Form# 30130	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
COLD AGGLUTININ DISEASE (CAD)			MC		ENJAYMO ¹	Use PA Form# 20420 1. Indicated to decrease the need for red blood cell transfusion due to hemolysis in adults with cold agglutinin disease (CAD).	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
CONGENITAL ADRENAL HYPERPLASIA			MC		CRENESSITY	Use PA Form# 30130	Creinessity - As adjunctive treatment to glucocorticoid replacement to control androgens in adults and pediatric patients 4 years of age and older with classic congenital adrenal hyperplasia (CAH).
PRIMARY HYPEROXALURIA TYPE 1 (PH1)			MC MC/DEL		OXLUMO ¹ RIVFLOZA	Use PA Form# 20420 1. PA is required to establish diagnosis and medical necessity.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Rivfloza: The patient has a diagnosis of Primary Hyperoxaluria Type I (PH1) confirmed via genetic testing (identification of alanine: glyoxylate aminotransferase gene (AGXT) mutation) AND urinary oxalate excretion $> 0.5\text{mmol}/1.73\text{ m}^2$ or urinary oxalate: creatinine ratio is above the upper limit of normal for age AND is at least 9 years of age AND medication is being prescribed by, or in consultation, with a nephrologist or urologist.
SICKLE CELL DISEASE	MC MC MC/DEL MC	DROXIA CASGEVY ^{2,3} HYDROXYUREA LYFGENIA ^{2,3}	MC MC MC/DEL MC		ADAKVEO ENDARI ¹ SIKLOS XROMI	Use PA Form# 20420 1. Evidence of other preferred L-glutamine products utilization and reason for failure. 2. For the treatment of patients ≥ 12 years of age. 3. PA required to confirm FDA approved	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
HUTCHINSON- GILFORD PROGERIA SYNDROME (HGPS)			MC		ZOKINVY ^{1,2}	Use PA Form# 20420 1. In patients 12 months of age and older with a body surface area (BSA) of 0.39m ² and above. 2. PA required to confirm FDA approved indication.	Zokinvy: To reduce the risk of mortality in Hutchinson-Gilford Progeria Syndrome (HGPS). For the treatment of processing-deficient Progeroid Laminopathies with either: Heterozygous LMNA mutation with progerin-like protein accumulation OR Homozygous or compound heterozygous ZMPSTE24 mutations.
OBSTRUCTIVE SLEEP APNEA			MC		ZEPBOUND	Use PA Form# 20420	Zepbound for adults with a BMI $\geq 30\text{ mg/kg}^2$ and diagnosis of moderate to severe OSA, confirmed by sleep study within the last 3 years documenting AHI ≥ 15 , AND in which CPAP is ineffective (AHI > 5 during therapeutic section of sleep study) or patient is unable to tolerate CPAP for at least 90 days AND for whom lifestyle modifications have been attempted for at least 3 months with failure to achieve weight loss. Note: Not for patients with T1DM, T2DM.

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
	MC/DEL MC MC/DEL	VALPROIC ACID SOL VALTOCO ² ZONISAMIDE	MC MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC MC MC/DEL MC/DEL MC/DEL	8 8 8 8 8 8 8 8 8 8 9 9 9 9	VIGAFYDE VIMPAT ⁴ VIMPAT SOL ⁴ XCOPRI ZARONTIN SYRP ZARONTIN CAP ZARONTIN SOL ZONISADE ZTALMY KEPPRA XR NEURONTIN TEGRETOL-XR TB12		
ANTI-PARKINSON DRUGS							
PARKINSONS - ANTICHOLINERGICS	MC/DEL MC MC/DEL	BENZTROPINE MESYLATE TABS COGENTIN SOLN TRIHEXYPHENIDYL				Use PA Form# 20420	
PARKINSONS - ADENOSINE RECEPTOR ANTAGONIST			MC/DEL		NOURIANZ	Use PA Form# 20420	Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. DDI: Avoid use of Nourianz with strong CYP3A4 inducers (e.g. carbamazepine, rifampin, phenytoin, St. John's wort).
PARKINSONS - COMT INHIBITORS			MC/DEL MC		COMTAN TABS ONGENTYS	Use PA Form# 20420	Preferred drug must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
PARKINSONS - SELECTED DOPAMIN AGONISTS	MC/DEL MC/DEL	PRAMIPEXOLE ROPINIROLE NEUPRO PATCH	MC/DEL MC MC/DEL	5 8 8	MIRAPEX TABS ¹ REQUIP TABS MIRAPEX ER	Use PA Form# 20420 1. As of 12/08 users of Mirapex will be grandfathered if diagnosis is Parkinson's.	Preferred drug must be tried and failed in step-order due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
PARKINSONS- MAOIS			MC		XADAGO	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
PARKINSONS - DOPAMINERGICS/ CARBII/ LEVO	MC/DEL	AMANTADINE HCL CAPS	MC/DEL		APOKYN	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL	AMANTADINE HCL TABS	MC		AZILECT ²		
	MC/DEL	BROMOCRIPTINE MESYLATE TABS	MC/DEL		CARBIDOPA/LEVODOPA RAPDIS		
	MC/DEL	BROMOCRIPTINE MESYLATE CAPS	MC		CREXONT ⁴		
	MC/DEL	CARBIDOPA/LEVODOPA TABS ³	MC		ELDEPRYL CAPS		
	MC/DEL	CARBIDOPA/LEVODOPA ER	MC		GOCOVRI		
	MC/DEL	CARBIDOPA/LEVO/ENTACAPONE TAB	MC/DEL		INBRIJA		
	MC	LARODOPA TABS	MC		KYNMOBI		
	MC/DEL	SELEGILINE CAPS HCL	MC/DEL		ONAPGO		
			MC/DEL		OSMOLEX ER		
			MC/DEL		PARLODEL CAPS		
			MC/DEL		PARLODEL TABS		
			MC		RYTARY		
			MC		SINEMET TABS		
			MC		SINEMET TBCR		
			MC		VYALEV		
PARKINSONS - COMBO.			MC/DEL		STALEVO ¹	Use PA Form# 20420	
			MC		CARBIDOPA/LEVODOPA/ENTACA ¹	1. Clinical PA is required to establish diagnosis and medical necessity.	
MUSCLE RELAXANTS							
MUSCLE RELAXANTS	MC/DEL	BACLOFEN TABS	MC/DEL	7	ORPHENADRINE CITRATE	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC/DEL	CHLORZOXAZONE TABS	MC/DEL	8	CARISOPRODOL 350MG TABS		
	MC/DEL	CYCLOBENZAPRINE HCL 5mg & 10mg TABS	MC/DEL	8	AMRIX		
	MC	LIORESAL INTRATHECAL KIT	MC/DEL	8	DANTRIUM CAPS		
	MC/DEL	METHOCARBAMOL TABS	MC	8	FLEQSVUY		
	MC/DEL	TIZANIDINE HCL TABS	MC	8	LIORESAL TABS		
			MC	8	LORZONE		
			MC/DEL	8	LYVISPAN		
			MC	8	METAXALONE		
			MC	8	NORFLEX TBCR		
			MC	8	OZOBAX		
			MC	8	ROBAXIN-750 TABS		
			MC	8	VECUROMIUM INJ		
			MC/DEL	8	ZANAFLEX TABS		
			MC/DEL	9	CARISOPRODOL 250MG TABS		
			MC/DEL	9	CHLORZOXAZONE 250mg TABS		
			MC/DEL	9	SKELAXIN TAB		
			MC/DEL	9	SOMA TABS		
			MC	9	TANLOR		
MUSCLE RELAXANT - COMBO.			MC/DEL		CARISOPRODOL/ASPIRIN TABS	Use PA Form# 20420	Individual components are available with PA described in the section above. 1. frequent or persistent early refills of non-controlled drugs; 2. multiple instances of early refill overrides due to reports of misplacement stolen, dropped in toilet or sink, distant travel, etc.
			MC/DEL		CARISOPRODOL/ASPIRIN/CODE		
			MC		NORGESIC TABS		
			MC/DEL		ORPHENADRINE COMPOUND		
			MC/DEL		ORPHENADRINE/ASA/CAFF		
			MC		ORPHENGESIC		
PARATHYROID HORMONE							
PARATHYROID HORMONE			MC		NATPARA ¹	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
			MC		YORVIPATH ¹		

Category	Coverage Indicator	Preferred Drugs		Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
VITAMINS								
VITAMINS	MC	CYANOCOBALAMIN SOLN	MC			AQUASOLE SOLN	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval.
	MC	FERIVA CAP	MC			AQUAVIT-E SOLN	Please refer to OTC list for covered products.	
	MC	FERIVAF A CAP	MC			DHT SOLN		
	MC/DEL	FOLIC ACID TABS	MC			FUSION PLUS CAP		
	MC/DEL	MEPHYTON TABS	MC			HEMOCYTE PLU CAP		DDI: B-12 will now be non-preferred and require prior authorization if it is currently being used in combination with either prevacid, pantoprazole, prilosec, or any currently non preferred PPI.
	MC/DEL	NIACIN	MC			INTEGRA CAP		
	MC	NIACOR TABS	MC			INTEGRA F CAP	Click here for the OTC List	Please refer to OTC list for covered products.
	MC/DEL	NICOTINIC ACID SR CPCR	MC			INTEGRA PLUS CAP		Preferred products that used to require diag codes still require diag codes unless indicated otherwise.
	MC	PYRIDOXINE HCL TABS	MC			NASCOBAL GEL		
	MC	TANDEM CAP	MC			TANDEM PLUS CAP		
	MC/DEL	THIAMINE HCL SOLN						
	MC/DEL	VITAMIN B-1 TABS						
	MC/DEL	VITAMIN B-12						
	MC	VITAMIN B-6 TABS						
	MC/DEL	VITAMIN C						
	MC/DEL	VITAMIN E CAPS						
	MC/DEL	VITAMIN E/D-ALPHA CAPS						
	MC	VITAMIN K1 SOLN						
	MC	V-R VITAMIN E CAPS						
VITAMIN D's	MC/DEL	CALCITRIOL CAPS ¹	MC			CALCIJEX	Use PA Form# 20420	Preferred products require dialysis/renal failure diagnosis.
	MC/DEL	ROCALTROL	MC/DEL			DOXERCALCIF CAP		Rayaldee requires clinical PA to verify stage 3 or 4 CKD.
	MC/DEL	VITAMIN D2 ²	MC/DEL			DOXERCALCIF INJ		
	MC/DEL	VITAMIN D3 ²	MC/DEL			PARICALCITROL CAP		
	MC/DEL	VITAMIN DROPS	MC/DEL			PARICALCITROL INJ		
	MC	PARICALCITOL CAPS	MC/DEL			HECTOROL (ORAL)		
			MC			HECTOROL (PARENTERAL)		
			MC			RAYALDEE		
			MC			ZEMPLAR INJ		
			MC			ZEMPLAR CAPS		
ENZYMES								
POMPE DISEASE AGENTS			MC			NEXVIAZYME ¹	Use PA Form# 20420	All preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
			MC			LUMIZYME		Pombili and Opfolda are for the treatment of adult patients with late-onset Pompe disease (lysosomal acid alpha-glucosidase [GAA] deficiency) weighing ≥40kg and
			MC			OPFOLDA		who are not improving on their current enzyme replacement therapy (ERT).
			MC			POMBILITI		
MISC MULTI-VITAMINS								
VITAMINS - MISC.	MC	CENTRUM TABS	MC			ADEKS	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval.
	MC	CENTRUM JR/IRON CHEW	MC/DEL			ADVANCED NATALCARE TABS		
	MC	CENTRUM-LUTEIN TABS	MC			AQUADEKS		
	MC	CEROVITE ADVANCED FO TABS	MC			CENTRUM JR/EXTRA C CHEW		
	MC/DEL	CHEWABLE MULTIVIT/FL CHEW	MC			CENTRUM PERFORMANCE TABS		
	MC	COD LIVER OIL CAPS	MC			CENTRUM SILVER TABS		
	MC/DEL	COMPLETE NATAL DHA (ORAL)	MC			DALYVITE LIQD		
		COMBO PKG	MC			EMBREX 600 MISC		
	MC	COMPLETE SENIOR TABS	MC			FERRALET 90		
	MC	DAILY MULTI VIT/IRON	MC			IBERET		
	MC/DEL	DIALYVITE 1MG	MC			MATERNA TABS		
	MC/DEL	DIALYVITE 800MG	MC			MAXARON		
	MC/DEL	FULL SPECTRUM B	MC			MULTIRET FOLIC -500 TBCR		
	MC	M.V.I-12 INJ	MC/DEL			NATAFORT TABS		
	MC	MULTI-VIT/FLUORIDE	MC/DEL			NATALCARE CFE 60 TABS ¹		
	MC/DEL	NATALCARE RX TABS	MC/DEL			NATALCARE GLOSS TABS ¹		
	MC/DEL	NEPHRONEX	MC			NATALCARE PIC TABS ¹		
	MC/DEL	NIVA-PLUS (ORAL) TABLET	MC			NATALCARE PIC FORTE TABS ¹		
	MC/DEL	ONE DAILY TABS	MC/DEL			NATALCARE PLUS TABS ¹		
	MC/DEL	ONE-DAILY MULTIVITAMINS	MC			NATALCARE THREE TABS ¹		
	MC/DEL	ONE-TABLET-DAILY	MC/DEL			NATACHEW CHEW		

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required			Criteria
	MC/DEL	POLY-VIT/IRON/FLUORID SOLN	MC		NATALFIRST TABS			
MINERALS								
	MC/DEL	POLY-VITAMIN/FLUORIDE SOLN	MC		NATATAB RX TABS			
	MC/DEL	POLY-VITAMINS/IRON SOLN	MC/DEL		NEPHPLEX RX TABS			
	MC	PRENATA (ORAL) TAB CHEW	MC/DEL		NEPHROCAPS CAPS			
	MC/DEL	PRENATAL TABS ¹	MC/DEL		NEPHRO-VITE TABS			
	MC/DEL	PRENATAL FORMULA 3 TABS ¹	MC		NESTABS RX TABS			
	MC/DEL	PRENATAL PLUS TABS ¹	MC/DEL		NIFEREX			
	MC/DEL	PRENATAL PLUS NF TABS ¹	MC/DEL		OCUVITE TABS			
	MC	PRENATAL PLUS/27MG IRON ¹	MC		POLY-VI-FLOR SOLN			
	MC	PRENATAL PLUS/IRON TABS ¹	MC		POLY-VI-SOL SOLN			
	MC	PRENATAL VITAMIN PLUS LOW IRON (ORAL) TABLET	MC		POLY-VI-SOL/IRON SOLN			
	MC/DEL	PRENATAL RX/BETA-CAROTENE ¹	MC		POLY-VITAMIN DROPS SOLN			
	MC/DEL	PREPLUS (ORAL) TABLET	MC		PRECARE			
	MC/DEL	RENAL CAPS	MC		PREFERA OB			
	MC/DEL	RENAPHRO CAPS	MC		PREMESIS RX TABS			
	MC	STRESS TAB NF TABS	MC		PRENATABS CBF TABS ¹			
	MC	THERAPEUTIC-M TABS	MC		PRENATAL CARE TABS ¹			
	MC	THERAVITE LIQD	MC/DEL		PRENATAL MR 90 TBCR ¹			
	MC/DEL	TRINATAL RX 1 (ORAL) TABLET	MC		PRENATAL MTR/SELENIUM TABS ¹			
	MC/DEL	TRIVEEN-DUO DHA (ORAL) COMBO. PKG	MC		PRENATAL OPTIMA ADVANCE TABS ¹			
	MC/DEL	TRI-VITAMIN/FLUORIDE SOLN	MC/DEL		PRENATAL PC 40 TABS ¹			
	MC	VITA CON FORTE CAPS	MC		PRENATAL RX TABS ¹			
	MC	VITAPLEX PLUS TABS	MC		PRENATE ¹			
			MC		PRENATE ELITE ¹			
			MC		PRIMACARE MISC			
			MC		PROTEGRA CAPS			
			MC		STUARTNATAL PLUS 3 TABS ¹			
			MC		TRI-VI-SOL SOLN			
			MC		TRI-VI-SOL/IRON SOLN			
			MC/DEL		ULTRA NATALCARE TABS			
			MC		ULTRA-NATAL TABS ¹			
			MC		VICON FORTE CAPS			
			MC		VINATAL FORTE TABS ¹			
			MC		VINATE ¹			
			MC/DEL		VINATE ADVANCED TABS ¹			
MISCELLANEOUS MINERALS								
	MC	CALCARB	MC		ANEMAGEN	Use PA Form# 20420		
	MC	CALCI-MIX CAPSULE CAPS	MC		CALCET TABS	Please refer to OTC list.		
	MC	CALCIQUID SYRP	MC/DEL		CALCIUM 600-D TABS			
	MC	CALCITRATE/VITAMIN D TABS	MC		CALCIUM/VITAMIN D TABS			
	MC/DEL	CALCIUM	MC		CALTRATE 600 PLUS/VIT D TABS			
	MC/DEL	CALCIUM CARBONATE	MC		CALTRATE PLUS TABS			
	MC/DEL	CALCIUM CITRATE TABS	MC		CHROMAGEN			
	MC/DEL	CALCIUM GLUCONATE TABS	MC		CITRACAL PLUS TABS			
	MC/DEL	CALCIUM LACTATE TABS	MC		CONTRIN CAPS			
	MC	CALCIUM/MAGNESIUM TABS	MC		FEOGEN FORTE CAPS			
	MC/DEL	CALCIUM/VITAMIN D TABS	MC		FEROCON CAPS			
	MC	CALTRATE 600 TABS	MC/DEL		FERREX 150 CAPS			
	MC/DEL	CHEWABLE CALCIUM CHEW	MC		FERRO-SEQUELS TBCR			
	MC	CITRACAL TABS	MC		FE-TINIC CAPS			
	MC	CITRACAL + D TABS	MC		FE-TINIC 150 FORTE CAPS			
	MC	CITRUS CALCIUM TABS	MC/DEL		FLUOR-A-DAY SOLN			
	MC	CITRUS CALCIUM 1500 + D TABS	MC		HEMOCYTE TABS			
	MC	EFFERVESCENT POTASSIUM TBDF	MC/DEL		K-DUR TBCR			
	MC/DEL	FEOSTAT CHEW	MC		KLOR-CON PACK			
	MC	FERATAB TABS	MC		K-LYTE			
	MC/DEL	FER-GEN-SOL SOLN	MC/DEL		K-PHOS TABS NEUTRAL			
	MC	FER-IRON SOLN	MC		K-TABS TBCR			
	MC	FERRONATE TABS	MC		K-VESCENT PACK			

Category	Coverage Indicator	Preferred Drugs		Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
MISC. ELECTROLYTES/NUTRITIONALS								
ELECTROLYTES/ NUTRITIONALS	MC	INTRALIPID EMUL ¹	MC		BOOST ¹	Use PA Form# 20420 & SGA Form	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Certain drugs require specific diagnoses for approval.	
	MC	P.T.E. -5 SOLN ¹	MC		CASEC POWD ¹		Medical foods are not to be authorized solely for the purpose of enhancing nutrient intake or managing body weight if the participant is able to eat conventional foods adequately. Medical foods may be approved if the member has a medical condition which precludes or restricts the use of conventional foods and necessitates the use of a formula. Concurrent Stimulant therapy is not an acceptable medical reason/condition for use of medical foods for enhancing nutrient intake or managing body weight.	
	MC	SEA-OMEGA CAPS ¹	MC		CHOICE DM LIQD ¹			
	MC		MC		DELIVER 2.0 LIQD ¹			
	MC		MC		DOJOLVI			
	MC		MC		ENFAMIL ¹			
	MC		MC		ENSURE ¹			
	MC		MC		GLUCERNA ¹			
	MC		MC		ISOCAL LIQD ¹			
	MC		MC		KINDERCAL TF LIQD ¹			
	MC		MC		KINDERCAL TF/FIBER LIQD ¹			
	MC		MC		L-CARNITINE CAPS ¹			
	MC		MC		LIPISORB LIQD ¹			
	MC		MC		LOVAZA ^{1,2}			
	MC		MC		MODULEN IBD POWD ¹			
	MC		MC		NUTRAMIGEN POWD ¹			
	MC		MC		NUTREN ¹			
	MC		MC		NUTRITIONAL SUPPLEMENT LIQD ¹			
	MC		MC		NUTRIVENT 1.5 LIQD ¹			
	MC		MC		PEPTAMEN ¹			
	MC		MC		PHENYLADE ¹			
	MC		MC		PHENYL-FREE ¹			
	MC		MC		PKU 3 POWD ¹			
	MC		MC		PREGESTIMIL POWD ¹			
	MC		MC		PROBALANCE LIQD ¹			
	MC		MC		PROSOBEE ¹			
	MC		MC		SCANDISHAKE PACK ¹			
	MC		MC		VASCEPA			
ERYTHROPOEITINS	MC	EPOGEN SOLN	MC	8	ARANESP SOLN ¹	Use PA Form# 10520	Non-Preferred drugs must be tried and failed in step-order, due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Please see the EPO PA form for other approval and renewal criteria.	
	MC	MIRCERA SYRINGE	MC	8	PROCRIT SOLN ¹			
	MC	RETACRIT						
GRANULOCYTE CSF								
GRANULOCYTE CSF	MC	FULPHILA	MC	8	FYLNETRA	Use PA Form# 20520	See approval criteria detailed on Granulocyte Colony Stimulating Factor PA form.	
	MC	NEUPOGEN SYRINGE	MC	8	GRANIX SYRINGE			
	MC	NEUPOGEN VIAL	MC	8	GRANIX VIAL			
	MC/DEL	NYVEPRIA SYRINGE	MC	8	LEUKINE			
			MC/DEL	8	NIVESTYM			
			MC	8	ROLVEDON			
			MC	8	RYZNEUTA			
			MC	8	STIMUFEND			
			MC/DEL	8	ZARXIO			
			MC/DEL	8	ZIEXTENZO			
			MC	9	NEULASTA ¹			
GAUCHER DISEASE								
GAUCHER DISEASE			MC		CERDELGA ¹	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Exceeding days supply limits for LMWH class requires PA.	
			MC		YARGESA ¹			
							Yargesa: As monotherapy for the treatment of adult patients with mild to moderate type 1 Gaucher disease for whom enzyme replacement therapy is not a therapeutic option (e.g., due to allergy, hypersensitivity, or poor venous access).	
NIEMANN-PICK DISEASE AGENTS								
NIEMANN-PICK DISEASE AGENTS			MC		AQNEURSA ¹	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.	
			MC		MIPLYFFA ¹			

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
ANTICOAGULANTS / PLATELET AGENTS							
ANTICOAGULANTS	MC COUMADIN TABS MC/DEL ENOXAPARIN ¹ MC ELIQUIS MC ELIQUIS STARTER PACK MC HEPARIN SODIUM/NACL 0.9% SOLN MC HEP-LOCK SOLN MC INNOHEP MC HEPARIN LOCK SOLN MC/DEL HEPARIN LOCK FLUSH SOLN MC/DEL HEPARIN SODIUM SOLN MC/DEL HEPARIN SODIUM LOCK FLUSH SOLN MC/DEL PRADAXA MC/DEL JANTOVEN MC/DEL WARFARIN SODIUM TABS MC/DEL XARELTO MC/DEL XARELTO STARTER PACK	MC COUMADIN TABS MC/DEL ENOXAPARIN ¹ MC ELIQUIS MC ELIQUIS STARTER PACK MC HEPARIN SODIUM/NACL 0.9% SOLN MC HEP-LOCK SOLN MC INNOHEP MC HEPARIN LOCK SOLN MC/DEL HEPARIN LOCK FLUSH SOLN MC/DEL HEPARIN SODIUM SOLN MC/DEL HEPARIN SODIUM LOCK FLUSH SOLN MC/DEL PRADAXA MC/DEL JANTOVEN MC/DEL WARFARIN SODIUM TABS MC/DEL XARELTO MC/DEL XARELTO STARTER PACK	MC COUMADIN TABS MC/DEL ENOXAPARIN ¹ MC ELIQUIS MC ELIQUIS STARTER PACK MC HEPARIN SODIUM/NACL 0.9% SOLN MC HEP-LOCK SOLN MC INNOHEP MC HEPARIN LOCK SOLN MC/DEL HEPARIN LOCK FLUSH SOLN MC/DEL HEPARIN SODIUM SOLN MC/DEL HEPARIN SODIUM LOCK FLUSH SOLN MC/DEL PRADAXA MC/DEL JANTOVEN MC/DEL WARFARIN SODIUM TABS MC/DEL XARELTO MC/DEL XARELTO STARTER PACK		ARIIXTRA SOLN FONDAPARINUX FRAGMIN INJ FRAGMIN VIAL LOVENOX SOLN LOVENOX 300² LOVENOX SUBQ SYRINGE PRADAXA ORAL PELLETS⁴ IPRIVASK SAVAYSAS³	Use PA form# 20420 1. Enoxaparin therapy durations greater than 7 days every 30 days require PA. 2. Use other strengths available to obtain desired dose. 3. Diagnosis required. 4. For the treatment of patients aged 3 months to less than 12 years of age.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Exceeding days supply limits for LMWH class requires PA. DDI: Warfarin will require prior authorization if being used in combination with fluconazole, miconazole, or voriconazole. DDI: Warfarin will require prior authorization if being used in conjunction with gemfibrozil or fenofibrate. DDI: Rifampin will require prior authorization if being used in combination with savaysa.
ANTIHEMOPHILIC AGENTS	MC ALPHANATE MC ALPHANINE SD MC/DEL ALPROLIX VIAL MC/DEL BEBULIN VIAL MC/DEL BENEFIX SOLR MC/DEL HELIXATE FS KIT MC HEMOFIL - M MC HUMATE-P SOLR MC/DEL IXINITY VIAL MC/DEL JIVI ³ MC KOATE-DVI MC KONYNE - 80 MC/DEL KOVALTRY MC/DEL REBINYN MC MONARC - M MC MONOCLOATE - P MC MONONINE MC/DEL NOVOEIGHT MC NOVOSEVEN SOLR MC NUWIQ MC/DEL PROFILNINE MC RECOMBINATE SOLR MC REFACTO MC/DEL RIXUBIS VIAL MC WILATE INJ MC/DEL XYNTHA	MC/DEL ALPHANATE MC ALPHANINE SD MC ALPROLIX VIAL MC/DEL BEBULIN VIAL MC/DEL BENEFIX SOLR MC/DEL HELIXATE FS KIT MC/DEL HEMOFIL - M MC/DEL HUMATE-P SOLR MC/DEL IXINITY VIAL MC/DEL JIVI ³ MC KOATE-DVI MC KONYNE - 80 MC/DEL KOVALTRY MC/DEL REBINYN MC MONARC - M MC MONOCLOATE - P MC MONONINE MC/DEL NOVOEIGHT MC NOVOSEVEN SOLR MC NUWIQ MC/DEL PROFILNINE MC RECOMBINATE SOLR MC REFACTO MC/DEL RIXUBIS VIAL MC WILATE INJ MC/DEL XYNTHA	MC/DEL ALPHANATE MC ALPHANINE SD MC ALPROLIX VIAL MC/DEL BEBULIN VIAL MC/DEL BENEFIX SOLR MC/DEL HELIXATE FS KIT MC/DEL HEMOFIL - M MC/DEL HUMATE-P SOLR MC/DEL IXINITY VIAL MC/DEL JIVI ³ MC KOATE-DVI MC KONYNE - 80 MC/DEL KOVALTRY MC/DEL REBINYN MC MONARC - M MC MONOCLOATE - P MC MONONINE MC/DEL NOVOEIGHT MC NOVOSEVEN SOLR MC NUWIQ MC/DEL PROFILNINE MC RECOMBINATE SOLR MC REFACTO MC/DEL RIXUBIS VIAL MC WILATE INJ MC/DEL XYNTHA	ADYNOVATE VIAL ADVATE^{1,2,5} ALTUVIPIO⁴ AFSTYLA BEQVEZ ESPEROCT ELOCTATE HEMGENIX IDELVION KOGENATE FS⁵ RECOMBINATE VIAL⁵ ROCTAVIAN⁴ SEVENFACT	Use PA Form# 20420 1. Only if other products unavailable. 2. Advate may be available with PA in cases of large volume dosing in patients with poor venous access. 3. Not indicated for use in children <12 years of age due to greater risk for hypersensitivity reactions and is not indicated for use in previously untreated patients. 4. Clinical PA required for appropriate diagnosis. 5. Established users will be grandfathered.	Non-preferred will only be approved if other preferred products are unavailable. Beqvez - FDA Approved Indication: An adeno-associated virus vector-based gene therapy indicated for the treatment of adults with moderate to severe hemophilia B (congenital factor IX deficiency) who: <ul style="list-style-type: none"> • Currently use factor IX prophylaxis therapy, or • Have current or historical life-threatening hemorrhage, or • Have repeated, serious spontaneous bleeding episodes, and, • Do not have neutralizing antibodies to adeno-associated virus serotype Rh74var (AAVRh74var) capsid as detected by an FDA-approve test. Hemgenix is an adeno-associated viral vector-based gene therapy for IV infusion after dilution. For treatment of adults with Hemophilia B (congenital Factor IX deficiency) who: Currently use Factor IX prophylaxis therapy, or have current or historical life-threatening hemorrhage, or have repeated, serious spontaneous bleeding episodes.	
NON-FACTOR REPLACEMENT THERAPY	MC HEMLIBRA	MC/DEL HEMLIBRA	MC/DEL HEMLIBRA		ALHEMO HYMPAVZI QFITLIA QFITLIA PEN	Use PA Form# 20420	Subsequent changes made to Antihemophilic Agents: Factor Therapy to move Hemlibra to Non-Factor Therapy
PLATELET AGGREGATION INHIBITORS	MC/DEL ASPIRIN MC ASPIRIN-DIPYRIDAMOLE ER CPMP 12HR MC/DEL BRILINTA 90mg MC/DEL DIPYRIDAMOLE TABS MC/DEL CLOPIDOGREL 75MG MC/DEL PRASUGREL HCL TAB	MC/DEL ASPIRIN-DIPYRIDAMOLE ER CPMP 12HR MC/DEL BRILINTA 90mg MC/DEL DIPYRIDAMOLE TABS MC/DEL CLOPIDOGREL 75MG MC/DEL PRASUGREL HCL TAB	7 8 8 8 8 8	TICLOPIDINE HCL TABS BRILINTA 60mg DURLAZA EFFIENT PERSANTINE TABS PLAVIX TABS ZONTIVITY	Use PA Form# 20715 for Plavix, Effient & Brilinta Use PA form# 20420 for other requests 1. Dosing limits apply, see Dose Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.	
							A special PA may be obtained at the pharmacy for members scheduled for "stent" placement or have had placement if in the last 12months. Please indicate on prescription date of stent placement. Brilinta - Concomitant use with strong CYP3A4 inhibitors should be avoided (including ketoconazole, itraconazole, atazanavir, and telithromycin). Doses of simvastatin and lovastatin >40mg should be avoided. DDI: exists for using maintenance ASA dose >100mg, as it reduces the effectiveness of Brilinta. DDI: Plavix will require prior authorization if being used in combination with omeprazole, esomeprazole, cimetidine, fluconazole, ketoconazole, intelence, fluoxetine, ticlopidine, and fluvoxamine.

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
PLATELET AGGR. INHIBITORS / COMBO'S - MISC.	MC/DEL MC/DEL	CILOSTAZOL PENTOXIFYLLINE ER TBCR	MC/DEL MC/DEL MC/DEL MC MC		AGRYLIN CAPS ANAGRELIDE CAPS PLETAL TABS TRENTAL TBCR YOSPRALA	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
HEMATOLOGICALS							
MONOCLONAL ANTIBODY			MC MC MC/DEL MC MC MC/DEL MC MC MC/DEL MC MC		BKEMV EMPAVELI ENSPRYNG EPYSQLI FABHALTA GAMIFANT PIASKY SOLIRIS ULTOMIRIS UPLIZNA VOYDEYA	Use PA Form# 20420	A diagnosis of Paroxysmal nocturnal hemoglobinuria (PNH) using the HAM test or flow cytometry is required. In addition, the patient must show evidence of having received a meningitis vaccine at least 2 weeks prior to the start of therapy. Gamifant is recommended for the treatment of adult and pediatric (newborn and older) patients with primary hemophagocytic lymphohistiocytosis (HLH) with refractory, recurrent, or progressive disease or intolerance with conventional HLH therapy. Fabhalta and Ultomiris are recommended for the treatment of adults with paroxysmal nocturnal hemoglobinuria (PNH). Bkemv and Epysqli have updated criteria for a diagnosis of generalized myasthenia gravis (gMG): must have confirmation that patients are anti-acetylcholine receptor (AChR) antibody positive.
IMMUNE GLOBULIN	MC MC/DEL MC MC/DEL MC/DEL MC/DEL MC	BIVIGAM ¹ CUTAQUIG ¹ GAMMAGARD S-D ¹ HIZENTRA ¹ PANZYGA ¹ PRIVIGEN ¹	MC MC MC/DEL MC MC/DEL MC MC/DEL		ALYGLO ASCENIV ² CUVITRU GAMMAPLEX INJ HYQVIA OCTAGAM INJ ¹ XEMBIFY	Use PA Form# 20420 1. Clinical PA required. 2. For the treatment of patients between 12 to 17 years of age.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Alyglo is indicated for treatment of primary humoral immunodeficiency in adults ages 17 or older. Cutaquig is indicated as replacement therapy for primary humoral immunodeficiency (PI) in adults. Xembify is indicated for treatment of primary humoral immunodeficiency (PI) in patients 2 years of age and older. Asceniv indicated for the treatment of primary humoral immunodeficiency (PI) in adults and adolescents (12 to 17 years of age). PI includes but is not limited to the humoral immune defect in congenital agammaglobulinemia, common variable immunodeficiency (CVID), X-linked agammaglobulinemia, Wiskott-Aldrich syndrome, and severe combined immunodeficiencies (SCID).
HEREDITARY ANGIOEDEMA	PROPHYLAXIS		PROPHYLAXIS		Use PA Form# 20420		Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Haegarda is indicated for routine prophylaxis to prevent Hereditary Angioedema (HAE) attacks in adolescent and adult patients
	MC MC MC MC/DEL	CINRYZE ¹ HAEGARDA ¹ ORLADEYO ^{1,2} TAKHYRO ¹	MC MC	8 8	ANDEMBRY DAWNZERA ²	1. Clinical PA is required to establish diagnosis and medical necessity. 2. For the treatment of patients \geq 12 years of age	
HEMATOLOGICAL AGENTS-THROMBOPOIETIN RECEPTOR AGONISTS	TREATMENT		TREATMENT		Use PA Form# 20420		Doptelet and Mulpelta : For the treatment of thrombocytopenia in adults with chronic liver disease who are scheduled to undergo a procedure.
	MC/DEL MC MC/DEL	BERINERT KIT ¹ FIRAZYR ¹ RUCONEST VIAL ¹	MC/DEL MC	8 8	KALBITOR VIAL EKTERLY ²	1. Clinical PA is required to establish diagnosis and medical necessity. 2. For the treatment of patients \geq 12 years of age	
HEMATOLOGICAL AGENTS-IgAN	MC MC	PROMACTA ¹ NPLATE ¹	MC MC/DEL MC/DEL		ALVAIZ DOPTELET MULPLETA	Use PA Form# 20420 1. Clinical PA required. Must see prior trial with insufficient response to corticosteroids and immunoglobulins.	Doptelet and Mulpelta : For the treatment of thrombocytopenia in adults with chronic liver disease who are scheduled to undergo a procedure. All preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs (in step-order) will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists PA required to confirm FDA-approved indication. Vanrafia is for adults with biopsy proven primary IgAN AND eGFR \geq 30 cc/min/1.73m ² AND urine protein \geq 1 g/day AND on stable dose of maximally tolerated renin-angiotensin system inhibitor.
ANEMIA- BETA THALASSEMIA			MC MC		REBLOZYL ZYNTEGLO	Use PA Form# 20420	Reblozyl is indicated for three (3) treatments of anemia in adults: 1. in adult patients with beta thalassemia who require regular red blood cell (RBC) transfusions; 2. without previous erythropoiesis stimulating agent use (ESA-naïve) in adult patients with very low- to intermediate-risk myelodysplastic syndromes (MDS) who may require regular RBC transfusions; and 3. failing an ESA and requiring 2 or more RBC units over 8 weeks in adult patients with very low- to intermediate-risk MDS with ring sideroblasts (MDS-RS) or with myelodysplastic/myeloproliferative neoplasm with ring sideroblasts and thrombocytosis (MDS/MPN-RS-T). It is not indicated for use as a substitute for RBC transfusions in patients who require immediate correction of anemia. Zynteglo is indicated for the treatment of adult and pediatric patients with β -thalassemia who require regular red blood cell (RBC) transfusions.
HEMATOLOGIC DISORDER TREATMENT AGENTS			MC/DEL MC MC		CABLIVI WAYRILZ ¹ TAVALISSE	Use PA Form# 20420 1. Clinical PA is required to establish diagnosis and medical necessity.	Tavalisse is recommended for patients at risk of bleeding when one line of therapy (steroids, IVIG, splenectomy) has failed. Cablivi is recommended for the treatment of adult patients with acquired thrombotic thrombocytopenic purpura (aTTP), in combination with plasma exchange and immunosuppressive therapy. Wayrilz : Baseline platelet count is less than 30,000/mcL and prescribed in consultation or by a hematologist/oncologist.
COMPLEMENT RECEPTOR ANTAGONIST			MC		TAVNEOS	Use PA Form# 20420	
WHIM SYNDROME AGENTS			MC		XOLREMDI	Use PA Form# 20420	Xolremdi : In patients 12 years of age and older with WHIM syndrome (warts, hypogammaglobulinemia, infections, and myelokathexis) to increase the number of circulating mature neutrophils and lymphocytes.

Category	Coverage Indicator	Preferred Drugs		Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
Dermatological								
ISOTRETINION, ACNE		AMNESTEEM ¹ CLARAVIS ¹ MYORISAN ¹ ZENATANE ¹			MC	ABSORICA ABSORICA LD	Use PA Form# 20420 1. Users 24 or under, PA will not be required.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - ACNE PREPARATIONS		ERYDERM SOLN ERYTHROMYCIN GEL ERYTHROMYCIN SOLN EVOCLIN ISOTRETINOIN METRONIDAZOLE GEL ² METRONIDAZOLE LOTN ² TRETINOIN .025%, .01% GEL ¹ TRETINOIN CREA ^{1,2}			MC/DEL	AKLIEF ⁶ ALTINAC CREA AMZEEQ ⁶ AVITA CREA BENZAC BENZACLIN GEL ³ BENZAGEL-10 GEL BENZEFOAM BREVOXYL CLEOCIN-T ² CLINAC BPO GEL CLINDETS SWAB DESQUAM-E GEL DESQUAM-X DIFFERIN 0.3% GEL DIFFERIN EMGEL GEL EPIDUO EPSOLAY ERYCETTE PADS FINEVIN CREA METROCREAM CREA ² METROGEL GEL ² METROLOTION LOTN ² NEOBENZ MICRO PLIXDA RHOFADE SODIUM SULFACET/SULF LOTN SOOLANTRA ⁴ TRIAZ TWYNEO VELTIN WINLEVI ⁵ ZENCIA WASH ZETACET ZILXI	Use PA Form# 10220 for Brand Name requests Use PA Form# 20420 for all other requests 1. Users 24 or under, PA will not be required. 2. Dosing limits allow one package per month. Refer to Dose Consolidation List. 3. Only available if component ingredients are unavailable. 4. Dosing limits apply, see Dosing Consolidation List. 5. Not approved for use in children <12 years of age. 6. For the treatment of patients ≥ 9 years of age.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL- ATOPIC DERMATITIS		PIMECROLIMUS PROTOPIC OINT TACROLIMUS OINT ADBRV ^{2,4} EBGLYSS ^{2,3,4} DUPIXENT ^{1,2,4} EUCRISA ^{2,4} OPZELURA ^{2,3,4} RINVOQ ⁵			MC	ANZUPGO	Use PA Form# 20420 1. Avoid live vaccines if treated with Dupixent. 2. Clinical PA required. 3. For the treatment of patients ≥ 12 years of age. 4. Preferred only after a trial and failure of TCI. 5. Clinical PA is required to establish diagnosis and medical necessity.	Preferred drugs also indicated for this condition, including topical steroids, cyclosporin AND calcineurin inhibitors must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Note: If unable to use TCIs then a trial of Eucrisa could be recommended before Dupixent. ANZUPGO: use of Anzupgo in combination with other JAK inhibitors or potent immunosuppressants is not recommended.

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
TOPICAL - CORTICOSTEROIDS		LOW POTENCY			LOW POTENCY	Use PA Form# 20420	At least 1 drug from each potency of preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
	MC	DERMA-SMOOTH-E FS BODY	MC/DEL		ACLOVATE		
	MC/DEL	HYDROCORTISONE CREA	MC		DESONATE GEL		
	MC	HYDROCORTISONE LOTN	MC/DEL		FLUOCINOLONE ACETONIDE		
	MC	HYDROCORTISONE LOTN	MC/DEL		FLUOCINOLONE		
	MC	TEXACORT SOLN	MC		HALOG		
			MC		HYDROCORTISONE POWD		
			MC		LIDA MANTLE HC CREA		
			MC		PROCTOCORT CREA		
			MC/DEL		VERDESO		
		MEDIUM POTENCY			MEDIUM POTENCY		
	MC/DEL	DESOXIMETASONE 0.05% CREA/GEL	MC/DEL		BESER LOTION ³		
	MC	FLUTICASONE PROPIONATE CREA/OINT	MC		CLODERM CREA		
	MC	HYDROCORTISONE BUTYRATE	MC/DEL		CORDRAN		
	MC	HYDROCORTISONE OINT	MC/DEL		CUTIVATE CREA / OINT		
	MC	HYDROCORTISONE VALERATE	MC/DEL		CUTIVATE LOTN		
	MC	MOMETASONE FUROATE OINT	MC/DEL		DERMATOP		
	MC	TRIAMCINOLONE ACETONIDE .025-.1%	MC		ELOCON OINT		
			MC/DEL		KENALOG AERS		
			MC/DEL		LOCOID		
			MC		LUXIQ FOAM		
			MC/DEL		PANDEL CREA		
			MC		TOPICORT		
			MC		TOPICORT LP CREA		
			MC/DEL		TOVET FOAM ³		
			MC		WESTCORT		
		HIGH POTENCY			HIGH POTENCY		
	MC/DEL	DESONIDE ¹	MC		AMCINONIDE CREA		
	MC	TRIAMCINOLONE ACETONIDE .5%	MC		BETAMETHASONE DIPROPIONATE		
			MC/DEL		DESOXIMETASONE 0.25% CREA/OINT		
		VERY HIGH POTENCY			VERY HIGH POTENCY		
	MC/DEL	AUGMENTED BETA DIP	MC/DEL		CLOBETASOL PROPIONATE LOTN		
	MC/DEL	BETAMETHASONE VALERATE	MC/DEL		CLOBETASOL PROPIONATE SHAMPOO 0.05%		
	MC	DIFLORASONE DIACETATE	MC/DEL		CORMAX		
			MC/DEL		DIPROLENE		
			MC/DEL		IMPEKLO ⁴		
			MC/DEL		LEXETTE		
			MC/DEL		OLUX FOAM		
			MC/DEL		PSORCON		
			MC/DEL		PSORCON E		
			MC		SERNIVO SPRAY ²		
			MC/DEL		TEMOVATE		
			MC		ULTRAVATE		
		MISCELLANEOUS					
	MC	PROCTO-KIT CREA 1%					
TOPICAL - STEROID LOCAL ANESTHETICS			MC		EPIFOAM FOAM	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - STEROID COMBINATIONS	MC	DERMA-SMOOTH-E FS SCALP	MC		CARMOL-HC CREA	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.

CATEGORY	Coverage Indicator	PREFERRED DRUGS	Coverage Indicator	Step Order	NON-PREFERRED DRUGS PA Required		Criteria
TOPICAL - EMOLLIENTS	MC/DEL MC MC	AMMONIUM LACTATE CREA ¹ AMMONIUM LACTATE LOTN 12% ¹ VITAMIN A & D MEDICATED OINT	MC MC MC MC MC		LAC-HYDRIN CREA ¹ LAC-HYDRIN LOTN 12% MEDERMA GEL MIMYX RENOVA CREA	Use PA Form# 20420 1. Dosing limits still apply, see Dose Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - ENZYMES / KERATOLYTICS / UREA			MC MC MC		CARMOL 40 CREA SALEX CREA SALEX LOTN	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Ziox, Panafil and Papain products have been removed from the PDL due to FDA safety concerns regarding drugs containing Papain.
TOPICAL - GENITAL WARTS	MC/DEL	IMIQUIMOD 5% ²	MC/DEL MC/DEL MC/DEL MC MC	5 8 8 8 8	PODOFILOX SOLN CONDYLOX ¹ ALDARA ¹ PICATO VEREGEN ¹	Use PA Form# 20420 1. Non-preferred products must be used in specified order. 2. Dosing limits still apply, see Dose Consolidation List.	
TOPICAL - LOCAL ANESTHETICS	MC MC/DEL MC/DEL MC/DEL MC MC/DEL MC/DEL MC/DEL MC/DEL MC/DEL	AF CAPSICUM OLEORESIN CREA CAPSAICIN CREA CAPSAICIN PATCH DIBUCAIN OINT ELA-MAX ¹ LIDOCAINE/PRilocaine CREA ¹ LIDOCAINE CREAM LIDOCAINE GEL LIDOCAINE PTCH 5%	MC/DEL MC/DEL MC MC MC MC/DEL		EMLA PADS EMLA CREA LIDA MANTLE CREA PONTOCAINE SOLN SYNERA ZOSTRIX ZTLIDO ²	Use PA Form# 20420 1. Lidocaine/Prilocaine cream and Ela-Max products require PA for users over 18 years of age. 2. Dosing limits still apply, see Dose Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - DEPIGMENTING AGENTS			MC MC MC MC/DEL MC/DEL MC MC MC	8 8 8 8 8 8 8 9	ALUSTRA CREA EPIQUIN MICRO GLYQUIN CREA HYDROQUINONE CREA HYDROQUINONE/SUNSCREENS SOLAQUIN FORTE CREA TRI-LUMA CREA ELDOQUIN	Use PA Form# 20420	As per Medicaid Policy, cosmetic drugs are not covered. Non-cosmetic clinical applications will be considered by prior authorization on a case by case basis.
TOPICAL - SCABICIDES AND PEDICULICIDES	MC/DEL MC MC/DEL MC/DEL MC	ACTICIN CREA LICE KILLING SHAM LICE TREATMENT CREME RINS LIQD PERMETHRIN LOTN NATROBA ¹	MC MC MC/DEL MC MC MC/DEL		ELIMITE CREA EURAX LINDANE MALATHION OVIDE LOTN SPINOSAD SUSP	Use PA Form# 20420 1. Dosing limits apply, see Dosage Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
TOPICAL - WOUND / DECUBITUS CARE			MC MC MC		FILSUEZ REGRANEX GEL VYJUVEK	Use PA Form# 20420	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. Regranex will be approved for diabetic patients in good control (hgbA1c <8), who are not smoking, with a stage III or IV WOCN AND NPUAP lower extremity diabetic ulcer and with an adequate blood supply (TcP 02 >30, ABI>0.7 or ASP> 70), and where the underlying cause has been corrected. The wound must be free of infection and have been previously treated with preferred standard therapies for at least 2 months. Maximum approval for 20 weeks. Accuzyme and Ethezyme products have been removed from the PDL due to FDA concerns regarding drugs containing papain. Filsuvez: The patient has a diagnosis of dystrophic or junctional epidermolysis bullosa. The patient is at least 6 months old and does not have current evidence or history of squamous cell carcinoma or active infection in the area requiring filsuvez application. The patient has used standard wound care treatments, including silicone or foam dressings without wound resolution Vyjuvek: For the treatment of wounds in patients 6 months of age and older with dystrophic epidermolysis bullosa (DEB) with mutation(s) in the collagen type VII alpha 1 chain (COL7A1) gene.
TOPICAL - ASTRINGENTS / PROTECTANTS			MC MC MC		MOISTURIN DRY SKIN CREA PROSHIELD PLUS SKIN PROTE CREA SURGILUBE GEL	Use PA Form# 20420 1. Dosing limits apply, see Dosage Consolidation List.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists.
HYPERHIDROSIS THERAPY - AXILLARY	MC	XERAC AC SOLN	MC	8	SOFDRA ^{1,2}	1. Clinical PA is required to establish diagnosis and medical necessity. 2. For adults and pediatric patients 9 years of age and older.	Preferred drugs must be tried and failed due to lack of efficacy or intolerable side effects before non-preferred drugs will be approved, unless an acceptable clinical exception is offered on the Prior Authorization form, such as the presence of a condition that prevents usage of the preferred drug or a significant potential drug interaction between another drug and the preferred drug(s) exists. SOFDRA: prescribed by a dermatologist.

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required		Criteria
ANTINEOPLASTIC AGENTS- LHRH ANALOGS	MC/DEL MC/DEL MC/DEL MC/DEL	LUPRON DEPOTSYRINGEKIT ¹ LUPRON DEPOT- PED KIT ¹ (1-month) LUPRON DEPOT-PED SYRINGEKIT (3-month) TRIPTODUR VIAL	MC/DEL MC/DEL MC/DEL MC		FIRMAGON ² SUPPRELIN LA (IMPLANT) KIT TRELSTAR VANTAS ²	Use PA Form# 20420 1. Dosing limits apply, please refer to Dosage Consolidation List. 2. PA required to confirm FDA approved indication.	
ANTINEOPLASTIC AGENTS - TYROSINE KINASE INHIBITORS			MC MC/DEL MC		SPRYCEL ¹ TYKERB ² GLEEVEC ¹	Use PA Form# 20420 1. Verification of diagnosis is required. 2. PA required to confirm FDA approved indication and to monitor for potential drug-drug interactions.	
ANTINEOPLASTICS- MISCELLANEOUS	MC MC/DEL MC/DEL	AMIFOSTINE MERCAPTOPURINE OXALIPLATIN	MC MC/DEL MC/DEL MC MC/DEL MC/DEL		DOCEFREZ ELOXATIN ETHYOL LEUPROLIDE PURINETHOL ZOLINZA	Use PA Form# 20420	
ANTINEOPLASTICS- MONOCLONAL ANTIBODIES	MC/DEL	TRAZIMERA	MC/DEL MC/DEL MC MC/DEL MC MC MC/DEL		ENHERTU HERCEPTIN HERCESSI HERZUMA KANJINTI OGIVRI ONTRUZANT	Use PA Form# 20420	
CANCER							
CANCER	MC	ALIMTA	MC		ABECMA	Use PA Form# 20420	All non-preferred: A clinical PA is required to confirm appropriate clinical indication for the individual drug request. Specific to each drug all age, clinical testing requirements, previous step therapies, adjunctive drug therapy requirements, and response without disease progression will be also be evaluated for clinical appropriateness. The standard for the appropriate indication will include the FDA label as well as current NCCN guidelines. Scemblix is for the treatment of adult patients with: Philadelphia chromosome-positive chronic myeloid leukemia (Ph+ CML) in chronic phase (CP), previously treated with two or more tyrosine kinase inhibitors (TKIs).
	MC/DEL	ANASTROZOLE TABS	MC		AKEEGA	1. PA required to confirm appropriate diagnosis and testing.	
	MC	ERBITUX	MC		ALECENSA	2. Avoid CYP3A drug interaction.	
	MC	IMATINIB MESYLATE	MC/DEL		ALIQOPA ³	3. Clinical PA required for appropriate diagnosis.	
	MC/DEL	LETROZOLE	MC		ALUNBRIG ¹		
	MC	RUXIENCE	MC		ALYMSYS		
	MC/DEL	VIDAZA	MC/DEL		ARIMIDEX		
	MC	ZIRABEV	MC		AUCATZYL		
			MC		AUGTYRO		
			MC		AVMAPKI-FAKZYNJA		
			MC		AYVAKIT		
			MC/DEL		AVASTIN		
			MC/DEL		BALVERSA		
			MC		BAVENCIO ^{1,8}		
			MC		BEIZRAY		
			MC/DEL		BENDEKA ³		
			MC/DEL		BESPONSA ³		
			MC		BESREMI ¹		
			MC		BIZENGRI		
			MC		BLENREP		
			MC/DEL		BOSULIF		
			MC/DEL		BRAFTOVI ¹		
			MC		BREYANZI		
			MC		BRUKINSA		
			MC		CABOMETYX ³		
			MC		CAMCEVI		
			MC/DEL		CALQUENCE ³		
			MC		COMETRIQ ^{3,4,5}		
			MC		COTELLIC		
			MC/DEL		COPIKTRA		
			MC		DANZITEN		
			MC/DEL		DARZALEX ³		
			MC/DEL		DATROWAY		
			MC/DEL		DAURISMO		
			MC/DEL		ELREXFIO		

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required			Criteria
			MC/DEL		EMPLICITI(IV) ⁸			
			MC		EMRELIS			
			MC		EPKINLY			
			MC/DEL		ERLEADA			
			MC/DEL		ERIVEDGE			
			MC		EXKIVITY			
			MC		FARYDAK			
			MC/DEL		FEMARA			
			MC		FOLOTYN			
			MC		FOTIVDA			
			MC		FRUZAQLA			
			MC		GAVRETO			
			MC/DEL		GILOTrif ^{4,5}			
			MC		GOMEKLI			
			MC		GRAFAPEX			
			MC/DEL		HERNEXEOS			
			MC/DEL		IBRANCE			
			MC		IBTROZI			
			MC		ICLUSIG ³			
			MC/DEL		IDHIFA ³			
			MC		IMBRUVICA			
			MC		IMDELLTRA			
			MC/DEL		IMFINZI			
			MC/DEL		IMJUDO			
			MC		IMKELDI			
			MC		IMLYGIC			
			MC		INLURIYO			
			MC/DEL		INLYTA			
			MC/DEL		INREBIC			
			MC		INQOVI			
			MC		ITOVEBI			
			MC		IWILFIN			
			MC		JAKAFI			
			MC		JAYPIRCA ^{1,2}			
			MC		JEMPERLI			
			MC		JOBEVNE			
			MC/DEL		KEYTRUDA ¹			
			MC		KEYTRUDA QLEX			
			MC		KIMMTRAK			
			MC		KISQALI ¹			
			MC/DEL		KOSELUGO			
			MC		KRAZATI ³			
			MC		KYMRIAH ^{3,9}			
			MC		KYPROLIS ¹			
			MC		LARTRUVO ¹			
			MC		LAZCLUZE			
			MC		LENVIMA			
			MC/DEL		LIBTAYO ¹			
			MC		LONSURF			
			MC/DEL		LORBRENA			
			MC		LOQTORZI			
			MC		LUMAKRAS			
			MC/DEL		LUMOXITI ¹			
			MC		LUNSUMIO ¹			
			MC		LYNOZYFIC			
			MC		LYNPARZA ¹			
			MC		LYTGOBI			
			MC		NEXAVAR ¹			
			MC		NERLYNX ³			

Category	Coverage Indicator	Preferred Drugs	Coverage Indicator	Step Order	Non-Preferred Drugs PA Required			Criteria
			MC		NINLARO(PO)			
			MC/DEL		NUBEQA			
			MC		MARGENZA			
			MC/DEL		MEKINIST ^{3,4}			
			MC/DEL		MEKTOVI ¹			
			MC		MODEYSO			
			MC		MONJUVI			
			MC/DEL		MYLOTARG ³			
			MC/DEL		MVASI			
			MC		ODOMZO ^{1,2,5}			
			MC		OGSIVEO			
			MC		OJEMDA			
			MC		OJJAARA			
			MC		OMISIRGE			
			MC		ONUREG			
			MC/DEL		OPDIVO ³			
			MC		OPDIVO QVANTIG			
			MC		OPDUALAG			
			MC		ORGOVYX			
			MC		ORSERDU ^{2,3}			
			MC		PADCEV			
			MC		PEMAZYRE			
			MC		PEPAXTO			
			MC		PHESGO			
			MC		PHYRAGO			
			MC/DEL		PIQRAY			
			MC		POLIVY			
			MC		POMALYST			
			MC		PORTRAZZA ³			
			MC		QINLOCK			
			MC		RETEVMO			
			MC		REVUFORJ			
			MC/DEL		ROMVIMZA			
			MC		REZLIDHIA			
			MC/DEL		ROZLYTREK			
			MC		RUBRACA			
			MC		RITUXAN			
			MC		RYBREVANT			
			MC		RYDAPT			
			MC		RYLAZE			
			MC		RYTELO			
			MC/DEL		SARCLISA			
			MC		SCEMBLIX ¹			
			MC/DEL		STIVARGA			
			MC/DEL		SUTENT ^{1,2}			
			MC/DEL		SYLATRON			
			MC		TABRECTA			
			MC		TALVEY			
			MC/DEL		TAFINLAR ^{3,4,5,6}			
			MC		TAZVERIK			
			MC/DEL		TALZENNA ¹			
			MC/DEL		TAGRISSO			
			MC		TECARTUS			
			MC		TECELRA			
			MC		TECENTRIQ ¹			
			MC		TECENTRIQ HYBREZA			
			MC		TEPMETKO			
			MC		TEVIMBRA			
			MC/DEL		TIBSOVO ¹			

New drugs are initially non-preferred until reviewed by the DUR Committee and the State. According to State policy, any drug requiring specific diagnosis still requires the specific diagnosis unless otherwise noted within this document.

Last update 1/1/2026

PDL DOSAGE CONSOLIDATION LIST

Tabs/Caps/Patches: Quantities in units

Shaded areas are non-preferred agents - Quantities of these

Sprays/Inhalers/Nebulizers: Quantities in GM, ML, OR MCG

non-preferred agents are available up the limit only with

Injectibles: Quantities in ML

prior authorization

Drug Name	Strength	Limit/Day	Limit/Days	Drug Name	Strength	Limit/Day	Limit/Days
ABILIFY SOLUTION	1MG/ML	30ML	1020/34	ATROVENT HFA	17MCG	12 INHALATIONS	25.8/34
ACCUPRIL	5MG	1	35/35	ATROVENT 30ML	0.03%	12 SPRAYS	30/30
ACCUPRIL	10MG	1	35/35	ATROVENT 15ML	0.06%	16 SPRAYS	45/30
ACCUPRIL	20MG	1	35/35	AVANDIA	2MG	1.5	53/35
ACEON	2MG	1	35/35	AVANDIA	4MG	1	35/35
ACEON	4MG	1	35/35	AVAPRO	75MG	1.5	53/35
ACTONEL	5MG	1	35/35	AVAPRO	150MG	1	35/35
ACTONEL	35MG	1/WK	5/35	AXERT (Step 8)	6.25MG		12/30
ACTOS	All Strengths	1	35/35	AXERT (Step 8)	12.5MG		12/30
ADDERALL XR	5MG	3	90/30	AZELEX	20%		1 TUBE/18
ADDERALL XR	10MG	3	90/30	AZILECT	All Strengths	1	35/35
ADDERALL XR	15MG	3	90/30	BACTROBAN CREAM			1 TUBE/30
ADDERALL XR	20MG	2	60/30	BECONASE AQ	42MCG	8 INHALATIONS	50/30
ADDERALL XR	30MG	1	35/35	BENICAR-HCT	All Strengths	1	30/30
ADEMPAS	All Strengths	1	35/35	BENAZEPRI	5MG	1	35/35
ADVAIR DISKUS	All Strengths	2	60/30	BENAZEPRI	10MG	1.5	53/35
ADVAIR HFA	All Strengths	4	120/30	BENAZEPRI	20MG	1	35/35
ADZENYS XR	All Strengths	1	30/30	BENAZEP/HCTZ	5-6.25	1	35/35
AEROBID	250MCG	8 INHALATIONS	21/35	BENAZEP/HCTZ	10/12.5	1	35/35
AEROBID-M	250MCG	8 INHALATIONS	21/35	BEVESPI AERO		4 INHALATIONS	120/30
ALAVERT-NON DROW	TAB	1	96/96	BONIVA	2.5MG	1	35/35
ALENDRONATE	All Strengths	1/WK	35/35	BOTOX (ADULTS)	100U/ML	1 session/90 days	600U/90
ALTABAX	5GM		1 TUBE/30	BOTOX (CHILDREN>12)	100U/ML	1 session/90 days	400U/90
ALTABAX	15GM		1 TUBE/30	BREO ELLIPTA	100/25MCG	1 INHALATIONS	60/60
ALTABAX	30GM		1 TUBE/30	BRILINTA	All Strengths	2	70/35
ALTACE	1.25MG	1	35/35	BRINTELLIX	All Strengths	1	35/35
ALTACE	2.5MG	1	35/35	BUTRANS		1 patch/WK	4/28
ALTACE	5MG	1	35/35	BYETTA	5mcg inj	0.04ML	1.2ML/30
AMARYL	1MG	1	35/35	BYETTA	10mcg inj	0.08ML	2.4ML/30
AMARYL	2MG	1	35/35	CALAN SR	120MG	1	35/35
AMBIEN	5MG		12/34	CALAN SR	180MG	2	70/35
AMBIEN	10MG		12/34	CALAN SR	240MG	2	70/35
AMBIEN CR	6.25MG		12/34	CARDIZEM CD	120MG/24	1	35/35
AMBIEN CR	12.5MG		12/34	CARDIZEM CD	180MG/24	1	35/35
AMERGE (Step 8)	1MG		12/30	CARDIZEM CD	240MG/24	1	35/35
AMERGE (Step 8)	2.5MG	2.5MG	12/30	CARDIZEM CD	300MG/24	1	35/35
AMLODIPINE	2.5MG	1.5	53/35 DAYS	CARDIZEM CD	360MG/24	1	35/35
AMLODIPINE	5MG	1.5	53/35 DAYS	CARDIZEM LA	120MG/24	1	35/35
AMMONIUM LACTATE CREA	12%		1 TUBE/10	CARDIZEM LA	180MG/24	1	35/35
AMMONIUM LACTATE LOTN	12%		1TUBE/8	CARDIZEM LA	240MG/24	1	35/35
AMPHETAMINE/DEXTROAMPHET ER	5MG	3	90/30	CARDIZEM LA	300MG/24	1	35/35
AMPHETAMINE/DEXTROAMPHET ER	10MG	3	90/30	CARDIZEM LA	360MG/24	1	35/35
AMPHETAMINE/DEXTROAMPHET ER	15MG	3	90/30	CARDURA	1MG	1	35/35
AMPHETAMINE/DEXTROAMPHET ER	20MG	2	60/30	CARDURA	2MG	1.5	53/35
AMPHETAMINE/DEXTROAMPHET ER	30MG	1	90/90	CARDURA	4MG	1.5	53/35
AMPHETAMINE SALT	5,10,15MG	3	105/35	CARTIA XT	120MG	1	90/90
AMPHETAMINE SALT	20MG	2	70/35	CARTIA XT	180MG	1	90/90
AMPHETAMINE SALT	30MG	1	35/35	CARTIA XT	240MG	1	90/90
ANDRODERM	2.5MG	2	60/30	CARTIA XT	300MG	1	90/90
ANDRODERM	5MG	1	30/30	CATAPRES-TTS1	0.1 MG/24HR		5/35
ARAVA	10MG	1	35/35	CATAPRES- TTS2	0.2 MG/24HR		5/35
ARCAPTA	75MCG	1 INHALATION	35/35	CATAPRES- TTS3	0.3 MG/24HR		5/35
ARICEPT	5MG	1	35/35	CEFIXIME	400MG	2	2/7
ARICEPT	10MG	1	35/35	CELEBREX	100MG	1	35/35
ARIPIPRAZOLE	2MG	2	180/90	CELEBREX	200MG	2	70/35
ARIPIPRAZOLE	5MG	2	180/90	CELEBREX	400MG	1	35/35
ARIPIPRAZOLE	10MG	2	180/90	CELEXA	20mg	0.5	17/34
ARIPIPRAZOLE	15MG	2	180/90	CELEXA	40mg	1	51/34
ARIPIPRAZOLE	20MG	1.5	135/90	CITALOPRAM	10MG	2	180/90
ARIPIPRAZOLE	30MG	1	90/90	CITALOPRAM	20MG	2	180/90
ARIIXTRA INJECTION	2.5MG/0.5ML		7/30	CITALOPRAM	40MG	1	90/90
ARIIXTRA INJECTION	5MG/0.4ML		7/30	CLARINEX	REDI TAB	1	35/35
ARIIXTRA INJECTION	7.5MG/0.6ML		7/30	CLEOCIN-T		1 PACKAGE	1/30
ARIIXTRA INJECTION	10MG/0.8ML		7/30	CLINDAMYCIN PHOSPHATE		1 PACKAGE	1/30
ARMONAIR	All Strengths	1 INHALATION	60U/30	COMBIVENT	103-18MCG	12 INHALATIONS	30/35
ASMANEX 30 UNITS	220MCG	1 INHALATION	30U/30	Drug Name	Strength	Limit/Day	Limit/Days
ASMANEX 60 UNITS	220MCG	2 INHALATIONS	60U/30	EFFEXOR XR	37.5MG	1	35/35
ASMANEX 120 UNITS	220MCG	4 INHALATIONS	120U/30	EFFEXOR XR	75MG	1	35/35
ATACAND	4MG	1.5	53/35	EMSAM	All Strengths	1	34/34
ATACAND	8MG	1.5	53/35	ENALAPRIL	2.5	1	90/90

Drug Name	Strength	Limit/Day	Limit/Days	Drug Name	Strength	Limit/Day	Limit/Days
ATACAND	16MG	1	35/35	ENALAPRIL	5MG	1.5	135/90
ATRIPLA	600MG	1	35/35	ENALAPRIL	10MG	1.5	135/90
ATOMOXETINE	All Strengths	1	90/90	ENALAPR/HCTZ	5-12.5	1	90/90
COMETRIQ	80MG	1	35/35	ENBREL	25MG/ML		8/28
COMETRIQ	20MG	3	105/35	ENBREL SURECLICK			8/28
CONCERTA	18MG	1	30/30	ESTAZOLAM	1MG		10/30
CONCERTA	27MG	1	30/30	ESTAZOLAM	2MG		10/30
CONCERTA	36MG	2	60/30	ESTRING MIS	2MG		1/90
COPAXONE INJ	20MG		1/32	EVENITY		12 DOSES/LIFETIME	12 DOSES/LIFETIME
COPAXONE KIT	20MG/ML		1/30	EVOTAZ	All Strengths	1	30/30
COREG CR	All Strengths	1	34/34	FELODIPINE	2.5MG	1	90/90
COSENTYX	150MG	1	1/30	FELODIPINE	5MG	1.5	135/90
CRESTOR	5MG	1	35/35	FENTANYL	25MCG/HR		11/33
CRESTOR	10MG	1	35/35	FENTANYL	50MCG/HR		11/33
CRESTOR	20MG	1	35/35	FENTANYL	75MCG/HR		11/33
CRESTOR	40MG	1	35/35	FENTANYL	100MCG/HR		22/33
CYMBALTA	All Strengths	1	35/35	FETZIMA	All Strengths	1	35/35
DALMANE	15MG		10/30	FINASTERIDE	5MG	1	90/90
DALMANE	30MG		10/30	FLONASE	50MCG	4 SPRAYS	32/34
DAYPRO	600MG	2	70/35	FLOVENT HFA 44MCG	44MCG	4 INHALATIONS	10.6/30
DAYTRANA	10mg/9hr (27.5mg)	1	34/34	FLOVENT HFA 110MCG	110MCG	4 INHALATIONS	12/30
DAYTRANA	15mg/9hr (41.3mg)	1	34/34	FLOVENT HFA 220MCG	220MCG	8 INHALATIONS	24/30
DAYTRANA	20mg/9hr (55.0mg)	1	34/34	FLOVENT DISKUS	50MCG, 100MCG	4 INHALATIONS	60/30
DAYTRANA	30mg/9hr (82.5mg)	1	34/34	FLOVENT DISKUS	250MCG	3 INHALATIONS	120/30
DDAVP	5ML		15/34	FLUCONAZOLE	150MG		1/7
DENAVIR CREAM			2gm/30	FLUNISOLIDE SOLN	0.025%	16 SPRAYS	75/30
DEPO-PROVERA	150MG/ML		1/90	FLUOXETINE CAP	40MG	2	180/90
DEPO-PROVERA	400MG/ML		2.5/90	FLUOXETINE CAP	20MG	4	360/90
DEPO-TESTOSTERONE	200MG/ML		20/90	FLUOXETINE CAP	10MG	3	270/90
DESMOPRESSIN	0.1MG	12	420/35	FLURAZEPAM	15MG		10/30
DESMOPRESSIN	0.2MG	6	210/35	FLURAZEPAM	30MG		10/30
DESONIDE	0.05%		2 TUBES/30	FLUTICASONE SPR		4 SPRAYS	48/90
DESOWEN	0.05%		2 TUBES/30	FLUVOXAMINE	25MG	3	270/90
DETROL LA	2MG	1	35/35	FLUVOXAMINE	50MG	3	270/90
DEXEDRINE	All Strengths	3	90/30	FOCALIN	All Strengths	3	105/35
DEXILANT	All Strengths	1	35/35	FOCALIN XR	All Strengths	1	35/35
DEXTROAMPHETAMINE	All Strengths	3	90/30	FORFIVO XL	All Strengths	1	35/35
DICLOFENAC 1% GEL	1% GEL		2 TUBES/30	FOSAMAX	5MG	1	35/35
DIFLUCAN	150MG		1/7	FOSAMAX	10MG	1	35/35
DILACOR XR	240MG/24	1	35/35	FOSAMAX	70MG	1/WK	5/35
DILACOR XR	120MG/24	1	35/35	FOSAMAX	40MG	2/WK	10/35
DILACOR XR	180MG/24	1	35/35	FOSINOPRIL	10MG	1.5	135/90
DILTIA - XT	120MG/24	1	90/90	FOSINOPRIL	20MG	2	180/90
DILTIA - XT	180MG	1	90/90	FRAGMIN INJ	10000U/ML	2ML	14/7
DILTIA - XT	240MG/24	1	90/90	FRAGMIN INJ	2500U/.2ML	0.4ML	2.80/7
DILTIAZEM CAP ER	120MG	1	90/90	FRAGMIN INJ	25000U/ML	0.8ML	5.6/7
DILTIAZEM CAP XR	120MG	1	90/90	FRAGMIN INJ	5000U/.2ML	0.4ML	2.80/7
DILTIAZEM CAP	120MG/24	1	90/90	FRAGMIN INJ	7500U/.3ML	0.6ML	4.2/7
DILTIAZEM CAP	180MG/24	1	90/90	FROVA TAB (Step 8)	2.5MG		12/30
DILTIAZEM CAP ER	240MG	1	90/90	FULYZAQ	125MG	2	70/35
DILTIAZEM CAP XR	240MG	1	90/90	FUZEON	KIT	1	1/30
DILTIAZEM XR CAP	240MG/24	1	90/90	FYCOMP	All Strengths	1	35/35
DILTIAZEM CAP	240MG/24	1	90/90	GABAPENTIN	300MG	9	810/90
DILTIAZEM CAP	300MG/24	1	90/90	GABAPENTIN	400MG	9	810/90
DILTIAZEM CAP	360MG/24	1	90/90	GABAPENTIN	600MG	6	540/90
DIOVAN	80MG	1	35/35	GABAPENTIN	800MG	4	360/90
DIOVAN - HCT	80 - 12.5	1	35/35	GEODON	20MG	2	70/35
DITROPAN XL	5MG	1	35/35	GEODON	40MG	2	70/35
DITROPAN XL	10MG	2	70/35	GEODON	60MG	2	70/35
DORAL	7.5MG		10/30	GEODON	80MG	2	70/35
DOXAZOSIN	1MG	1	90/90	GEODON	INJ	2	70/35
DOXAZOSIN	2MG	1.5	135/90	GILOTrif	All Strengths	1	35/35
DOXAZOSIN	4MG	1.5	135/90	GLIMEPIRIDE	1MG	1	90/90
DRYSOL SOL	20%		1 BOTTLE/30DAYS	GLIMEPIRIDE	2MG	1	90/90
DURAGESIC PATCHES	12.5MCG/HR		11/33	GLUCOSE TES STRP		12	420/35
DURAGESIC PATCHES	25MCG/HR		11/33	GLUCAGEN INJ. HYPOKIT			2/30
DURAGESIC PATCHES	50MCG/HR		11/33	GLYCOLAX*	255GM		255GM/90
DURAGESIC PATCHES	75MCG/HR		11/33	* Available for once daily dosing to members under the age of 18 years			
DURAGESIC PATCHES	100MCG/HR		22/33	Drug Name	Strength	Limit/Day	Limit/Days
DULOXETINE	20MG	3	270/90	LUNESTA	2MG		12/34
DULOXETINE	30MG	3	270/90	LUNESTA	3MG		12/34
DULOXETINE	60MG	2	180/90	LUPRON DEPOT INJ	11.25MG	KIT	1/90
EDEX	All Strengths		1/30	LUPRON DEPOT INJ	22.5	KIT	1/90
Drug Name	Strength	Limit/Day	Limit/Days				

Drug Name	Strength	Limit/Day	Limit/Days	Drug Name	Strength	Limit/Day	Limit/Days
ILARIS			2/28	LUPRON DEPOT INJ	30MG		1/90
HALCION	0.125MG		10/35	LUPRON DEPOT INJ	30MG	KIT	1/90
HALCION	0.25		10/35	LYRICA	25,50,75MG	3	102/35
HUMIRA	40mg/0.8ml		4/28	LYRICA	100,150,200MG	3	102/35
HYDROXYZINE TAB	All Strengths	3	270/90	LYRICA	225,300MG	2	70/35
HYTRIN	1MG	1	35/35	MAVIK	1MG	1	35/35
HYTRIN	5MG	1	35/35	MAVIK	2MG	1	35/35
HYZAAR	50-12.5	1	35/35	MAXAIR AUTO	200MCG	12 INHALATIONS	14/30
IMDUR	30MG	1.5	53/35	MAXALT (step 8)	5MG		12/30
IMDUR	60MG	1.5	53/35	MAXALT (step 8)	10MG		12/30
IMITREX (step 8)	25MG		12/30	MAXALT MLT (step 1)	5MG		12/30
IMITREX (step 8)	50MG		12/30	MAXALT MLT (step 1)	10MG		12/30
IMITREX (step 8)	100MG		12/30	MEDROXYPR AC	150MG/ML		1/90
IMITREX VIAL	All Strengths		6 boxes/30	MELOXICAM TABS	All Strengths	1	90/90
IMITREX CARTRIDGE	All Strengths		12/30	METADATE ER	10,20MG	3	90/30
IMITREX NASAL SPRAY	All Strengths		12/30	METFORMIN ER	500MG	4	360/90
IMITREX PEN INJCTR	All Strengths		12/30	METHYLINE	All Strengths	3	90/30
IMIQUIMOD	5%		12/28	METHYLPHENIDATE ER	36mg	2	180/90
INTAL	800MCG	8 INHALATIONS	28.4/34	METHYLPHENIDATE	All Strengths	3	90/30
INVOKANA	All Strengths	1	35/35	METROCREAM		1 PACKAGE	1/30
IPRATROPIUM 30ML	0.03%	12 SPRAYS	90/90	METROGEL		1 PACKAGE	1/30
IPRATROPIUM 15ML	0.06%	16 SPRAYS	135/90	METROLOTION		1 PACKAGE	1/30
ISOPTIN SR	180MG	2	70/35	METRONIDAZOLE GEL		1 PACKAGE	1/30
IRBESARTAN	All Strengths	1	90/90	METRONIDAZOLE LOTION		1 PACKAGE	1/30
ISOPTIN SR	240MG	2	70/35	MEVACOR	10MG	1.5	53/35
ISOSORBIDE MONO	30MG	2	180/90	MEVACOR	20MG	1.5	53/35
ISOSORBIDE MONO	60 MG	1.5	135/90	MIACALCIN		3.75ml	1 bottle/34
JANUMET	All Strengths	2	70/35	MICARDIS	All Strengths	1	30/30
JANUVIA	All Strengths	1	35/35	MICARDIS-HCT	All Strengths	1	30/30
JUVISYNC	All Strengths	1	35/35	MIGRANAL NASAL SPRAY	All Strengths		12/30
KETOPROFEN	100MG	2	180/90	MIRALAX	255G	8.5G	1 bottle/30
KETOPROFEN	200MG	1	90/90	MIRALAX	17G/PACKET	0.5 packet	15 packets/30
KETOROLAC	10MG	4.8	24/30	MIRTAZAPINE	15mg	3	270/90
KHEDEZLA	All Strengths	1	35/35	MOBIC	7.5 MG	1	35/35
LAC-HYDRIN CREAM	12%		1TUBE/30	MOBIC	15MG	1	35/35
LAMICTAL	25MG	6	210/35	MOEXIPRIL	7.5	1.5	135/90
LAMICTAL	25MG CHW	6	210/35	MONOPRIL	10MG	1.5	53/35
LAMICTAL	100MG	2	70/35	MONOPRIL	20MG	2	70/35
LAMISIL	250MG	1	35/35	MUPIROCIN			1 TUBE/30
LAMOTRIGINE	25MG	6	540/90	NABUMETONE	500MG	2	180/90
LAMOTRIGINE	100MG	2	180/90	NABUMETONE	750MG	2	180/90
LANSOPRAZOLE CAPS	All Strengths	2	180/90	NARATRIPTAN			12/30
LATUDA	All Strengths	1	17/34	NASACORT AERS	55 MCG	4 SPRAYS	9.3/25
LEFLUNOMIDE	10MG	1	90/90	NASACORT AQ	55MCG	4 SPRAYS	17/30
LEVAQUIN	250MG	1	35/35	NATROBA		120ML	1 bottle/30
LEXAPRO	5MG	0.5	15/30	NAYZILAM	All Strengths		5/30
LIPITOR	10MG	1	35/35	NEUPOGEN INJ	300MCG/ML		10/30
LIPITOR	20MG	1	35/35	NEUPOGEN INJ	480MCG/1.6		16/30
LIPITOR	40MG	1.5	53/35	NEUPOGEN INJ	300MCG/.5ML		5/30
LISINOP/HCTZ	10/12.5MG	1	90/90	NEUPOGEN INJ	480MCG/.8ML		8/30
LINEZOLID	600mg		28/60	NEURONTIN	300MG	9	315/35
LINZESS	All Strengths	1	35/35	NEURONTIN	600MG	9	315/35
LOSARTAN	All Strengths	1	90/90	NEXIUM	20MG	1	35/35
LOSARTAN- HCT	All Strengths	1	90/90	NEXIUM	40MG	2	70/35
LOTENSIN	5MG	1	35/35	NEXIUM SUS	All Strengths	1	30/30
LOTENSIN	10MG	1.5	35/35	NIFEDIPINE CR	90MG	1	90/90
LOTENSIN	20MG	1	53/35	NIFEDIPINE ER	60MG	1	90/90
LOTENSIN - HCT	5 - 6.25	1	35/35	NIFEDIPINE ER	30MG	1	90/90
LOTENSIN - HCT	10 - 12.5	1	35/35	NIFEDIPINE ER	60MG	1	90/90
LOVASTATIN	10MG	1.5	135/90	RELPAX	All Strengths		12/30
LOVASTATIN	20MG	1.5	135/90	REMODULIN	All Strengths		1 MDV/30
LOVENOX INJ	30MG/.3ML	0.6	14 injections/7	RESTORIL	7.5MG		10/30
LOVENOX INJ	40MG/.4ML	0.8	14 injections/7	RESTORIL	15MG		10/30
LOVENOX INJ	60MG/.6ML	1.2	14 injections/7	RESTORIL	30MG		10/30
LOVENOX INJ	80MG/.8ML	1.6	14 injections/7	REVLIMID	All Strengths	1	35/35
LOVENOX INJ	100MG/ML	2	14 injections/7	REYVOW	All Strengths		4/30
LOVENOX INJ	120MG/.8ML	1.6	14 injections/7	RHINOCORT AQ	32MCG	8 SPRAYS	18/30
LOVENOX INJ	150MG/ML	2	14 injections/7	REFRESH PLUS		15 ML	1 bottle/30
LUNESTA	1MG		12/34	REFRESH PLUS		30 ML	2 bottles/30
NIFEDIPINE ER	90MG	1	90/90	REFRESH TEARS		15 ML	1 bottle/30
NIFEDIPINE ER,CR	30MG	1	90/90	REFRESH TEARS		30 ML	2 bottles/30
NORVASC	2.5MG	1.5	53/35 DAYS	RESCULA			2 bottles/35
NORVASC	5MG	1.5	53/35 DAYS	REYATAZ	All Strengths	1	35/35
NURTEC ODT	All Strengths		8/30	RISPERDAL	0.5MG	1.5	53/35

Drug Name	Strength	Limit/Day	Limit/Days	Drug Name	Strength	Limit/Day	Limit/Days
NUVARING		1/MO	1/28	RISPERDAL	0.25MG	1.5	53/35
ODOMZO	200mg	1	30/30	RISPERDAL	1MG	1.5	53/35
OLMESARTAN	All Strengths	1	90/90	RISPERDAL	2MG	1.5	53/35
OLANZAPINE	2.5MG	3	270/90	RISPERDAL	3MG	2	70/35
OLANZAPINE	5MG	3	270/90	RISPERDAL	4MG	2	70/35
OLANZAPINE	7.5MG	3	270/90	RISPERDAL INJ	25MG		2/28
OLANZAPINE	10MG	3	270/90	RISPERDAL INJ	37.5		2/28
OLANZAPINE	15MH	2	180/90	RISPERDAL INJ	50MG		2/28
OLANZAPINE	20MG	1.5	135/90	RISPERDAL M-TAB	0.5MG	1.5	53/35
OLANZAPINE ODT	All Strengths	1	90/90	RISPERDAL M-TAB	1MG	1.5	53/35
OMEPRAZOLE	10MG	2	180/90	RISPERDAL M-TAB	2MG	4	140/35
OMEPRAZOLE	20MG	2	180/90	RISPERDAL SOL.	1MG/ML	8ML	280/35
OMEPRAZOLE	40MG	2	180/90	RISPERIDONE	0.5MG	3	270/90
OMNARIS	50MCG	4 sprays	12.5/30	RISPERIDONE	0.25MG	3	270/90
OPSUMIT	All Strengths	1	35/35	RISPERIDONE	1MG	3	270/90
ORUVAIL	100MG	2	70/35	RISPERIDONE	2MG	3	270/90
ORUVAIL	200MG	1	35/35	RISPERIDONE	3MG	2	180/90
OXAPROZIN	600MG	2	180/90	RISPERIDONE	4MG	2	180/90
OXYCODONE ER	10,20,40MG	2	70/35	RISPERIDONE SOL.	1MG/ML	8ML	280/35
OXYCODONE ER	80MG	4	140/35	RITALIN LA	All Strengths	1	35/35
OXYCONTIN**	10,20,30,40MG	2	70/35	RITALIN LA	30mg	2	70/35
OXYCONTIN**	80MG	4	140/35	SAVELLA	All Strengths	2	70/35
PANTOPRAZOLE	All Strengths	2	180/90	SEREVENT DISKUS	50MCG	2 INHALATIONS	60/30
PAROXETINE	10MG	2	180/90	SEROQUEL	100MG		45/30
PAROXETINE	20MG	2	180/90	SEROQUEL XR	150MG	1	35/35
PAXIL	10MG	1.5	53/35	SEROQUEL XR	200MG	1	35/35
PAXIL	20MG	1	35/35	SEROQUEL XR	300MG	2	70/35
PEGASYS KIT		KIT	1/28	SEROQUEL XR	400MG	2	70/35
PLAN B			2/15 or 4/30	SERTRALINE	25MG	3	270/90
PLENDIL	2.5MG	1	35/35	SERTRALINE	50MG	3	270/90
PLENDIL	5MG	1.5	53/35	SERTRALINE	100MG	3	270/90
PRAVACHOL	10MG	1	35/35	SIMVASTATIN	5MG	1	35/35
PRAVACHOL	20MG	1	35/35	SIMVASTATIN	10MG	1.5	53/35
PRAVACHOL	40MG	1	35/35	SIMVASTATIN	20MG	1.5	53/35
PRAVACHOL	80MG	1	35/35	SIMVASTATIN	40MG	1.5	53/35
PRAVASTATIN	10MG	1	35/35	SIMVASTATIN	80MG	1	35/35
PRAVASTATIN	20MG	1	35/35	SINGULAIR	4MG	1	35/35
PRAVASTATIN	40MG	2	180/90	SINGULAIR	5MG	1	35/35
PRAVASTATIN	80MG	1	35/35	SINGULAIR	10MG	1	35/35
PREVPAC MIS	500MG-30MG		14/30	SONATA	5MG		12/34
PRILOSEC OTC	20MG	2	168/84	SONATA	10MG		12/34
PRINIVIL	2.5MG	1	35/35	SPIRIVA	HANDIHLR	1 INHALTION	30/30
PRINIVIL	5MG	1	35/35	SPORANOX SOL	10MG/ML	10ML/ML	300cc/30
PRINIVIL	10MG	1.5	53/35	SPORANOX PULSEPAK	F		30/30
PRINIVIL	20MG	1.5	53/35	SPORANOX	100MG		30/30
PRINZIDE	10-12.5	1	35/35	STADOL INJ	1MG/ML		9/35
PROAIR HFA	90mcg	12 INHALATIONS	17/34	STADOL INJ	2MG/ML		9/35
PROTONIX	20MG	2	70/35	STRATTERA	All Strengths	1	35/35
PROTONIX	40MG	2	70/35	SUPRAX	400MG	1	1/7
PROZAC	10MG	1.5	53/35	XIFYRM	All Strengths	1	90/90
PULMICORT	200MCG	8 INHALATIONS	1/25	XOPENEX HFA		12 INHALATIONS	2 INHALERS/34
PULMICORT FLEX	All Strengths	8 Inhalations	2/30	XOPENEX NEB		12CC	408/34
QUETIAPINE	25MG	3	270/90	ZALEPLON	All Strengths		30/30
QUETIAPINE	50MG	3	270/90	ZECURITY	6.5		4/28
QUETIAPINE	100MG	3	270/90	ZEMBRACE	All Strengths		3boxes/30
QUETIAPINE	200MG	3	270/90	ZESTORETIC	10-12.5	1	35/35
QUINAPRIL	5MG	1	90/90	ZESTRIL	2.5MG	1	35/35
QUINAPRIL	10MG	1	90/90	ZESTRIL	5MG	1	35/35
QUINAPRIL	20MG	1	90/90	ZESTRIL	10MG	1.5	53/35
QVAR AERS	All Strengths	8 Inhalations	14.6/25	ZESTRIL	20MG	1.5	53/35
RANITIDINE SYRUP***	15MG/ML	20ML	700ML/35	ZETONNA	37MCG	2	60/30
RELAFEN	500MG	2	70/35	ZIPRASIDONE	20MG	3	270/90
RELAFEN	750MG	2	70/35	ZIPRASIDONE	40MG	3	270/90
REMERON	15MG	1.5	53/35	ZOCOR	5MG	1	35/35
SULAR	10MG	1.5	53/35	ZOCOR	10MG	1.5	53/35
SULAR	20MG	1	35/35	ZOCOR	20MG	1.5	53/35
SUMATRIPTAN PEN INJ	All Strengths		12/30	ZOCOR	40MG	1.5	53/35
SUMATRIPTAN NASAL SPRAY	All Strengths		12/30	ZOFRAN*	4MG	3	90/30
SUMATRIPTAN SYRINGE	All Strengths		12/30	ZOFRAN*	8MG	1.5	45/30
SUMATRIPTAN TAB	All Strengths		12/30	ZOFRAN*	24MG	0.5	15/30
SYNViSC INJ	8MG/ML		2/30	ZOFRAN*	4MG/5ML	15ML	450/30
SYRINGES		10	1000/100	ZOLMITRIPTAN TAB	All Strengths		12/30
TAFINLAR	50MG	6	210/35	ZOLOFT	25MG	0.5	18/35
TAFINLAR	75MG	4	140/35	ZOLOFT	50MG	0.5	18/35

Drug Name	Strength	Limit/Day	Limit/Days
TAMIFLU CAPS	75MG		10/30
TAZTIA XT CAP	120MG/24	1	90/90
TAZTIA XT CAP	180MG/24	1	90/90
TAZTIA XT CAP	240MG/24	1	90/90
TAZTIA XT CAP	300MG/24	1	90/90
TAZTIA XT CAP	360MG/24	1	90/90
TELMISARTAN	All Strengths	1	90/90
TEMAZEPAM	7.5MG		10/30
TEMAZEPAM	15MG		10/30
TEMAZEPAM	30MG		10/30
TEQUIN	200MG	1	35/35
TERAZOSIN	1MG	1	90/90
TERAZOSIN	5MG	1	90/90
TERBINAFINE	250MG	1	35/35
TEST STRIPS	Blood Glucose	12	420/35
TAZAC	120MG/24	1	35/35
TAZAC	180MG/24	1	35/35
TAZAC	240MG/24	1	35/35
TAZAC	300MG/24	1	35/35
TAZAC	360MG/24	1	35/35
TAZAC	420MG/24	1	35/35
TILADE	1.75MG	8 INHALATIONS	48.6/35
TOPAMAX SPRINKLES	All Strengths	1	35/35
TOPROL XL	25MG	1.5	53/35
TOPROL XL	50MG	1.5	53/35
TRADJENTA	All Strengths	1	35/35
TRAMADOL	50MG	8	720/90
TRAMADOL/ APAP	37.5/325MG	8	720/90
TRELEGY ELLIPTA	All Strengths	1INHALATION	30U/30
TREXIMET	85/500	2.5	12/30
TRIAZOLAM	0.125MG		10/30
TRIAZOLAM	0.25MG		10/30
TROKENDI XR	25MG	1	35/35
TROKENDI XR	50MG	1	35/35
TROKENDI XR	100MG	1	35/35
TROKENDI XR	200MG	2	70/35
UBRELVY	All Strengths		10/30
ULTRAM	50MG	8	280/35
UNIVASC	7.5MG	1.5	53/35 DAYS
UTIBRON	7.5mcg/15.6mcg	2 INHALATIONS	60/30
VALTOCO	All Strengths		10/30
VALSARTAN-HCT	All Strengths	1	90/90
VASERETIC	5-12.5MG	1	35/35
VASOTEC	2.5MG	1	35/35
VASOTEC	5MG	1.5	53/35
VASOTEC	10MG	1.5	53/35
VENLAFAXINE TABS	25	3	270/90
VENLAFAXINE TABS	37.5	3	270/90
VENLAFAXINE TABS	100	3	270/90
VENLAFAXINE ER CAPS	37.5	3	270/90
VENLAFAXINE ER CAPS	75	3	270/90
VENLAFAXINE ER	150	2	180/90
VENTOLIN HFA	90MCG	12 INHALATIONS	36/34
VERAPAMIL ER, SR	120MG	1	90/90
VERAPAMIL ER, CR, SR	180MG	2	90/90
VERAPAMIL ER, CR, SR	240MG	2	90/90
VERELAN	180MG	1	35/35
VERELAN SR	120MG	1	35/35
VERELAN SR	180MG	1	35/35
VERELAN SR	240MG	2	70/35
VERAMYST	27.5MCG	4 sprays	10/30
VYEPTI	All Strengths		4/30
VYVANSE	All Strengths	1	35/35
VYVANSE CHEW	All Strengths	1	35/35

Drug Name	Strength	Limit/Day	Limit/Days
ZOLOFT	100MG	3	105/35
ZOLPIDEM (step 1)	5MG		30/30
ZOLPIDEM (step 1)	10MG		30/30
ZOMIG (Step 8)	5MG		12/30
ZTLIDO	All Strengths	3	90/30
ZYPREXA	2.5MG	1.5	53/35
ZYPREXA	5MG	1	35/35
ZYPREXA	7.5MG	1	35/35
ZYPREXA	10MG	1	35/35
ZYPREXA	15MG	1	35/35
ZYPREXA	20MG	1	35/35
ZYPREXA ZYDIS	5MG	1	35/35
ZYPREXA ZYDIS	10MG	1	35/35
ZYPREXA ZYDIS	15MG	1	35/35
ZYPREXA ZYDIS	20MG	1	35/35

*Cancer diagnosis with non-daily chemotherapy required

**Available without pa with CA and HO diag.

*** Ranitidine syrup available without PA to users less than 6 years old.

MDV=Multidose Vial

Pain Management Policy

[Back to Index](#)

Beginning January 2017, all current opiate users who are above the maximum combined daily dose of 100 MME must titrate their total daily dose of opioid medications below 30 MME. Also, the maximum daily supply of an opiate prescription for acute pain will be limited to 7-day supplies. The maximum day supply of an opiate prescription for chronic pain will be limited to 30-day supplies. As of July 1, 2017 all users of opioid medications must comply with the maximum combined daily dose of 100 MME.

However, for MaineCare members, effective January 1, 2017, opioid prescription(s) for more than a 7-day supply and/or more than 30 MME/ day will require a prior authorization. Please note that MaineCare implemented a 30 MME limit January 1, 2013 that is still effective.

The following are general exceptions: pain associated with cancer treatment, end-of-life and hospice care, palliative care, and symptoms related to HIV/AIDS. Per MaineCare criteria, the diagnosis of cancer must be written on the prescription. A palliative care exception for any MaineCare opioid prescription will require prior authorization (PA) with appropriate clinical documentation.

Post-surgical members may receive prior authorizations for opiates up to a 60 days in length if medical necessity is provided by the surgical provider.

An MME conversion chart is available at www.maineicarepdl.org. Click on "General Pharmacy Info."

Updated July 1, 2025